

Inventors: Peng Cho Tang, G. Davis Harris and Xiaoyuan Li

**INDOLINONE DERIVATIVES AS PROTEIN KINASE/PHOSPHATASE
INHIBITORS**

5

CROSS-REFERENCE

This application claims priority under 35 U.S.C. 119(e) to U.S. Provisional
Applications Serial No. 60/209,162, filed June 2, 2000, the disclosure of which is
10 incorporated herein by reference in its entirety.

BACKGROUND OF THE INVENTION

Field of Invention

The present invention relates to certain 2-indolinone compounds which
15 modulate the activity of protein kinases ("PKs") and phosphatases. The compounds
of this invention are therefore useful in treating disorders related to abnormal PK
activity. Pharmaceutical compositions comprising these compounds, methods of
treating diseases utilizing pharmaceutical compositions comprising these compounds
and methods of preparing them are also disclosed.

20 State of the Art

Cellular signal transduction is a fundamental mechanism whereby
extracellular stimuli are relayed to the interior of cells and subsequently regulate
diverse cellular processes. One of the key biochemical mechanisms of signal
transduction involves the reversible phosphorylation of proteins. Phosphorylation of
25 polypeptides regulates the activity of mature proteins by altering their structure and
function. Phosphate most often resides on the hydroxyl moiety (-OH) of serine,
threonine, or tyrosine amino acids in proteins.

Enzymes that mediate phosphorylation of cellular effectors generally fall into
two classes. The first class consists of protein kinases which transfer a phosphate
30 moiety from adenosine triphosphate to protein substrates. The second class consists
of protein phosphatases which hydrolyze phosphate moieties from phosphoryl
protein substrates. The converse functions of protein kinases and protein

phosphatases balance and regulate the flow of signals in signal transduction processes.

Protein kinases and protein phosphatases are generally divided into two groups - receptor and non-receptor type proteins. Most receptor-type protein tyrosine phosphatases contain two conserved catalytic domains, each of which encompasses a segment of 240 amino acid residues. Saito *et al.*, 1991, *Cell Growth and Diff.* 2:59-65. Receptor protein tyrosine phosphatases can be subclassified further based upon the amino acid sequence diversity of their extracellular domains. Saito *et al.*, supra; Krueger *et al.*, 1992, *Proc. Natl. Acad. Sci. USA* 89:7417-7421.

Protein kinases and protein phosphatases are also typically divided into three classes based upon the amino acids they act upon. Some catalyze the addition or hydrolysis of phosphate on serine or threonine only, some catalyze the addition or hydrolysis of phosphate on tyrosine only, and some catalyze the addition or hydrolysis of phosphate on serine, threonine, and tyrosine.

Tyrosine kinases can regulate the catalytic activity of other protein kinases involved in cell proliferation. Protein kinases with inappropriate activity are also involved in some types of cancer. Abnormally elevated levels of cell proliferation are associated with receptor and non-receptor protein kinases with unregulated activity.

In addition to their role in cellular proliferation, protein kinases are thought to be involved in cellular differentiation processes. Cell differentiation occurs in some cells upon nerve growth factor (NGF) or epidermal growth factor (EGF) stimulation. Cellular differentiation is characterized by rapid membrane ruffling, cell flattening, and increases in cell adhesion. Chao, 1992, *Cell* 68:995-997.

In an effort to discover novel treatments for cancer and other diseases, biomedical researchers and chemists have designed, synthesized, and tested molecules that inhibit the function of protein kinases. Some small organic molecules form a class of compounds that modulate the function of protein kinases. Examples of molecules that have been reported to inhibit the function of protein kinases are bis-monocyclic, bicyclic or heterocyclic aryl compounds (PCT WO 92/20642), vinylene-azaindole derivatives (PCT WO 94/14808), 1-cyclopropyl-4-


pyridyl-quinolones (U.S. Patent No. 5,330,992), styryl compounds (by Levitzki, et al., U.S. Patent No. 5,217,999, and entitled "Styryl Compounds which Inhibit EGF Receptor Protein Tyrosine Kinase), styryl-substituted pyridyl compounds (U.S. Patent No. 5,302,606), certain quinazoline derivatives (EP Application No. 0 566 266 A1), seleoindoles and selenides (PCT WO 94/03427), tricyclic polyhydroxylic compounds (PCT WO 92/21660), and benzylphosphonic acid compounds (PCT WO 91/15495).

The compounds that can traverse cell membranes and are resistant to acid hydrolysis are potentially advantageous therapeutics as they can become highly bioavailable after being administered orally to patients. However, many of these protein kinase inhibitors only weakly inhibit the function of protein kinases. In addition, many inhibit a variety of protein kinases and will therefore cause multiple side-effects as therapeutics for diseases.

Despite the significant progress that has been made in developing compounds for the treatment of cancer, there remains a need in the art to identify the particular structures and substitution patterns that form the compounds capable of modulating the function of particular protein kinases.

SUMMARY OF THE INVENTION

In a first aspect, the invention provides an indolinone compound having a structure set forth in formula (I):



wherein:

- (a) R_4 - R_6 , and R_8 - R_{10} are hydrogen;
- (b) R_1 , R_2 , and R_3 are each independently selected from the group consisting of hydrogen, halogen, carboxylic acid, optionally substituted ester, optionally

substituted amide, optionally substituted alkyl, optionally substituted alkoxy, trihalomethyl, optionally substituted aryl, and optionally substituted heteroaryl; and

- (c) R_7 is selected from the group consisting of substituted alkyl, and substituted alkoxy;

or a pharmaceutically acceptable salt thereof.


Preferably,

- (a) R_1 is selected from the group consisting of hydrogen and optionally substituted alkyl, more preferably hydrogen;

- (b) R_2 and R_3 are each independently selected from the group consisting of hydrogen, halo, carboxylic acid, optionally substituted heteroaryl, and optionally substituted phenyl. Preferably R_2 is hydrogen, halo, phenyl, or carboxylic acid, more preferably hydrogen, phenyl, -COOH, chloro, fluoro or bromo. Preferably, R_3 is hydrogen, halo, carboxylic acid, optionally substituted pyridyl, and phenyl optionally substituted with lower alkoxy or halo, more preferably phenyl, -COOH, pyridin-3-yl, 3-, or 4-methoxyphenyl or 4-fluorophenyl; and


- (c) R_7 is selected from the group consisting of lower alkyl substituted with a heteroaliphatic ring or dialkylamino; or lower alkoxy substituted with a heteroaliphatic ring or dialkylamino, preferably R_7 is lower alkyl substituted with a heteroaliphatic ring or dialkylamino; more preferably R^7 is 3-diethylaminopropyl or 3-pyrrolidin-1-yl-propyl, 2-dimethylaminoethoxy, 2-diethylaminoethoxy, 2-pyrrolidin-1-yl-ethoxy, or 2-morpholin-4-yl-ethoxy.

In a second aspect, the invention provides for an indolinone compound having a structure set forth in formula (II):



wherein:

- (a) R_{11} - R_{14} are hydrogen;
- (b) R_{15} and R_{16} are each independently selected from the group consisting of hydrogen, optionally substituted alkyl, optionally substituted aryl, and optionally substituted heteroaryl, or R_{15} and R_{16} taken together with the nitrogen atom to which they are attached form a ring structure selected from the group consisting of a five-membered or six-membered heteroaromatic ring, a five-membered or six-membered heteroaliphatic ring, a nine-membered fused bicyclic heteroaromatic ring, and a ten-membered fused bicyclic heteroaromatic ring; and
- (c) A is selected from the group consisting of formula (III), (IV), and (V):



wherein:

- (i) R_{19} - R_{25} and R_{27} - R_{31} are hydrogen;
- (ii) R_{17} and R_{18} are each independently selected from the group consisting of hydrogen, optionally substituted alkyl, and optionally substituted alkoxy provided that both R_{17} and R_{18} are not hydrogen; and
- (iii) R_{26} is selected from the group consisting of optionally substituted alkyl; or a pharmaceutically acceptable salt thereof.
- In one preferred embodiment:
- (i) R_{15} is hydrogen or alkyl, more preferably hydrogen or methyl;
- (ii) R_{16} is hydrogen, alkyl, phenyl optionally substituted with one or two substituents selected from halo or unsubstituted lower alkyl or 5 or 6 membered heteroaryl; more preferably hydrogen, methyl, isopropyl, phenyl, pyridin-3-yl, 3-chlorophenyl, or 4-chloro-2-fluorophenyl; or

(iii) R_{15} and R_{16} together with the nitrogen to which they are attached form 2,3-dihydroindol-1-yl, 2,3-dihydro-2H-quinolin-1-yl, or 2,3-dihydro-2H-isoisoquinolin-2-yl ring wherein said rings are optionally substituted with halo or alkyl, preferably R_{15} and R_{16} together with the nitrogen atom to which they are attached form 2,3-dihydroindol-1-yl, 2,3-dihydro-2H-quinolin-1-yl, 5-bromo-2,3-dihydro-2H-quinolin-1-yl, or 2,3-dihydro-2H-isoisoquinolin-2-yl;

(iv) A is group of formula III where:

R_{17} is hydrogen, methyl, or methoxy, preferably hydrogen; and

R_{18} is selected from the group consisting of lower alkoxy substituted with heteroalicyclic, preferably 2-pyrrolidin-1-yl-ethoxy and 2-morpholin-4-yl-ethoxy.

Another preferred group of compound is that wherein:

(i) R_{15} is hydrogen or alkyl, preferably hydrogen or methyl;

(ii) R_{16} is hydrogen, alkyl, phenyl optionally substituted with one or two substituents selected from halo or unsubstituted lower alkyl, or 5 or 6 membered heteroaryl; preferably hydrogen, methyl, isopropyl, phenyl, pyridin-3-yl, 3-chlorophenyl, or 4-chloro-2-fluorophenyl; or

R_{15} and R_{16} together with the nitrogen to which they are attached form 2,3-dihydroindol-1-yl, 2,3-dihydro-2H-quinolin-1-yl, or 2,3-dihydro-2H-isoisoquinolin-2-yl ring wherein said rings are optionally substituted with halo or alkyl, preferably 3-dihydroindol-1-yl, 2,3-dihydro-2H-quinolin-1-yl, 5-bromo-2,3-dihydro-2H-quinolin-1-yl, or 2,3-dihydro-2H-isoisoquinolin-2-yl; and

(iii) A is group of formula IV where R_{26} is selected from the group consisting of optionally substituted alkyl, preferably hydrogen or methyl.

Another preferred group of compounds is that wherein:

- (i) R_{15} is hydrogen or alkyl, preferably hydrogen or methyl;
- (iii) R_{16} is hydrogen, alkyl, phenyl optionally substituted with one or two substituents selected from halo or unsubstituted lower alkyl or 5 or 6 membered heteroaryl; more preferably hydrogen, methyl, isopropyl, phenyl, pyridin-3-yl, 3-chlorophenyl, or 4-chloro-2-fluorophenyl; or
- R_{15} and R_{16} together with the nitrogen to which they are attached form 2,3-dihydroindol-1-yl, 2,3-dihydro-2H-quinolin-1-yl, or 2,3-dihydro-2H-isoquinolin-2-yl ring wherein said rings are optionally substituted with halo or alkyl, preferably R_{15} and R_{16} together with the nitrogen atom to which they are attached form 2,3-dihydroindol-1-yl, 2,3-dihydro-2H-quinolin-1-yl, 5-bromo-2,3-dihydro-2H-quinolin-1-yl, or 2,3-dihydro-2H-isoisoquinolin-2-yl;

- (ii) A is group of formula V.

As used herein, the term “optionally substituted alkyl” refers to an aliphatic hydrocarbon group. The alkyl moiety may be a “saturated alkyl” group, which means that it does not contain any alkene or alkyne moieties. The alkyl moiety may also be an “unsaturated alkyl” moiety, which means that it contains at least one alkene or alkyne moiety. An “alkene” moiety refers to a group consisting of at least two carbon atoms and at least one carbon-carbon double bond, and an “alkyne” moiety refers to a group consisting of at least two carbon atoms and at least one carbon-carbon triple bond. The alkyl moiety, whether saturated or unsaturated, may be branched, non-branched, or cyclic. Preferably, the alkyl group has 1 to 20 carbon atoms (whenever it appears herein, a numerical range such as “1 to 20” refers to each integer in the given range; *e.g.*, “1 to 20 carbon atoms” means that the alkyl group may consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, etc., up to and including 20 carbon atoms, although the present definition also covers the occurrence of the term “alkyl” where no numerical range is designated). More preferably, it is a medium size alkyl having 1 to 10 carbon atoms. Most preferably, it is a lower alkyl having 1 to 4 carbon atoms. The alkyl group is optionally substituted with one, two, or three substituents individually and independently selected from cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, alkoxy, aryloxy, mercapto, alkylthio, arylthio, cyano, or halo. The term “substituted alkyl” means

that the alkyl group as defined above carries at least one of the substituents listed above.

The term "optionally substituted aryl" refer to an aromatic carbocyclic group of 6 to 12 ring atoms which has at least one ring having a conjugated π electron system (e.g., phenyl, naphthyl, tetrahydronaphthyl, and the like) which is optionally substituted with one, two, or three substituents independently selected from optionally substituted alkyl, halogen, trihalomethyl, hydroxy, alkoxy, carboxyl amino, amido, nitro, and ester.

"Optionally substituted phenyl" refers to a phenyl group that is optionally substituted with one, two, or three substituents independently selected from optionally substituted alkyl, halogen, trihalomethyl, hydroxy, alkoxy, carboxyl amino, amido, nitro, and ester.

The term "cycloalkyl" refers to a saturated, cyclic group of 3 to 6 carbon atoms (e.g., cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl) which is optionally substituted with one, two, or three substituents independently selected from optionally substituted alkyl, halogen, trihalomethyl, hydroxy, alkoxy, carboxyl amino, amido, nitro, and ester.

The term "optionally substituted heteroaryl or heteroaromatic ring" refers to a ring system of 5 to 10 ring atoms in which one, two, three, or four of the atoms forming the backbone is a heteroatom selected from nitrogen, oxygen, or sulfur, the remaining being carbon e.g., pyridine, furan, pyrrole, indole, pyrazine, pyrimidine, tetrazole, imidazole, and the like. The heteroaromatic/heteroaryl ring may be single or fused bicyclic ring and wherein one of the rings may be partially or fully saturated e.g., 2,3-dihydroindole, 2,3-dihydroquinoline, 2,3-dihydroisoquinoline, and the like. The heteroaryl/heteroaromatic ring is optionally substituted with one, two, or three substituents independently selected from optionally substituted alkyl, halogen, trihalomethyl, hydroxy, alkoxy, carboxyl amino, amido, nitro, and ester.

The term "optionally substituted heteroaliphatic or heteroalicyclic" refers to a saturated ring system of 5 to 9 ring atoms in which in which one, two, three, or four of the atoms forming the backbone is a heteroatom selected from nitrogen, oxygen, or sulfur, the remaining being carbon e.g., piperazine, piperidine, pyrrolidine,

morpholine, tetrahydrofuran and the like. The heteroaliphatic ring is optionally substituted with one, two, or three substituents independently selected from optionally substituted alkyl, halogen, trihalomethyl, hydroxy, alkoxy, carboxyl amino, amido, nitro, and ester.

5 The term "halogen" refers to an atom selected from the group consisting of fluorine, chlorine, bromine, and iodine.

The term "trihalomethyl" refers to the $-C(X)_3$ group, where X is a halogen group as defined above e.g., trifluoromethyl, trichloromethyl, tribromomethyl, and the like.

10 The term "optionally substituted alkoxy" is a group of formula -O-alkyl wherein alkyl is as defined above. The term "substituted alkoxy" means that the alkyl group as defined above carries at least one of the substituents listed above. The term "alkoxy" means that the alkyl chain in the alkoxy group as defined above is not substituted.

15 When X is hydrogen, then the alkoxy group becomes a "hydroxy" group, *i.e.*, -OH.

A "nitro" is a substituent of formula $-NO_2$.

The term "alkylthio" refers to -SR group where R is unsubstituted alkyl as defined above e.g., methylthio, ethylthio, and the like.

20 The term "aryloxy" refers to -OR group where R is aryl group as defined above, e.g., phenoxy, and the like.

The term "arylthio" refers to -SR group where R is aryl group as defined above e.g., phenylthio, and the like.

25 The term "dialkylamino" means -NRR where each R is an unsubstituted alkyl group of 1-6 carbon atoms e.g., dimethylamino, diethylamino, and the like.

The term "optionally substituted ester" refers to -COOR where R is an alkyl group as defined above.

The term "optionally substituted amide" refers to $-CONR^aR^b$ where R^a and R^b are hydrogen or unsubstituted lower alkyl.

30 "Pharmaceutically acceptable salt" refers to those salts which retain the biological effectiveness and properties of the free bases and which are obtained by

reaction with inorganic or organic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, malic acid, citric acid, maleic acid, succinic acid, tartaric acid, and the like.


- 5 Some of the preferred compounds of the invention that have the generic structure of formula I are listed in Table 1.

Table 1

Compound Number	Compound Name
IN-001	3-[5-(3-diethylamino-propyl)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-002	5-bromo-3-[5-(3-diethylamino-propyl)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-003	3-[5-(3-diethylamino-propyl)-1 <i>H</i> -indol-2-ylmethylene]-6-phenyl-1,3-dihydro-indol-2-one
IN-004	3-[5-(3-diethylamino-propyl)-1 <i>H</i> -indol-2-ylmethylene]-5-phenyl-1,3-dihydro-indol-2-one
IN-005	3-[5-(2-dimethylamino-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-5-phenyl-1,3-dihydro-indol-2-one
IN-006	5-phenyl-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-007	3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-008	5-bromo-3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-009	3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-6-phenyl-1,3-dihydro-indol-2-one
IN-010	3-[5-(2-dimethylamino-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-011	5-bromo-3-[5-(2-dimethylamino-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-012	3-[5-(2-dimethylamino-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-6-phenyl-1,3-dihydro-indol-2-one
IN-013	3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-014	5-bromo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-015	6-phenyl-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-016	3-[5-(2-diethylamino-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-017	5-bromo-3-[5-(2-diethylamino-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-018	3-[5-(2-diethylamino-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-6-phenyl-1,3-dihydro-indol-2-one
	3-[5-(2-diethylamino-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-5-phenyl-1,3-dihydro-indol-2-one

Compound Number	Compound Name
IN-019	one
IN-020	3-[5-(3-pyrrolidin-1-yl-propyl)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-021	2-oxo-3-[5-(3-pyrrolidin-1-yl-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indol-5-carboxylic acid
IN-022	2-oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-5-carboxylic acid
IN-023	2-oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-6-carboxylic acid
IN-024	4-(2-hydroxy-ethyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-025	6-pyridin-3-yl-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-026	6-(4-methoxy-phenyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-027	6-(3-methoxy-phenyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-028	6-(2-methoxy-phenyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-029	6-(4-fluoro-phenyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-030	3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-carboxylic acid
IN-031	3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-6-carboxylic acid
IN-032	3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-5-phenyl-1,3-dihydro-indol-2-one
IN-033	4-(2-hydroxy-ethyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-034	3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-6-pyridin-3-yl-1,3-dihydro-indol-2-one
IN-035	6-(4-methoxy-phenyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-036	6-(3-methoxy-phenyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-037	6-(2-methoxy-phenyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-038	6-(4-fluoro-phenyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one

The above compounds have the structure of formula X, with the R₁₀₁, R₁₀₂, R₁₀₃, and R₁₁₂ substituents as defined in Table 2.



038602-060401

Table 2

Compound Number	R ₁₀₁	R ₁₀₂	R ₁₀₃	R ₁₁₂
IN-001	H	H	3-diethylamino-propyl	H
IN-002	Br	H	3-diethylamino-propyl	H
IN-003	H	Phenyl	3-diethylamino-propyl	H
IN-004	Phenyl	H	3-diethylamino-propyl	H
IN-005	Phenyl	H	2-dimethylamino-ethoxy	H
IN-006	Phenyl	H	2-pyrrolidin-1-yl-ethoxy	H
IN-007	H	H	2-morpholin-4-yl-ethoxy	H
IN-008	Br	H	2-morpholin-4-yl-ethoxy	H
IN-009	H	Phenyl	2-morpholin-4-yl-ethoxy	H
IN-010	H	H	2-dimethylamino-ethoxy	H
IN-011	Br	H	2-dimethylamino-ethoxy	H
IN-012	H	Phenyl	2-dimethylamino-ethoxy	H
IN-013	H	H	2-pyrrolidin-1-yl-ethoxy	H
IN-014	Br	H	2-pyrrolidin-1-yl-ethoxy	H
IN-015	H	Phenyl	2-pyrrolidin-1-yl-ethoxy	H
IN-016	H	H	2-diethylamino-ethoxy	H
IN-017	Br	H	2-diethylamino-	H

Compound Number	R ₁₀₁	R ₁₀₂	R ₁₀₃	R ₁₁₂
			ethoxy	
IN-018	H	Phenyl	2-diethylamino-ethoxy	H
IN-019	Phenyl	H	2-diethylamino-ethoxy	H
IN-020	H	H	3-pyrrolidin-1-yl-propyl	H
IN-021	-COOH	H	3-pyrrolidin-1-yl-propyl	H
IN-022	-COOH	H	2-pyrrolidin-1-yl-ethoxy	H
IN-023	H	-COOH	2-pyrrolidin-1-yl-ethoxy	H
IN-024	H	H	2-pyrrolidin-1-yl-ethoxy	2-hydroxy-ethyl
IN-025	H	Pyridin-3-yl	2-pyrrolidin-1-yl-ethoxy	H
IN-026	H	4-Methoxy Phenyl	2-pyrrolidin-1-yl-ethoxy	H
IN-027	H	3-Methoxy Phenyl	2-pyrrolidin-1-yl-ethoxy	H
IN-028	H	2-Methoxy Phenyl	2-pyrrolidin-1-yl-ethoxy	H
IN-029	H	4-Fluoro Phenyl	2-pyrrolidin-1-yl-ethoxy	H
IN-030	-COOH	H	2-morpholin-4-yl-ethoxy	H
IN-031	H	-COOH	2-morpholin-4-yl-ethoxy	H
IN-032	Phenyl	H	2-morpholin-4-yl-ethoxy	H
IN-033	H	H	2-morpholin-4-yl-ethoxy	2-hydroxy-ethyl
IN-034	H	Pyridin-3-yl	2-morpholin-4-yl-	H

Compound Number	R ₁₀₁	R ₁₀₂	R ₁₀₃	R ₁₁₂
			ethoxy	
IN-035	H	4-Methoxy Phenyl	2-morpholin-4-yl-ethoxy	H
IN-036	H	3-Methoxy Phenyl	2-morpholin-4-yl-ethoxy	H
IN-037	H	2-Methoxy Phenyl	2-morpholin-4-yl-ethoxy	H
IN-038	H	4-Fluoro Phenyl	2-morpholin-4-yl-ethoxy	H

Some of the compounds of the invention that have the generic structure of formula II are listed in Table 3.

09874002860

Table 3

Compound Number	Compound Name
IN-039	2-oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid amide
IN-040	3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid amide
IN-041	2-oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid methylamide
IN-042	2-oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid dimethylamide
IN-043	2-oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid isopropylamide
IN-044	2-oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid phenylamide
IN-045	2-oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid pyridin-3-ylamide
IN-046	5-(2,3-dihydro-indole-1-sulfonyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-047	2-oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid (3-chloro-phenyl)-amide
IN-048	2-oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid (3-chloro-phenyl)-methyl-amide
IN-049	2-oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid (4-chloro-2-fluoro-phenyl)-amide
IN-050	5-(3,4-dihydro-2 <i>H</i> -quinoline-1-sulfonyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-051	5-(3,4-dihydro-1 <i>H</i> -isoquinoline-2-sulfonyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-052	5-(5-bromo-2,3-dihydro-indole-1-sulfonyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-053	3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid methylamide
IN-054	3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid dimethylamide
IN-055	3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid isopropylamide
IN-056	3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid phenylamide
IN-057	3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid pyridin-3-ylamide
IN-058	5-(2,3-dihydro-indole-1-sulfonyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-059	3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid (3-chloro-phenyl)-amide
IN-060	3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid (3-chloro-phenyl)-methyl-amide
IN-061	3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid (4-chloro-2-fluoro-phenyl)-amide
IN-062	5-(3,4-dihydro-2 <i>H</i> -quinoline-1-sulfonyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
	5-(3,4-dihydro-1 <i>H</i> -isoquinoline-2-sulfonyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-

Compound Number	Compound Name
IN-063	2-ylmethylene]-1,3-dihydro-indol-2-one
IN-064	5-(5-bromo-2,3-dihydro-indole-1-sulfonyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
IN-065	3-(1 <i>H</i> -indol-3-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid amide
IN-066	3-(2-methyl-1 <i>H</i> -indol-3-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid amide
IN-067	3-(1 <i>H</i> -indol-5-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid amide
IN-068	3-(1 <i>H</i> -indol-3-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid methylamide
IN-069	3-(2-methyl-1 <i>H</i> -indol-3-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid methylamide
IN-070	3-(1 <i>H</i> -indol-5-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid methylamide
IN-071	3-(1 <i>H</i> -indol-2-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid methylamide
IN-072	3-(1 <i>H</i> -indol-3-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid dimethylamide
IN-073	3-(2-methyl-1 <i>H</i> -indol-3-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid dimethylamide
IN-074	3-(1 <i>H</i> -indol-5-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid dimethylamide
IN-075	3-(1 <i>H</i> -indol-2-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid dimethylamide
IN-076	3-(4-methoxy-1 <i>H</i> -indol-2-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid methylamide

Some of the above compounds have the structure of formula XI, with the R_{104} , R_{105} , R_{106} , and R_{113} substituents as defined in Table 4, while others have the structure of formula XII, with the R_{107} , R_{108} , and R_{109} substituents as defined in Table 5, and still others have the structure of formula XIII, with the R_{110} and R_{111} substituents as defined in Table 6.





Table 4

Compound Number	R ₁₀₄	R ₁₀₅	R ₁₀₆	R ₁₁₃
IN-039	H	H	2-pyrrolidin-1-yl-ethoxy	H
IN-040	H	H	2-morpholin-4-yl-ethoxy	H
IN-041	H	CH ₃	2-pyrrolidin-1-yl-ethoxy	H
IN-042	CH ₃	CH ₃	2-pyrrolidin-1-yl-ethoxy	H
IN-043	H	Isopropyl	2-pyrrolidin-1-yl-ethoxy	H
IN-044	H	Phenyl	2-pyrrolidin-1-yl-ethoxy	H
IN-045	H	Pyridin-3-yl	2-pyrrolidin-1-yl-ethoxy	H
IN-046	R ₁₀₄ and R ₁₀₅ taken together form a 2,3-dihydroindol-1-yl ring		2-pyrrolidin-1-yl-ethoxy	H
IN-047	H	3-Chloro-phenyl	2-pyrrolidin-1-yl-ethoxy	H
IN-048	CH ₃	3-Chloro-phenyl	2-pyrrolidin-1-yl-ethoxy	H
IN-049	H	4-Chloro-2-fluorophenyl	2-pyrrolidin-1-yl-ethoxy	H
IN-050	R ₁₀₄ and R ₁₀₅ taken together form a 3,4-dihydro-2H-quinolin-1-yl ring		2-pyrrolidin-1-yl-ethoxy	H
IN-051	R ₁₀₄ and R ₁₀₅ taken together form a 3,4-dihydro-2H-isoquinolin-2-yl ring		2-pyrrolidin-1-yl-ethoxy	H
IN-052	R ₁₀₄ and R ₁₀₅ taken together form a 5-bromo-2,3-dihydroindol-1-yl ring		2-pyrrolidin-1-yl-ethoxy	H

Compound Number	R ₁₀₄	R ₁₀₅	R ₁₀₆	R ₁₁₃
IN-053	H	CH ₃	2-morpholin-4-yl-ethoxy	H
IN-054	CH ₃	CH ₃	2-morpholin-4-yl-ethoxy	H
IN-055	H	Isopropyl	2-morpholin-4-yl-ethoxy	H
IN-056	H	Phenyl	2-morpholin-4-yl-ethoxy	H
IN-057	H	Pyridin-3-yl	2-morpholin-4-yl-ethoxy	H
IN-058	R ₁₀₄ and R ₁₀₅ taken together form a 2,3-dihydroindol-1-yl ring		2-morpholin-4-yl-ethoxy	H
IN-059	H	3-Chloro-phenyl	2-morpholin-4-yl-ethoxy	H
IN-060	CH ₃	3-Chloro-phenyl	2-morpholin-4-yl-ethoxy	H
IN-061	H	4-Chloro-2-fluorophenyl	2-morpholin-4-yl-ethoxy	H
IN-062	R ₁₀₄ and R ₁₀₅ taken together form a 3,4-dihydro-2 <i>H</i> -quinolin-1-yl ring		2-morpholin-4-yl-ethoxy	H
IN-063	R ₁₀₄ and R ₁₀₅ taken together form a 3,4-dihydro-2 <i>H</i> -isoquinolin-2-yl ring		2-morpholin-4-yl-ethoxy	H
IN-064	R ₁₀₄ and R ₁₀₅ taken together form a 5-bromo-2,3-dihydroindol-1-yl ring		2-morpholin-4-yl-ethoxy	H
IN-071	H	CH ₃	H	H
IN-075	CH ₃	CH ₃	H	H
IN-076	H	CH ₃	H	Methoxy




Compound Number	R ₁₀₇	R ₁₀₈	R ₁₀₉
IN-065	H	H	H
IN-066	H	H	CH ₃
IN-068	CH ₃	H	H
IN-069	CH ₃	H	CH ₃
IN-072	CH ₃	CH ₃	H
IN-073	CH ₃	CH ₃	CH ₃



Compound Number	R ₁₁₀	R ₁₁₁
IN-067	H	H
IN-070	CH ₃	H
IN-074	CH ₃	CH ₃

In a further aspect, the invention relates to an indolinone compound having a structure set forth in formula VI:













5 where


- (a) R_{41} is selected from the group consisting of hydrogen, amide, and sulfonamide;
- (b) R_{42} and R_{44} are each independently selected from the group consisting of hydrogen, halogen, and alkoxy;
- 10 (c) R_{43} is selected from the group consisting of hydrogen, alkyl, halogen, hydroxy, alkoxy, perhaloalkoxy, nitro, sulfone, and sulfonamide;
- (d) R_{45} is selected from the group consisting of hydrogen, alkyl, nitro, and amide; and
- (e) R_{46} is selected from the group consisting of hydrogen, alkyl, carboxylic acid, and amine.

15 Preferrably, for the compounds of formula VI,


- (a) R_{42} and R_{44} are each independently selected from the group consisting of hydrogen, chloro, and methoxy;
- (b) R_{43} is selected from the group consisting of, hydrogen, fluoro, chloro, bromo, methoxy, hydroxy, methyl, ethyl, nitro, trifluoromethoxy, $-\text{SO}_2\text{CH}_3$, and $-\text{NHSO}_2\text{CH}_3$,
- 20 (c) R_{45} is selected from the group consisting of hydrogen, methyl, $-\text{NO}_2$, and $-\text{NHC}(\text{O})\text{C}(\text{CH}_3)_3$, and
- (d) R_{46} is selected from the group consisting of hydrogen, methyl, $-\text{CH}_2\text{N}(\text{CH}_3)_2$, $-\text{CH}_2\text{N}(\text{CH}_2\text{CH}_3)_2$, $-\text{CH}_2\text{CH}_2\text{C}(\text{O})\text{OH}$, and $-\text{CH}_2\text{CH}_2\text{CH}_2\text{C}(\text{O})\text{OH}$.

25 The preferred compounds of the invention that have the structure set forth in formula VI are selected from the group consisting of







, (12)




,




, (15)




,




, (18)




,



, (21)




,




,



(24)





,




(22)



(24)





and (26)



5

where R_{41} is selected from the group consisting of $-\text{SO}_2\text{N}(\text{X}_1)_2$, $-\text{NHSO}_2\text{X}_1$, $-\text{NHC}(\text{O})\text{X}_1$, where X_1 is hydrogen or alkyl.


In another aspect, the invention relates to a combinatorial library of at least five indolinone compounds that can be formed by reacting an oxindole with an aldehyde where the oxindole has a structure set forth in formula VII or VIII:



5 where

- (a) R_{51} - R_{54} , R_{64} , and R_{65} are hydrogen;
- (b) R_{55} and R_{56} are each independently selected from the group consisting of hydrogen, optionally substituted alkyl, optionally substituted aryl, and optionally substituted heteroaryl, or when taken together R_{55} and R_{56} form an optionally substituted five-membered or six-membered heteroaliphatic ring;
- (c) R_{61} - R_{63} are each independently selected from the group consisting of hydrogen, halogen, carboxylic acid, optionally substituted aryl, optionally substituted heteroaryl, and amide; and

where the aldehyde has a structure set forth in formula IX



15 where

- (d) R_{71} and R_{73} - R_{75} are hydrogen;
- (e) R_{72} is selected from the group consisting of hydrogen, optionally substituted alkyl, and optionally substituted alkoxy.

20 A "combinatorial library" refers to all the compounds formed by the reaction of each compound of one dimension with a compound in each of the other dimensions in a multi-dimensional array of compounds. In the context of the present invention, the array is two dimensional and one dimension represents all the oxindoles of the invention and the second dimension represents all the aldehydes of the invention. Each oxindole may be reacted with each and every aldehyde in order to form an indolinone compound. All indolinone compounds formed in this way are

within the scope of the present invention. Also within the scope of the present invention are smaller combinatorial libraries formed by the reaction of some of the oxindoles with all of the aldehydes, all of the oxindoles with some of the aldehydes, or some of the oxindoles with some of the aldehydes.

5 Preferably in the combinatorial library of the invention:

(a) R_{55} and R_{56} are each independently selected from the group consisting of hydrogen, methyl, isopropyl, 2-methoxyethyl, benzyl, 4-fluorobenzyl, 2-methoxyphenyl, 3-fluorophenyl, 4-fluorophenyl, 3-chlorophenyl, 3-pyridyl, or when taken together R_{55} and R_{56} form a ring selected from the group consisting of pyrrolidine, 4-methylpiperazine;

(b) R_{61} - R_{63} are each independently selected from the group consisting of hydrogen, bromo, phenyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 3-pyridyl, -COOH, -C(O)NHCH₂CH₃, -C(O)NHCH₂C(O)OH, -C(O)NHCH(CH₃)C(O)OH,

15 -C(O)NHCH(CH(CH₃)₂)C(O)OH, and  ;

(c) R_{72} is selected from the group consisting of hydrogen 2-diethylamino-ethoxy, 3-diethylamino-1-yl-propyl, and 3-pyrrolidin-1-yl-propyl.

Even more preferably, in the combinatorial library of the invention the aldehyde is selected from the group consisting of 2-formyl-1*H*-indole, 2-formyl-5-(3-diethylamino-propyl)-1*H*-indole, 2-formyl-5-(2-dimethylaminoethoxy)-1*H*-indole, 2-formyl-5-(2-dimethylaminethoxy)-1*H*-indole, 2-formyl-5-(2-morpholin-4-yl-ethoxy)-1*H*-indole, 2-formyl-5-(2-pyrrolidine-1-yl-ethoxy)-1*H*-indole, 2-methyl-1*H*-indole-3-carbaldehyde, 1*H*-indole-5-carbaldehyd, and 2-formyl-5-(3-pyrrolidin-1-yl-propyl)-1*H*-indol;

25 and the oxindole is selected from the group consisting of 2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid amide, 2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid methylamide, 2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid dimethylamide, 2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid (4-chloro-2-fluoro-phenyl)-ethyl-amide, 5-(2,3-dihydro-indole-1-sulfonyl)-1,3-dihydro-indol-2-one, 2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid (4-chloro-2-fluoro-phenyl)-methyl-amide, 2-oxo-2,3-dihydro-

- 1H-indole-5-sulfonic acid (4-chloro-2-fluoro-phenyl)-amide, 5-phenyl-1,3-dihydro-indol-2-one, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid [2-(4-methoxy-phenyl)-2-oxo-ethyl]-amide, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid (3-methoxy-phenyl)-amide, N-(2-oxo-2,3-dihydro-1H-indol-5-yl)-benzamide,
- 5 cyclopentanecarboxylic acid (2-oxo-2,3-dihydro-1H-indol-5-yl)-amide, (2-oxo-2,3-dihydro-1H-indol-5-yl)-carbamic acid benzyl ester, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid 4-fluoro-benzylamide, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid [3-(2-oxo-pyrrolidin-1-yl)-propyl]-amide, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid (1,2,3,4-tetrahydro-naphthalen-1-yl)-amide, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid (furan-2-ylmethyl)-amide, 3-(2-oxo-2,3-dihydro-1H-indole-5-sulfonylamino)-thiophene-2-carboxylic acid methyl ester, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid (3-chloro-phenyl)-amide, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid m-tolylamide, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid (2-hydroxy-ethyl)-amide, 4-(2-oxo-2,3-dihydro-1H-indole-5-sulfonyl)-piperazine-1-carboxylic acid ethyl
- 15 ester, 4-(2-oxo-2,3-dihydro-1H-indole-5-sulfonyl)-piperazine-1-carbaldehyde, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid (2-methoxy-ethyl)-amide, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid propylamide, 5-(4-methyl-piperazine-1-sulfonyl)-1,3-dihydro-indol-2-one, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid benzylamide, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid (2-methoxy-phenyl)-amide, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid cyclopropylamide, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid pyridin-3-ylamide, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid phenylamide, 5-(pyrrolidine-1-sulfonyl)-1,3-dihydro-indol-2-one, 5-(4-acetyl-piperazine-1-sulfonyl)-1,3-dihydro-indol-2-one, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid cyclohexylamide, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid (2-
- 25 morpholin-4-yl-ethyl)-amide, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid cyclobutylamide, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid (1-phenyl-ethyl)-amide, 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid cyclopentylamide, 5-(2-pyrrolidin-1-yl-acetyl)-1,3-dihydro-indol-2-one, N-(2-oxo-2,3-dihydro-1H-indole-5-yl)-acetamide, (2-oxo-2,3-dihydro-1H-indole-5-yl)-carbamic acid tert-butyl ester, 2-
- 30 oxo-2,3-dihydro-1H-indole-5-sulfonic acid isopropylamide, toluene-4-sulfonic acid 2-(2-oxo-2,3-dihydro-1H-indole-4-yl)-ethyl ester, 2-oxo-2,3-dihydro-1H-indole-4-

carboxylic acid ethylamide, 4-phenyl-1,3-dihydro-indol-2-one, 2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid (4-methoxy-phenyl)-amide, 4-(2-morpholin-4-yl-ethyl)-1,3-dihydro-indol-2-one, 4-(2-pyrrolidin-1-yl-ethyl)-1,3-dihydro-indol-2-one, 4-[2-(4-methyl-piperazin-1-yl)-ethyl]-1,3-dihydro indol-2-one, 4-[2-(3-bromo-phenoxy)-ethyl]-1,3-dihydro-indol-2-one, 4-(2-bromo-ethyl)-1,3-dihydro-indol-2-one, 4-(2-bromo-ethyl)-1,3-dihydro-indol-2-one, 2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid (6-methoxy-4'-methylsulfanyl-biphenyl-3-yl)-amide, 2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid (4-methoxy-3-thiophen-3-yl-phenyl)-amide, 2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid biphenyl-3-ylamide, 2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid (6-methoxy-3'-trifluoromethyl-biphenyl-3-yl)-amide, 2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid (4'-fluoro-6-methoxy-biphenyl-3-yl)-amide, 2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid (6-methoxy-[1,1',4',1'']terphenyl-3-yl)-amide, 2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid (6,4'-dimethoxy-biphenyl-3-yl)-amide, 2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid (3-chloro-4-methoxy-phenyl)-amide, isopropyl-carbamic acid 2-(2-oxo-2,3-dihydro-1H-indol-4-yl)-ethyl ester, 4-[2-(3-trifluoromethyl-phenoxy)-ethyl]-1,3-dihydro-indol-2-one, 4-[2-(5-chloro-pyridin-3-yloxy)-ethyl]-1,3-dihydro-indol-2-one, 4-[2-(4-methoxy-phenoxy)-ethyl]-1,3-dihydro-indol-2-one, 4-(2-ethoxy-ethyl)-1,3-dihydro-indol-2-one, 4-(2-methoxy-ethyl)-1,3-dihydro-indol-2-one, ethyl-carbamic acid 2-(2-oxo-2,3-dihydro-1H-indol-4-yl)-ethyl ester, biphenyl-2-yl-carbamic acid 2-(2-oxo-2,3-dihydro-1H-indol-4-yl)-ethyl ester, 6-pyridin-3-yl-1,3-dihydro-indol-2-one, 6-phenyl-1,3-dihydro-indol-2-one, 6-(2-methoxy-phenyl)-1,3-dihydro-indol-2-one, 6-(3-methoxy-phenyl)-1,3-dihydro-indol-2-one, 6-(4-methoxy-phenyl)-1,3-dihydro-indol-2-one, and cyclohexyl-carbamic acid 2-(2-oxo-2,3-dihydro-1H-indol-4-yl)-ethyl ester.

Some of the compounds of the invention that can be formed by the above comibinatorial library are listed in Table 7.

Table 7

Compound Number	Compound Name
CL-001	3-[5-(3-diethylamino-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid dimethyl amide
CL-002	3-[5-(3-diethylamino-propyl)-1 <i>H</i> -indol-2-ylmethylene]-5-(4-methyl-piperazine-1-sulfonyl)-1,3-dihydro-indol-2-one
CL-003	3-[5-(3-diethylamino-propyl)-1 <i>H</i> -indol-2-ylmethylene]-5-(pyrrolidine-1-sulfonyl)-1,3-dihydro-indol-2-one
CL-004	3-[5-(3-diethylamino-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid (2-methoxy-ethyl)-amide
CL-005	3-[5-(3-diethylamino-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid pyridin-3-ylamide
CL-006	3-[5-(3-diethylamino-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid (2-methoxy-phenyl)amide
CL-007	3-[5-(3-diethylamino-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid (3-chlorophenyl)amide
CL-008	3-[5-(3-diethylamino-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid (4-fluorobenzyl)amide
CL-009	3-[5-(3-diethylamino-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid amide
CL-010	3-[5-(3-diethylamino-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid isopropylamide
CL-011	3-(1 <i>H</i> -indol-2-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid benzylamide
CL-012	3-(1 <i>H</i> -indol-2-ylmethylene)-5-(pyrrolidine-1-sulfonyl)-1,3-dihydro-indol-2-one
CL-013	3-(1 <i>H</i> -indol-2-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid (2-methoxyethyl)amide
CL-014	3-(1 <i>H</i> -indol-2-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid (2-methoxyphenyl)amide
CL-015	3-(1 <i>H</i> -indol-2-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid (3-chlorophenyl)amide
CL-016	3-(1 <i>H</i> -indol-2-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid (4-fluorobenzyl)amide
CL-017	3-(1 <i>H</i> -indol-2-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid amide
CL-018	3-(1 <i>H</i> -indol-2-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid methylamide
CL-019	3-(1 <i>H</i> -indol-2-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid dimethylamide
CL-020	3-(1 <i>H</i> -indol-2-ylmethylene)-2-oxo-2,3-dihydro-1 <i>H</i> -indole-5-sulfonic acid isopropylamide
CL-021	5-bromo-3-[5-(2-diethylaminoethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
CL-022	3-[5-(2-diethylaminoethoxy)-1 <i>H</i> -indol-2-ylmethylene]-6-(3-methoxyphenyl)-1,3-dihydro-indol-2-one
CL-023	3-[5-(2-diethylaminoethoxy)-1 <i>H</i> -indol-2-ylmethylene]-6-(4-methoxyphenyl)-1,3-dihydro-indol-2-one
CL-024	3-[5-(2-diethylaminoethoxy)-1 <i>H</i> -indol-2-ylmethylene]-6-phenyl-1,3-dihydro-indol-2-one
	3-[5-(2-diethylaminoethoxy)-1 <i>H</i> -indol-2-ylmethylene]-6-(2-methoxyphenyl)-1,3-

Compound Number	Compound Name
CL-025	dihydro-indol-2-one
CL-026	3-[5-(2-diethylaminoethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-4-carboxylic acid ethylamine
CL-027	6-bromo-3-[5-(2-diethylaminoethoxy)-1 <i>H</i> -indol-2-ylmethylene]-1,3-dihydro-indol-2-one
CL-028	3-[5-(2-diethylaminoethoxy)-1 <i>H</i> -indol-2-ylmethylene]-6-pyridin-1,3-dihydro-indol-2-one
CL-029	3-[5-(2-diethylaminoethoxy)-1 <i>H</i> -indol-2-ylmethylene]-5-phenyl-1,3-dihydro-indol-2-one
CL-030	3-[5-(2-diethylaminoethoxy)-1 <i>H</i> -indol-2-ylmethylene]-2-oxo-2,3-dihydro-1 <i>H</i> -indole-4-carboxylic acid (3-chloro-4-methoxyphenyl)-amide
CL-031	2-oxo-3-[5-(3-pyrrolidin-1-yl-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-5-carboxylic acid
CL-032	2-oxo-3-[5-(3-pyrrolidin-1-yl-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-6-carboxylic acid
CL-033	({2-oxo-3-[5-(3-pyrrolidin-1-yl-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-5-carbonyl}-amino)-acetic acid
CL-034	2-({2-oxo-3-[5-(3-pyrrolidin-1-yl-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-6-carbonyl}-amino)-acetic acid
CL-035	({2-oxo-3-[5-(3-pyrrolidin-1-yl-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-5-carbonyl}-amino)-propionic acid
CL-036	3-methyl-2-({2-oxo-3-[5-(3-pyrrolidin-1-yl-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-5-carbonyl}-amino)-butyric acid
CL-037	2-({2-oxo-3-[5-(3-pyrrolidin-1-yl-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-6-carbonyl}-amino)-propionic acid
CL-038	3-methyl-2-({2-oxo-3-[5-(3-pyrrolidin-1-yl-propyl)-1 <i>H</i> -indol-2-ylmethylene]-2,3-dihydro-1 <i>H</i> -indole-6-carbonyl}-amino)-butyric acid

Some of the above compounds have the structure of formula XIV, with the R_{114} , R_{115} , and R_{116} substituents as defined in Table 8, while others have the structure of formula XV, with the R_{107} , R_{108} , and R_{109} substituents as defined in Table 9.

5







Table 8


Compound Number	R ₁₁₄	R ₁₁₅	R ₁₁₆
CL-001	CH ₃	CH ₃	3-diethylamino-propyl
CL-002	R ₁₁₄ and R ₁₁₅ together form 4-methyl-piperazin-1-yl		3-diethylamino-propyl
CL-003	R ₁₁₄ and R ₁₁₅ together form pyrrolidin-1-yl		3-diethylamino-propyl
CL-004	2-methoxyethyl	H	3-diethylamino-propyl
CL-005	pyridin-3-yl	H	3-diethylamino-propyl
CL-006	2-methoxyphenyl	H	3-diethylamino-propyl
CL-007	3-chlorophenyl	H	3-diethylamino-propyl
CL-008	4-fluorobenzyl	H	3-diethylamino-propyl
CL-009	H	H	3-diethylamino-propyl
CL-010	isopropyl	H	3-diethylamino-propyl
CL-011	benzyl	H	H
CL-012	R ₁₁₄ and R ₁₁₅ together form pyrrolidin-1-yl		H
CL-013	2-methoxyethyl	H	H
CL-014	2-methoxyphenyl	H	H
CL-015	3-chlorophenyl	H	H
CL-016	4-fluorobenzyl	H	H
CL-017	H	H	H
CL-018	CH ₃	H	H
CL-019	CH ₃	CH ₃	H
CL-020	isopropyl	H	H



**Table 9**

Compound Number	R ₁₁₇	R ₁₁₈	R ₁₁₉	R ₁₂₀
CL-021	H	Br	H	2-Diethylaminoethoxy
CL-022	H	H	3-Methoxyphenyl	2-Diethylaminoethoxy
CL-023	H	H	4-Methoxyphenyl	2-Diethylaminoethoxy
CL-024	H	H	Phenyl	2-Diethylaminoethoxy
CL-025	H	H	2-Methoxyphenyl	2-Diethylaminoethoxy
CL-026		H	H	2-Diethylaminoethoxy
CL-027	H	H	Br	2-Diethylaminoethoxy
CL-028	H	H	Pyridin-3-yl	2-Diethylaminoethoxy
CL-029	H	Phenyl	H	2-Diethylaminoethoxy
CL-030		H	H	2-Diethylaminoethoxy
CL-031	H	-COOH	H	3-Pyrrolidin-1-yl-propyl


Compound Number	R ₁₁₇	R ₁₁₈	R ₁₁₉	R ₁₂₀
CL-032	H	H	-COOH	3-Pyrrolidin-1-yl-propyl
CL-033	H		H	3-Pyrrolidin-1-yl-propyl
CL-034	H	H		3-Pyrrolidin-1-yl-propyl
CL-035	H		H	3-Pyrrolidin-1-yl-propyl
CL-036	H			3-Pyrrolidin-1-yl-propyl
CL-037	H	H		3-Pyrrolidin-1-yl-propyl
CL-038	H	H		3-Pyrrolidin-1-yl-propyl

In another aspect, the invention relates to a method of synthesizing a compound of formula I, II, or VI, which method comprises reacting an oxindole having the structure set forth in formula XV, XVI, or XVII, with an aldehyde or a ketone, having a structure set forth in formula XVIII, XIX, or XX. R₁-R₁₆, R₄₁-R₄₆, and A in formulae XV-XX are as defined herein.





5



R' is hydrogen or alkyl, as defined herein.

10

piperidine. In other embodiments the base may be the hydroxide anion, preferably used as its sodium or potassium salt.

The synthesis of the compounds of the invention takes place in a solvent. The solvent of the reaction is preferably a protic solvent or an aprotic solvent.

- 5 “Protic solvents” are those that are capable of donating a proton to a solute. Examples of protic solvents include, but are not limited to, alcohols and water. “Aprotic solvents” are those solvents that, under normal reaction conditions, do not donate a proton to a solute. Typical organic solvents, such as hexane, toluene, benzene, methylene chloride, dimethylformamide, dimethylsulfoxide, chloroform, 10 tetrahydrofuran, are some of the examples of aprotic solvents. Other aprotic solvents are also within the scope of use by the present invention. In some preferred embodiments, the solvent of the reaction is an alcohol, which may preferably be isopropanol or most preferably ethanol. Water is another preferred protic solvent. Dimethylformamide, known in the chemistry art as DMF, and dimethylsulfoxide, 15 known in the chemistry art as DMSO, are preferred aprotic solvents.

- The synthetic method of the invention calls for the reaction to take place at elevated temperatures which are temperatures that are greater than room temperature. More preferably, the elevated temperature is preferably about 30-150 °C, more preferably is about 70-100 °C, and most preferably is about 70-90 °C, 20 which is about the temperature at which ethanol boils (*i.e.*, the boiling point of ethanol). By “about” a certain temperature it is meant that the temperature range is preferably within 10 °C of the listed temperature, more preferably within 5 °C of the listed temperature, and most preferably within 2 °C of the listed temperature. Therefore, by way of example, by “about 80 °C” it is meant that the temperature 25 range is preferably 80±10 °C, more preferably 80±5 °C, and most preferably 80±2 °C.

- The synthetic method of the invention may be accompanied by the step of screening a library for a compound of the desired activity and structure - thus, providing a method of synthesis of a compound by first screening for a compound 30 having the desired properties and then chemically synthesizing that compound.

In another aspect, the invention features a pharmaceutical composition comprising (i) a physiologically acceptable carrier, diluent, or excipient; and (ii) an indolinone compound of the invention, as described herein. It is understood that the indolinone compound of the invention may be one of formula I, II, or VI, or a compound obtained from the combinatorial library of the invention.

The term "pharmaceutical composition" refers to a mixture of an indolinone compound of the invention with other pharmaceutically acceptable diluents, excipients, or carriers. The pharmaceutical composition facilitates administration of the compound to an organism. Pharmaceutical compositions can be obtained by reacting compounds with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, methanesulfonic acid, ethanesulfonic acid, *p*-toluenesulfonic acid, salicylic acid and the like.

The term "pharmaceutically acceptable" defines a carrier or diluent that does not abrogate the biological activity and properties of the compound.

The term "carrier" defines a chemical compound that facilitates the incorporation of a compound into cells or tissues. For example dimethyl sulfoxide (DMSO) is a commonly utilized carrier as it facilitates the uptake of many organic compounds into the cells or tissues of an organism.

The term "diluent" defines chemical compounds diluted in water that will dissolve the compound of interest as well as stabilize the biologically active form of the compound. Salts dissolved in buffered solutions are utilized as diluents in the art. One commonly used buffered solution is phosphate buffered saline because it mimics the salt conditions of human blood. Since buffer salts can control the pH of a solution at low concentrations, a buffered diluent rarely modifies the biological activity of a compound.

The invention also features a method of modulating the function of a PK or a protein phosphatase (PP) with an indolinone compound of the invention, comprising the step of contacting cells expressing the PK or a PP with the compound. It is understood that the indolinone compound of the invention may be one of formula I, II, or VI, or a compound obtained from the combinatorial library of the invention.

The term "function" refers to the cellular role of PKs or PPs. These proteins include members that regulate many steps in signaling cascades, including cascades controlling cell growth, migration, differentiation, gene expression, muscle contraction, glucose metabolism, cellular protein synthesis, and regulation of the cell cycle.

The term "catalytic activity", in the context of the invention, defines the rate at which a protein kinase phosphorylates a substrate or a protein phosphatase dephosphorylates a substrate. Catalytic activity can be measured, for example, by determining the amount of a substrate converted to a product as a function of time.

Phosphorylation or dephosphatase of a substrate occurs at the active-site of a protein kinase. The active-site is normally a cavity in which the substrate binds to the protein kinase or phosphatase and is connected to the product.

The term "substrate" as used herein refers to a molecule phosphorylated by a PK or dephosphorylated by a PP. The substrate is preferably a peptide and more preferably a protein.

The term "activates" refers to increasing the cellular function of a PK or PP. This function is preferably the interaction with a natural binding partner and most preferably catalytic activity.

The term "inhibit" refers to decreasing the cellular function of a PK or PP. The PK or PP function is preferably the interaction with a natural binding partner and most preferably catalytic activity.

The term "modulates" refers to altering the function of a protein kinase by increasing or decreasing the probability that a complex forms between a protein kinase and a natural binding partner. A modulator preferably increases the probability that such a complex forms between the PK or PP and the natural binding partner, more preferably increases or decreases the probability that a complex forms between the PK or PP and the natural binding partner depending on the concentration of the compound exposed to the PK or PP, and most preferably decreases the probability that a complex forms between the PK or PP and the natural binding partner. A modulator preferably activates the catalytic activity of a PK or PP, more preferably activates or inhibits the catalytic activity of a PK or PP

depending on the concentration of the compound exposed to the PK or PP, or most preferably inhibits the catalytic activity of a PK or PP.

The term “complex” refers to an assembly of at least two molecules bound to one another. Signal transduction complexes often contain at least two protein
5 molecules bound to one another.

The term “natural binding partner” refers to polypeptides that bind to a PK or PP in cells. Natural binding partners can play a role in propagating a signal in a PK or PP signal transduction process. A change in the interaction between a PK or PP and a natural binding partner can manifest itself as an increased or decreased
10 probability that the interaction forms, or an increased or decreased concentration of the PK or PP /natural binding partner complex.

A PK or PP natural binding partner can bind to a PK’s or PP’s intracellular region with high affinity. High affinity represents an equilibrium binding constant on the order of 10^{-6} M or less. In addition, a natural binding partner can also
15 transiently interact with a protein kinase intracellular region and chemically modify it. PK or PP natural binding partners are chosen from a group that includes, but is not limited to, SRC homology 2 (SH2) or 3 (SH3) domains, other phosphoryl tyrosine binding (PTB) domains, guanine nucleotide exchange factors, protein phosphatases, and other protein kinases. Methods of determining changes in
20 interactions between PK or PP and their natural binding partners are readily available in the art.

The term “contacting” as used herein refers to mixing a solution comprising an indolinone compound of the invention with a liquid medium bathing the cells of the methods. The solution comprising the compound may also comprise another
25 component, such as dimethylsulfoxide (DMSO), which facilitates the uptake of the indolinone compound or compounds into the cells of the methods. The solution comprising the indolinone compound may be added to the medium bathing the cells by utilizing a delivery apparatus, such as a pipet-based device or syringe-based device.

30 The indolinone compounds of the invention preferably modulate the activity of the PK or PP *in vitro*. These compounds preferably show positive results in one

or more *in vitro* assays for an activity corresponding to treatment of the disease or disorder in question (such as the assays described in the Examples below). It is understood that the indolinone compound of the invention may be one of formula I, II, or VI, or a compound obtained from the combinatorial library of the invention.

5 The invention also features a method of identifying indolinone compounds that modulate the function of PK or PP, comprising the following steps: (a) contacting cells expressing the PK or PP with the compound; and (b) monitoring an effect upon the cells. The effect upon the cells is preferably a change or an absence of a change in cell phenotype, more preferably it is a change or an absence of a
10 change in cell proliferation, even more preferably it is a change or absence of a change in the catalytic activity of the PK or PP, and most preferably it is a change or absence of a change in the interaction between the PK or PP with a natural binding partner, as described herein. It is understood that the indolinone compound of the invention may be one of formula I, II, or VI, or a compound obtained from the
15 combinatorial library of the invention.

 The term "monitoring" refers to observing the effect of adding the compound to the cells of the method. The effect can be manifested in a change in cell phenotype, cell proliferation, PK or PP catalytic activity, or in the interaction between a PK or PP and a natural binding partner.

20 The term "effect" describes a change or an absence of a change in cell phenotype or cell proliferation. "Effect" can also describe a change or an absence of a change in the catalytic activity of the pro PK or PP. "Effect" can also describe a change or an absence of a change in an interaction between the PK or PP and a natural binding partner.

25 The term "cell phenotype" refers to the outward appearance of a cell or tissue or the function of the cell or tissue. Examples of cell phenotype are cell size (reduction or enlargement), cell proliferation (increased or decreased numbers of cells), cell differentiation (a change or absence of a change in cell shape), cell survival, apoptosis (cell death), or the utilization of a metabolic nutrient (*e.g.*,
30 glucose uptake). Changes or the absence of changes in cell phenotype are readily measured by techniques known in the art.

In a preferred embodiment, the invention features a method for identifying other compounds capable of modulating the activity of PK or PP using the indolinones of the invention as a reference which method comprises the following steps: (a) lysing the cells to render a lysate comprising PK or PP; (b) adsorbing the PK or PP to an antibody; (c) incubating the adsorbed PK or PP with a substrate or substrates in the presence or absence of test compound or an indolinone compound of the present invention; and (d) adsorbing the substrate or substrates to a solid support or antibody the effect upon the cells is then monitored and the step of monitoring the effect on the cells comprises measuring the phosphate concentration of the substrate or substrates. It is understood that the indolinone compound of the invention may be one of formula I, II, or VI, or a compound obtained from the combinatorial library of the invention.

The term "antibody" refers to an antibody (*e.g.*, a monoclonal or polyclonal antibody), or antibody fragment, having specific binding affinity to PK or PP or its fragment.

By "specific binding affinity" is meant that the antibody binds to target (PK or PP) polypeptides with greater affinity than it binds to other polypeptides under specified conditions. Antibodies having specific binding affinity to a PK or PP may be used in methods for detecting the presence and/or amount of a PK or PP in a sample by contacting the sample with the antibody under conditions such that an immunocomplex forms and detecting the presence and/or amount of the antibody conjugated to the PK or PP. Diagnostic kits for performing such methods may be constructed to include a first container containing the antibody and a second container having a conjugate of a binding partner of the antibody and a label, such as, for example, a radioisotope. The diagnostic kit may also include notification of an FDA approved use and instructions therefor.

The term "polyclonal" refers to antibodies that are heterogenous populations of antibody molecules derived from the sera of animals immunized with an antigen or an antigenic functional derivative thereof. For the production of polyclonal antibodies, various host animals may be immunized by injection with the antigen.

Various adjuvants may be used to increase the immunological response, depending on the host species.

“Monoclonal antibodies” are substantially homogenous populations of antibodies to a particular antigen. They may be obtained by any technique which provides for the production of antibody molecules by continuous cell lines in culture. Monoclonal antibodies may be obtained by methods known to those skilled in the art. See, for example, Kohler, et al., *Nature* 256:495-497 (1975), and U.S. Patent No. 4,376,110.

The term “antibody fragment” refers to a portion of an antibody, often the hypervariable region and portions of the surrounding heavy and light chains, that displays specific binding affinity for a particular molecule. A hypervariable region is a portion of an antibody that physically binds to the polypeptide target.

In yet another aspect, the invention features a method for treating a disease related to unregulated tyrosine kinase or phosphatase signal transduction, where the method includes the step of administering to a subject in need thereof a therapeutically effective amount of an indolinone compound as described herein. It is understood that the indolinone compound of the invention may be one of formula I, II, or VI, or a compound obtained from the combinatorial library of the invention.

The term “treating” or “treatment” does not necessarily mean total cure. Any alleviation of any undesired symptom of the disease to any extent or the slowing down of the progress of the disease can be considered treatment. Furthermore, treatment may include acts which may worsen the patient’s overall feeling of well being or appearance. For example, the administration of chemotherapy in cancer patients which may leave the patients feeling “sicker” is still considered treatment.

The invention also features a method of regulating tyrosine kinase or phosphatase signal transduction comprising administering to a subject a therapeutically effective amount of an indolinone compound as described herein. It is understood that the indolinone compound of the invention may be one of formula I, II, or VI, or a compound obtained from the combinatorial library of the invention.

Furthermore, the invention features a method of preventing or treating an abnormal condition in an organism, where the abnormal condition is associated with

0987400:060404

an aberration in a signal transduction pathway characterized by an interaction between a PK or PP and a natural binding partner, where the method comprises the following steps: (a) administering an indolinone compound as described herein; and (b) promoting or disrupting the abnormal interaction. The organism is preferably a mammal and the abnormal condition is preferably cancer. It is understood that the indolinone compound of the invention may be one of formula I, II, or VI, or a compound obtained from the combinatorial library of the invention. The abnormal condition may also preferably be selected from the group consisting of hypertension, depression, generalized anxiety disorder, phobias, post-traumatic stress syndrome, avoidant personality disorder, sexual dysfunction, eating disorders, obesity, chemical dependencies, cluster headache, migraine, pain, Alzheimer's disease, obsessive-compulsive disorder, panic disorder, memory disorders, Parkinson's disease, endocrine disorders, vasospasm, cerebellar ataxia, and gastrointestinal tract disorders.

The term "aberration", in conjunction with a signal transduction process, refers to a PK or PP that is over- or under-expressed in an organism, mutated such that its catalytic activity is lower or higher than wild-type PK or PP activity, mutated such that it can no longer interact with a natural binding partner, is no longer modified by another protein kinase or protein phosphatase, or no longer interacts with a natural binding partner.

The term "promoting or disrupting the abnormal interaction" refers to a method that can be accomplished by administering a compound of the invention to cells or tissues in an organism. A compound can promote an interaction between a PK or PP and natural binding partners by, for example, forming favorable interactions with multiple atoms at the complex interface. Alternatively, a compound can inhibit an interaction between a PK or PP and natural binding partners by compromising favorable interactions formed between atoms at the complex interface.

The summary of the invention described above is non-limiting and other features and advantages of the invention will be apparent from the following description of the preferred embodiments, and from the claims.

Utility

The present invention relates to compounds capable of regulating and/or modulating PK or PP signal transduction and more particularly receptor and non-receptor tyrosine kinase signal transduction.

5 Receptor tyrosine kinase mediated signal transduction is initiated by extracellular interaction with a specific growth factor (ligand), followed by receptor dimerization, transient stimulation of the intrinsic protein tyrosine kinase activity and phosphorylation. Binding sites are thereby created for intracellular signal transduction molecules and lead to the formation of complexes with a spectrum of
10 cytoplasmic signaling molecules that facilitate the appropriate cellular response (e.g., cell division, metabolic effects to the extracellular microenvironment). See, Schlessinger and Ullrich, 1992, *Neuron* 9:303-391.

It has been shown that tyrosine phosphorylation sites in growth factor receptors function as high-affinity binding sites for SH2 (*src* homology) domains of
15 signaling molecules. Fantl *et al.*, 1992, *Cell* 69:413-423; Songyang *et al.*, 1994, *Mol. Cell. Biol.* 14:2777-2785; Songyang *et al.*, 1993, *Cell* 72:767-778; and Koch *et al.*, 1991, *Science* 252:668-678. Several intracellular substrate proteins that associate with receptor tyrosine kinases have been identified. They may be divided
20 into two principal groups: (1) substrates which have a catalytic domain; and (2) substrates which lack such domain but serve as adapters and associate with catalytically active molecules. Songyang *et al.*, 1993, *Cell* 72:767-778. The specificity of the interactions between receptors and SH2 domains of their substrates is determined by the amino acid residues immediately surrounding the
25 phosphorylated tyrosine residue. Differences in the binding affinities between SH2 domains and the amino acid sequences surrounding the phosphotyrosine residues on particular receptors are consistent with the observed differences in their substrate phosphorylation profiles. Songyang *et al.*, 1993, *Cell* 72:767-778. These observations suggest that the function of each receptor tyrosine kinase is determined not only by its pattern of expression and ligand availability but also by the array of
30 downstream signal transduction pathways that are activated by a particular receptor. Thus, phosphorylation provides an important regulatory step which determines the

selectivity of signaling pathways recruited by specific growth factor receptors, as well as differentiation factor receptors.

5 Tyrosine kinase signal transduction results in, among other responses, cell proliferation, differentiation and metabolism. Abnormal cell proliferation may result in a wide array of disorders and diseases, including the development of neoplasia such as carcinoma, sarcoma, leukemia, glioblastoma, hemangioma, psoriasis, arteriosclerosis, arthritis and diabetic retinopathy (or other disorders related to uncontrolled angiogenesis and/or vasculogenesis).

10 This invention is therefore directed to compounds which regulate, modulate and/or inhibit tyrosine kinase signal transduction by affecting the enzymatic activity of the RTKs and/or the non-receptor tyrosine kinases and interfering with the signal transduced by such proteins. More particularly, the present invention is directed to compounds which regulate, modulate and/or inhibit the RTK and/or non-receptor tyrosine kinase mediated signal transduction pathways as a therapeutic approach to
15 cure many kinds of solid tumors, including but not limited to carcinoma, sarcoma, erythroblastoma, glioblastoma, meningioma, astrocytoma, melanoma and myoblastoma. Indications may include, but are not limited to brain cancers, bladder cancers, ovarian cancers, gastric cancers, pancreas cancers, colon cancers, blood cancers, lung cancers and bone cancers. The present invention is also directed to
20 compounds used in a therapeutic approach to cure non-solid tumor cancers, such as, leukemia.

I. Target Diseases to be Treated by the Compounds of the Invention

25 The compounds described herein are useful for treating disorders related to unregulated tyrosine kinase signal transduction, including cell proliferative disorders, fibrotic disorders and metabolic disorders.

Cell proliferative disorders which can be treated or further studied by the present invention include cancers, blood vessel proliferative disorders and mesangial cell proliferative disorders.

30 Blood vessel proliferative disorders refer to angiogenic and vasculogenic disorders generally resulting in abnormal proliferation of blood vessels. The

formation and spreading of blood vessels, or vasculogenesis and angiogenesis, respectively, play important roles in a variety of physiological processes such as embryonic development, corpus luteum formation, wound healing and organ regeneration. They also play a pivotal role in cancer development. Other examples
 5 of blood vessel proliferation disorders include arthritis, where new capillary blood vessels invade the joint and destroy cartilage, and ocular diseases, like diabetic retinopathy, where new capillaries in the retina invade the vitreous, bleed and cause blindness, and Von Hippel-Lindau (VHL) disease, which is an inherited multi-system disorder characterized by abnormal growth of blood vessels in certain parts
 10 of the body. Conversely, disorders related to the shrinkage, contraction or closing of blood vessels, such as restenosis, are also implicated.

Fibrotic disorders refer to the abnormal formation of extracellular matrix. Examples of fibrotic disorders include hepatic cirrhosis and mesangial cell proliferative disorders. Hepatic cirrhosis is characterized by the increase in
 15 extracellular matrix constituents resulting in the formation of a hepatic scar. Hepatic cirrhosis can cause diseases such as cirrhosis of the liver. An increased extracellular matrix resulting in a hepatic scar can also be caused by viral infection such as hepatitis. Lipocytes appear to play a major role in hepatic cirrhosis. Other fibrotic disorders implicated include atherosclerosis (*see, below*).

20 Mesangial cell proliferative disorders refer to disorders brought about by abnormal proliferation of mesangial cells. Mesangial proliferative disorders include various human renal diseases, such as glomerulonephritis, diabetic nephropathy, malignant nephrosclerosis, thrombotic microangiopathy syndromes, transplant rejection, and glomerulopathies. The PDGF-R has been implicated in the
 25 maintenance of mesangial cell proliferation. Floege *et al.*, 1993, *Kidney International* 43:47S-54S.

PKs have been associated with such cell proliferative disorders. For example, some members of the RTK family have been associated with the development of cancer. Some of these receptors, like the EGFR (Tuzi *et al.*, 1991,
 30 *Br. J. Cancer* 63:227-233; Torp *et al.*, 1992, *APMIS* 100:713-719) HER2/neu (Slamon *et al.*, 1989, *Science* 244:707-712) and the PDGF-R (Kumabe *et al.*, 1992,

Oncogene 7:627-633) are overexpressed in many tumors and/or persistently activated by autocrine loops. In fact, in the most common and severe cancers these receptor overexpressions (Akbasak and Suner-Akbasak *et al.*, 1992, *J. Neurol. Sci.* 111:119-133; Dickson *et al.*, 1992, *Cancer Treatment Res.* 61:249-273; Korc *et al.*, 1992, *J. Clin. Invest.* 90:1352-1360) and autocrine loops (Lee and Donoghue, 1992, *J. Cell. Biol.* 118:1057-1070; Korc *et al.*, *supra*; Akbasak and Suner-Akbasak *et al.*, *supra*) have been demonstrated. For example, the EGFR receptor has been associated with squamous cell carcinoma, astrocytoma, glioblastoma, head and neck cancer, lung cancer and bladder cancer. HER2 has been associated with breast, ovarian, gastric, lung, pancreas and bladder cancer. The PDGF-R has been associated with glioblastoma, lung, ovarian, melanoma and prostate cancer. The RTK c-met has been generally associated with hepatocarcinogenesis and thus hepatocellular carcinoma. Additionally, c-met has been linked to malignant tumor formation. More specifically, the RTK c-met has been associated with, among other cancers, colorectal, thyroid, pancreatic and gastric carcinoma, leukemia and lymphoma. Additionally, over-expression of the c-met gene has been detected in patients with Hodgkin's disease, Burkitt's disease, and the lymphoma cell line.

The IGF-IR, in addition to being implicated in nutritional support and in type-II diabetes, has also been associated with several types of cancers. For example, IGF-I has been implicated as an autocrine growth stimulator for several tumor types, *e.g.*, human breast cancer carcinoma cells (Arteaga *et al.*, 1989, *J. Clin. Invest.* 84:1418-1423) and small lung tumor cells (Macauley *et al.*, 1990, *Cancer Res.* 50:2511-2517). In addition, IGF-I, integrally involved in the normal growth and differentiation of the nervous system, appears to be an autocrine stimulator of human gliomas. Sandberg-Nordqvist *et al.*, 1993, *Cancer Res.* 53:2475-2478. The importance of the IGF-IR and its ligands in cell proliferation is further supported by the fact that many cell types in culture (fibroblasts, epithelial cells, smooth muscle cells, T-lymphocytes, myeloid cells, chondrocytes, osteoblasts, the stem cells of the bone marrow) are stimulated to grow by IGF-I. Goldring and Goldring, 1991, *Eukaryotic Gene Expression* 1:301-326. In a series of recent publications, Baserga even suggests that IGF-I-R plays a central role in the mechanisms of transformation

and, as such, could be a preferred target for therapeutic interventions for a broad spectrum of human malignancies. Baserga, 1995, *Cancer Res.* 55:249-252; Baserga, 1994, *Cell* 79:927-930; Coppola *et al.*, 1994, *Mol. Cell. Biol.* 14:4588-4595.

5 The association between abnormalities in RTKs and disease are not restricted to cancer, however. For example, RTKs have been associated with metabolic diseases like psoriasis, diabetes mellitus, wound healing, inflammation, and neurodegenerative diseases. These diseases include, but are not limited to hypertension, depression, generalized anxiety disorder, phobias, post-traumatic
10 stress syndrome, avoidant personality disorder, sexual dysfunction, eating disorders, obesity, chemical dependencies, cluster headache, migraine, pain, Alzheimer's disease, obsessive-compulsive disorder, panic disorder, memory disorders, Parkinson's disease, endocrine disorders, vasospasm, cerebellar ataxia, and gastrointestinal tract disorders. For example, the EGF-R is indicated in corneal and
15 dermal wound healing. Defects in the Insulin-R and the IGF-1R are indicated in type-II diabetes mellitus. A more complete correlation between specific RTKs and their therapeutic indications is set forth in Plowman *et al.*, 1994, *DN&P* 7:334-339.

Not only receptor type tyrosine kinases, but also many cellular tyrosine kinases (CTKs) including src, abl, fps, yes, fyn, lyn, lck, blk, hck, fgr, yrk (reviewed
20 by Bolen *et al.*, 1992, *FASEB J.* 6:3403-3409) are involved in the proliferative and metabolic signal transduction pathway and thus in indications of the present invention. For example, mutated src (v-src) has been demonstrated as an oncoprotein (pp60^{v-src}) in chicken. Moreover, its cellular homolog, the proto-oncogene pp60^{c-src} transmits oncogenic signals of many receptors. For example,
25 overexpression of EGF-R or HER2/neu in tumors leads to the constitutive activation of pp60^{c-src}, which is characteristic for the malignant cell but absent from the normal cell. On the other hand, mice deficient for the expression of c-src exhibit an osteopetrotic phenotype, indicating a key participation of c-src in osteoclast function and a possible involvement in related disorders. Similarly, Zap 70 is implicated in
30 T-cell signaling.

Furthermore, the identification of CTK modulating compounds to augment or even synergize with RTK aimed blockers is an aspect of the present invention.

Finally, both RTKs and non-receptor type kinases have been connected to hyperimmune disorders.

5 The compounds of the present invention are also effective in treating diseases that are related to the PYK-2 protein. This protein, its cellular function, and diseases related to them are set forth in detail in U.S. Patent Number 5,837,524, issued November 17, 1998, by Lev et al., and entitled "PYK2 RELATED PRODUCTS AND METHODS" (Lyon & Lyon Docket No. 209/070), U.S. Patent Number
10 5,837,815, issued November 17, 1998, by Lev et al., and entitled "PYK2 RELATED PRODUCTS AND METHODS" (Lyon & Lyon Docket No. 211/121), U.S. Patent Application Serial Number 08/987,689, filed December 9, 1997, by Lev et al., and entitled "PYK2 RELATED PRODUCTS AND METHODS" (Lyon & Lyon Docket No. 230/110), U.S. Patent Application Serial Number 09/165,062, filed October 1,
15 1998, by Lev et al., and entitled "PYK2 RELATED PRODUCTS AND METHODS" (Lyon & Lyon Docket No. 237/201), International Publication Number WO98/26054, published June 18, 1998, by Lev et al., and entitled "PYK2 RELATED PRODUCTS AND METHODS" (Lyon & Lyon Docket No. 222/126), and International Application Number US98/27871, filed December 31, 1998, by
20 Schlessinger et al., and entitled "PYK2 AND INFLAMMATION" (Lyon & Lyon Docket No. 236/075), all of which are hereby incorporated by reference herein in their entirety, including any drawings.

 In addition, some of the compounds of the present invention are effective against rheumatoid arthritis (RA). RA is a chronic inflammatory disease mediated by
25 multiple cell types and cellular processes. Included in these are the infiltration of macrophages and T cells, and the involvement of angiogenesis. The utility of small molecule inhibitors for the treatment of RA was investigated in a rat collagen induced arthritis model. Some of the compounds of the present invention inhibit the tyrosine kinases Flk-1/KDR, pyk2, and ZAP-70 to varying degrees in biochemical
30 kinase assays. In addition, the compounds are active in cellular assays targeted to cells implicated in the pathogenesis of RA: inhibition of T cell proliferation

mediated by ZAP-70 activity, inhibition of VEGF stimulated HUVEC proliferation mediated by Flk-1/KDR activation, and inhibition of TNF- α production from murine bone marrow derived macrophages mediated by pyk2 activation. Finally, in a rodent collagen induced arthritis model, which mimics the histological and pathological changes associated with human RA, some of the compounds of the invention are efficacious in inhibiting joint swelling when dosed daily from the time of collagen immunization.

II. The KDR/FLK-1 Receptor and VEGF

Normal vasculogenesis and angiogenesis play important roles in a variety of physiological processes such as embryonic development, wound healing, organ regeneration and female reproductive processes such as follicle development in the corpus luteum during ovulation and placental growth after pregnancy. Folkman and Shing, 1992, *J. Biological Chem.* 267:10931-34. However, many diseases are driven by persistent unregulated or inappropriate angiogenesis. For example, in arthritis, new capillary blood vessels invade the joint and destroy the cartilage. In diabetes, new capillaries in the retina invade the vitreous, bleed and cause blindness. Folkman, 1987, in: *Congress of Thrombosis and Haemostasis* (Verstraete, et. al, eds.), Leuven University Press, Leuven, pp.583-596. Ocular neovascularization is the most common cause of blindness and dominates approximately twenty (20) eye diseases.

Moreover, vasculogenesis and/or angiogenesis have been associated with the growth of malignant solid tumors and metastasis. A tumor must continuously stimulate the growth of new capillary blood vessels for the tumor itself to grow. Furthermore, the new blood vessels embedded in a tumor provide a gateway for tumor cells to enter the circulation and to metastasize to distant sites in the body. Folkman, 1990, *J. Natl. Cancer Inst.* 82:4-6; Klagsbrunn and Soker, 1993, *Current Biology* 3:699-702; Folkman, 1991, *J. Natl., Cancer Inst.* 82:4-6; Weidner et al., 1991, *New Engl. J. Med.* 324:1-5.

Several polypeptides with *in vitro* endothelial cell growth promoting activity have been identified. Examples include acidic and basic fibroblastic growth factor

(aFGF, bFGF), vascular endothelial growth factor (VEGF) and placental growth factor. Unlike aFGF and bFGF, VEGF has recently been reported to be an endothelial cell specific mitogen. Ferrara and Henzel, 1989, *Biochem. Biophys. Res. Comm.* 161:851-858; Vaisman *et al.*, 1990, *J. Biol. Chem.* 265:19461-19566.

Thus, the identification of the specific receptors to which VEGF binds is an important advancement in the understanding of the regulation of endothelial cell proliferation. Two structurally closely related RTKs have been identified to bind VEGF with high affinity: the flt-1 receptor (Shibuya *et al.*, 1990, *Oncogene* 5:519-524; De Vries *et al.*, 1992, *Science* 255:989-991) and the KDR/FLK-1 receptor, (also referred to as VEGFR2) discussed in the U.S. Patent Application Nos. 08/193,829 and 08/965,598. Consequently, it had been surmised that these RTKs may have a role in the modulation and regulation of endothelial cell proliferation.

Evidence, such as the disclosure set forth in copending U.S. Application Serial Nos. 08/193,829 and 08/965,598, strongly suggests that VEGF is not only responsible for endothelial cell proliferation, but also is a prime regulator of normal and pathological angiogenesis. *See generally*, Klagsburn and Soker, 1993, *Current Biology* 3:699-702; Houck *et al.*, 1992, *J. Biol. Chem.* 267:26031-26037. Moreover, it has been shown that KDR/FLK-1 and flt-1 are abundantly expressed in the proliferating endothelial cells of a growing tumor, but not in the surrounding quiescent endothelial cells. Plate *et al.*, 1992, *Nature* 359:845-848; Shweiki *et al.*, 1992, *Nature* 359:843-845.

III. Identification Of Agonists And Antagonists To The KDR/FLK-1 Receptor

In view of the deduced importance of RTKs in the control, regulation and modulation of endothelial cell proliferation and potentially vasculogenesis and/or angiogenesis, many attempts have been made to identify RTK "inhibitors" using a variety of approaches. These include the use of mutant ligands (U.S. Patent No. 4,966,849); soluble receptors and antibodies (International Publication No. WO 94/10202; Kendall and Thomas, 1994, *Proc. Natl. Acad. Sci. USA* 90:10705-10709; Kim *et al.*, 1993, *Nature* 362:841-844); and RNA ligands (Jellinek *et al.*, 1994, *Biochemistry* 33:10450-10456).

Furthermore, tyrosine kinase inhibitors (WO 94/03427; WO 92/21660; WO 91/15495; WO 94/14808; U.S. Patent No. 5,330,992; Mariani *et al.*, 1994, *Proc. Am. Assoc. Cancer Res.* 35:2268), and inhibitors acting on receptor tyrosine kinase signal transduction pathways, such as protein kinase C inhibitors have been identified
 5 (Schuchter *et al.*, 1991, *Cancer Res.* 51:682-687); Takano *et al.*, 1993, *Mol. Bio. Cell* 4:358A; Kinsella *et al.*, 1992, *Exp. Cell Res.* 199:56-62; Wright *et al.*, 1992, *J. Cellular Phys.* 152:448-57).

More recently, attempts have been made to identify small molecules which act as tyrosine kinase inhibitors. For example, bis monocyclic, bicyclic or
 10 heterocyclic aryl compounds (PCT WO 92/20642), vinylene-azaindole derivatives (PCT WO 94/14808) and 1-cyclopropyl-4-pyridyl-quinolones (U.S. Patent No. 5,330,992) have been described generally as tyrosine kinase inhibitors. Styryl compounds (U.S. Patent No. 5,217,999), styryl-substituted pyridyl compounds (U.S. Patent No. 5,302,606), certain quinazoline derivatives (EP Application No. 0 566
 15 266 A1), seleoindoles and selenides (PCT WO 94/03427), tricyclic polyhydroxylic compounds (PCT WO 92/21660) and benzylphosphonic acid compounds (PCT WO 91/15495) have been described as compounds for use as tyrosine kinase inhibitors for use in the treatment of cancer.

Consequently, there is an unmet need for the identification and generation of
 20 effective small compounds which selectively inhibit the signal transduction of the KDR/FLK-1 receptor in order to effectively and specifically suppress vasculogenesis.

Some of the compounds of the present invention demonstrate excellent activity in biological assays and thus these compounds and related compounds are
 25 expected to be effective in treating Flk related disorders such as those driven by persistent unregulated or inappropriate angiogenesis.

Pharmaceutical Formulations And Routes Of Administration

The compounds described herein can be administered to a human patient *per*
 30 *se*, or in pharmaceutical compositions where they are mixed with other active ingredients, as in combination therapy, or suitable carriers or excipient(s).

Techniques for formulation and administration of the compounds of the instant application may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition.

1. Routes Of Administration

5 Suitable routes of administration may, for example, include oral, rectal, transmucosal, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intravenous, intramedullary injections, as well as intrathecal, direct intraventricular, intraperitoneal, intranasal, or intraocular injections.

10 Alternately, one may administer the compound in a local rather than systemic manner, for example, via injection of the compound directly into a solid tumor, often in a depot or sustained release formulation.

 Furthermore, one may administer the drug in a targeted drug delivery system, for example, in a liposome coated with tumor-specific antibody. The liposomes will
15 be targeted to and taken up selectively by the tumor.

2. Composition/Formulation

 The pharmaceutical compositions of the present invention may be manufactured in a manner that is itself known, *e.g.*, by means of conventional
20 mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

 Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which
25 facilitate processing of the active compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

 For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution,

Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition,

stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

5 For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a
10 pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of *e.g.* gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds may be formulated for parenteral administration by
15 injection, *e.g.*, by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, *e.g.*, in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

20 Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes.
25 Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

30 Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

A pharmaceutical carrier for the hydrophobic compounds of the invention is a cosolvent system comprising benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer, and an aqueous phase. The cosolvent system may be the VPD co-solvent system. VPD is a solution of 3% w/v benzyl alcohol, 8% w/v of the nonpolar surfactant Polysorbate 80™, and 65% w/v polyethylene glycol 300, made up to volume in absolute ethanol. The VPD co-solvent system (VPD:D5W) consists of VPD diluted 1:1 with a 5% dextrose in water solution. This co-solvent system dissolves hydrophobic compounds well, and itself produces low toxicity upon systemic administration. Naturally, the proportions of a co-solvent system may be varied considerably without destroying its solubility and toxicity characteristics. Furthermore, the identity of the co-solvent components may be varied: for example, other low-toxicity nonpolar surfactants may be used instead of Polysorbate 80™; the fraction size of polyethylene glycol may be varied; other biocompatible polymers may replace polyethylene glycol, *e.g.*, polyvinyl pyrrolidone; and other sugars or polysaccharides may substitute for dextrose.

Alternatively, other delivery systems for hydrophobic pharmaceutical compounds may be employed. Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed, although usually at the cost of greater toxicity. Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing

the therapeutic agent. Various sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization may be employed.

The pharmaceutical compositions also may comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

Many of the PK modulating compounds of the invention may be provided as salts with pharmaceutically compatible counterions. Pharmaceutically compatible salts may be formed with many acids, including but not limited to hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms.

3. Effective Dosage

Pharmaceutical compositions suitable for use in the present invention include compositions where the active ingredients are contained in an amount effective to achieve its intended purpose. More specifically, a therapeutically effective amount means an amount of compound effective to prevent, alleviate or ameliorate symptoms of disease or prolong the survival of the subject being treated. Determination of a therapeutically effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

For any compound used in the methods of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. For example, a dose can be formulated in animal models to achieve a circulating concentration range that includes the IC_{50} as determined in cell culture (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of the PK activity). Such information can be used to more accurately determine useful doses in humans.

Toxicity and therapeutic efficacy of the compounds described herein can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between LD₅₀ and ED₅₀. Compounds which exhibit high therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See *e.g.*, Fingl *et al.*, 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p.1).

Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the kinase modulating effects, or minimal effective concentration (MEC). The MEC will vary for each compound but can be estimated from *in vitro* data; *e.g.*, the concentration necessary to achieve 50-90% inhibition of the kinase using the assays described herein.

Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration. However, HPLC assays or bioassays can be used to determine plasma concentrations.

Dosage intervals can also be determined using MEC value. Compounds should be administered using a regimen which maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90%.

In cases of local administration or selective uptake, the effective local concentration of the drug may not be related to plasma concentration.

The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.

4. Packaging

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. The pack or dispenser may also be accompanied with a notice associated with the container in form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the polynucleotide for human or veterinary administration. Such notice, for example, may be the labeling approved by the U.S. Food and Drug Administration for prescription drugs, or the approved product insert. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition. Suitable conditions indicated on the label may include treatment of a tumor, inhibition of angiogenesis, treatment of fibrosis, diabetes, and the like.

IV. Biological Activity of the Indolinone Compounds of the Invention

Some of the indolinone compounds of the present invention were tested for their ability to inhibit most of protein tyrosine kinase activity. The biological assays and results of these inhibition studies are reported herein. The methods used to measure indolinone compound modulation of protein kinase function are similar to those described in International Publication No. WO 98/07695, published March 26, 1998, by Tang et al., and entitled "Indolinone Combinatorial Libraries and Related Products and Methods for the Treatment of Disease," with respect to the high throughput aspect of the method. The WO 98/07695 publication is incorporated herein by reference in its entirety, including any drawings.

V. Pharmaceutical Compositions and Administration of Indolinone Compounds of the Invention

Methods of preparing pharmaceutical formulations of the compounds, methods of determining the amounts of compounds to be administered to a patient, and modes of administering compounds to an organism are disclosed in International Publication No. WO 98/07695, by Tang et al., and entitled "Indolinone Combinatorial Libraries and Related Products and Methods for the Treatment of Disease," and International Publication No. WO 96/22976, by Buzzetti et al., and entitled "Hydrosoluble 3-Arylidene-2-Oxindole Derivatives as Tyrosine Kinase Inhibitors," published August 1, 1996, both of which are incorporated herein by reference in their entirety, including any drawings. Those skilled in the art will appreciate that such descriptions are applicable to the present invention and can be easily adapted to it.

15 EXAMPLES

The examples below are non-limiting and are merely representative of various aspects and features of the present invention. The examples describe methods for synthesizing compounds of the invention and methods for measuring an effect of a compound on the function of protein tyrosine kinases.

20 The cells used in the methods are commercially available. The nucleic acid vectors harbored by the cells are also commercially available and the sequences of genes for the various protein kinases are readily accessible in sequence data banks. Thus, a person of ordinary skill in the art can readily recreate the cell lines in a timely manner by combining the commercially available cells, the commercially available nucleic acid vectors, and the protein kinase genes using techniques readily available to persons of ordinary skill in the art.

SYNTHETIC PROCEDURES

The following examples describe methods of synthesizing some of the compounds of the invention.

5 EXAMPLE 1: Preparation of 5-(2-dimethylamino-ethoxy)-1H-indole-2-carbaldehyde

A mixture of 5-hydroxy-1H-indole-2-carboxylic acid ethyl ester (2 g, 10 mmol), 2-chloro-N,N-dimethylacetamide (1.33 g, 11 mmol) and cesium carbonate (9.8 g, 30 mmol)) in dimethylformamide (20 mL) was stirred at room temperature
10 for 16 hours. The reaction was diluted with ethyl acetate (150 mL), washed with water (5x 50 mL) and brine, dried and concentrated. The residue was recrystallized from ethyl acetate and hexane to give 1.4 g of 5-dimethylcarbamoylmethoxy-1H-indole-2-carboxylic acid ethyl ester as a brown solid.

¹HNMR (300 MHz, DMSO-d₆) δ 11.74 (s, br, 1H, NH), 7.33 (d, *J* = 9.0 Hz, 1H, H-7), 7.06 (d, *J* = 2.4 Hz, 1H, H-4), 7.01 (d, *J* = 1.5 Hz, 1H, H-3), 6.94 (dd, *J* = 2.4 & 9.0 Hz, 1H, H-6), 4.74 (s, 2H, OCH₂CON(CH₃)₂), 4.31 (q, *J* = 7.1 Hz, 2H, OCH₂CH₃), 3.0 (s, 3H, NCH₃), 2.83 (s, 3H, NCH₃), 1.32 (t, *J* = 7.1 Hz, 3H, OCH₂CH₃).

MS-EI *m/z* 290 [M⁺].

20 Lithium aluminum hydride (1.7 g, 45 mmol) was added to 5-dimethylcarbamoylmethoxy-1H-indole-2-carboxylic acid ethyl ester (1.4 g, 4.8 mmol) in tetrahydrofuran (80 mL). The mixture was stirred at room temperature for 4 hours and cooled to 0 °C. It was then quenched with water (1.7 mL), 15% sodium hydroxide (1.7 mL) and water (1.7 mL). The precipitate was filtered off and the
25 filtrate was concentrated to give 0.6 g (54%) of [5-(2-dimethylamino-ethoxy)-1H-indol-2-yl]-methanol.

¹HNMR (300 MHz, DMSO-d₆) δ 10.79 (s, br, 1H, NH), 7.17 (d, *J* = 8.9 Hz, 1H, H-7), 6.95 (d, *J* = 2.1 Hz, H-4), 6.64 (dd, *J* = 2.1 & 8.9 Hz, 1H, H-6), 6.61 (d, *J* = 1.5 Hz, H-3), 5.18 (d, *J* = 5.5 Hz, 1H, CH₂OH), 4.54 (d, *J* = 5.5 Hz, 2H, CH₂OH),
30 3.98 (t, *J* = 6.0 Hz, 2H, OCH₂CH₂N), 2.59 (t, *J* = 6.0 Hz, 2H, OCH₂CH₂N), 2.20 (s, 6H, N(CH₃)₂).

Manganese (IV) oxide (3.3 g, 38 mmol) was added to 5-(2-dimethylamino-ethoxy)-1*H*-indol-2-yl]-methanol (0.6 g, 2.5 mmol) in dichloromethane (100 mL). The mixture was stirred at room temperature for 16 hours. The precipitate was filtered off and the filtrate was concentrated to give 0.41 g (71%) of 5-(2-dimethylamino-ethoxy)-1*H*-indole-2-carbaldehyde.

¹HNMR (360 MHz, DMSO-d₆) δ 11.78 (s, br, 1H, NH), 9.78 (s, 1H, CHO), 7.34 (d, *J* = 8.8 Hz, 1H, H-7), 7.25 (d, *J* = 1.1 Hz, H-3), 7.18 (d, *J* = 2.3 Hz, H-4), 6.98 (dd, *J* = 2.3 & 8.8 Hz, 1H, H-6), 4.04 (t, *J* = 5.9 Hz, 2H, OCH₂CH₂N), 2.63 (t, *J* = 5.9 Hz, 2H, OCH₂CH₂N), 2.22 (s, 6H, N(CH₃)₂).

EXAMPLE 2: Preparation of 5-(2-pyrrolidin-1-yl-ethoxy)-1H-indole-2-carbaldehyde

20 ¹H NMR (300 MHz, DMSO-*d*₆) δ 4.03 (s, 2H, BrCH₂CO), 3.45 (t, *J* = 6.6 Hz, 2H), 3.27 (t, *J* = 6.6 Hz, 2H), 1.74-1.89 (m, 4H).

¹HNMR (300 MHz, DMSO-d₆) δ 11.74 (s, br, 1H, NH), 7.34 (d, *J* = 8.7 Hz, 1H, H-7), 7.07 (d, *J* = 2.5 Hz, 1H, H-4), 7.02 (d, *J* = 1.5 Hz, 1H, H-3), 6.95 (dd, *J* = 2.5 & 8.7 Hz, 1H, H-6), 4.67 (s, 2H, OCH₂CON), 4.31 (q, *J* = 7.1 Hz, 2H,

OCH₂CH₃), 3.47 (t, J = 6.7 Hz, 2H), 3.31 (t, J = 6.7 Hz, 2H), 1.88 (t, J = 6.7 Hz, 2H), 1.76 (t, J = 6.7 Hz, 2H), 1.32 (t, J = 7.1 Hz, 3H, OCH₂CH₃).

MS-EI m/z 316 [M^+].

Lithium aluminum hydride (0.4 g, 10 mmol) was added to 5-(2-oxo-2-pyrrolidin-1-yl-ethoxy)-1*H*-indole-2-carboxylic acid ethyl ester (0.6 g, 1.9 mmol) in tetrahydrofuran (120 mL). The mixture was stirred at room temperature for 16 hours and cooled to 0 °C. It was then quenched with water (0.4 mL), 15% sodium hydroxide (0.4 mL) and water (0.4 mL). The precipitate was filtered off and the filtrate was concentrated to give 0.31 g (63%) of [5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indo-2-yl]-methanol as a white solid.

¹HNMR (360 MHz, DMSO- d_6) δ 10.74 (s, br, 1H, NH), 7.17 (d, J = 8.8 Hz, 1H, H-7), 6.94 (d, J = 2.5 Hz, H-4), 6.65 (dd, J = 2.5 & 8.8 Hz, 1H, H-6), 6.15 (d, J = 0.7 Hz, H-3), 5.12 (d, J = 5.5 Hz, 1H, CH₂OH), 4.55 (d, J = 5.5 Hz, 2H, CH₂OH), 4.01 (t, J = 6.0 Hz, 2H, OCH₂CH₂N), 2.76 (t, J = 6.0 Hz, 2H, OCH₂CH₂N), 2.53 (m, under DMSO, H-pyrrolidine), 1.67 (m, 4H, H-pyrrolidine).

MS m/z 261.1 [$M^+ + 1$].

Manganese (IV) oxide (1 g, 12 mmol) was added to [5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indo-2-yl]-methanol (0.31 g, 1.2 mmol) in dichloromethane (100 mL). The mixture was stirred at room temperature for 4 hours. The precipitate was filtered off and the filtrate was concentrated to give 0.19 g (61%) of 5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indole-2-carbaldehyde.

¹HNMR (360 MHz, DMSO- d_6) δ 11.78 (s, br, 1H, NH), 9.79 (s, 1H, CHO), 7.34 (d, J = 9.0 Hz, 1H, H-7), 7.25 (d, J = 1.1 Hz, H-3), 7.18 (d, J = 2.4 Hz, H-4), 6.98 (dd, J = 2.4 & 9.0 Hz, 1H, H-6), 4.06 (t, J = 5.8 Hz, 2H, OCH₂CH₂N), 2.79 (t, J = 5.8 Hz, 2H, OCH₂CH₂N), 2.52 (m, under DMSO, H-pyrrolidine), 1.67 (m, 4H, H-pyrrolidine).

MS-EI m/z 258 [M^+].

EXAMPLE 3: Preparation of 5-(2-morpholin-4-yl-ethoxy)-1H-indole-2-carbaldehyde

A mixture of 5-hydroxy-1H-indole-2-carboxylic acid ethyl ester (2 g, 10 mmol), tert-butyl bromoacetate (2.35 g, 12 mmol) and cesium carbonate (10.5 g, 30 mmol) in dimethylformamide (10 mL) was stirred at room temperature for 16 hours.

The reaction was diluted with ethyl acetate (150 mL), washed with water (5x 50 mL) and brine, dried and concentrated. The residue was recrystallized from ethyl acetate and hexane to give 2 g (63%) 5-tert-butoxycarbonylmethoxy-1H-indole-2-carboxylic acid ethyl ester.

¹HNMR (360 MHz, DMSO-d₆) δ 11.71 (s, br, 1H, NH), 7.34 (d, *J* = 9.0 Hz, 1H, H-7), 7.02-7.03 (m, 2H, H-3 & 4), 6.93 (dd, *J* = 2.3 & 9.0 Hz, 1H, H-6), 4.59 (s, 2H, OCH₂CO), 4.31 (q, *J* = 7.0 Hz, 2H, OCH₂CH₃), 1.41 (s, 9H, C(CH₃)₃), 1.32 (t, *J* = 7.0 Hz, 3H, OCH₂CH₃).

MS-EI *m/z* 319 [M⁺].

A mixture of 5-tert-butoxycarbonylmethoxy-1H-indole-2-carboxylic acid ethyl ester (2 g, 6 mmol) and trifluoroacetic acid (8 mL) in dichloromethane (8 mL) was stirred at room temperature for 2 hours. The reaction was concentrated to give 1.6 g (97%) of 5-carboxymethoxy-1H-indole-2-carboxylic acid ethyl ester as a light brown solid.

¹HNMR (360 MHz, DMSO-d₆) δ 11.70 (s, br, 1H, NH), 7.35 (d, *J* = 8.8 Hz, 1H, H-7), 7.04 (d, *J* = 2.2 Hz, 1H, H-4), 7.02 (m, 1H, H-3), 6.94 (dd, *J* = 2.2 & 8.8 Hz, 1H, H-6), 4.62 (s, 2H, OCH₂CO), 4.31 (q, *J* = 7.2 Hz, 2H, OCH₂CH₃), 1.32 (t, *J* = 7.2 Hz, 3H, OCH₂CH₃).

MS-EI *m/z* 263 [M⁺].

A mixture of 5-carboxymethoxy-1H-indole-2-carboxylic acid ethyl ester (1.5 g, 5.7 mmol) and 1-1'-carbonyldiimidazole (1.11 g, 6.85 mmol) in dimethylformamide (15 mL) was stirred at room temperature for 30 mins. To the mixture was then added morpholine (1 mL) and the stirring was continued for overnight. The reaction was concentrated and the residue was dissolved in dichloromethane (200 mL), washed with 1N hydrochloric acid, saturated sodium

bicarbonate and brine, dried and concentrated to give 1.6 g (85%) of 5-(2-morpholin-4-yl-2-oxo-ethoxy)-1*H*-indole-2-carboxylic acid ethyl ester.

¹HNMR (360 MHz, DMSO-*d*₆) δ 11.71 (s, br, 1H, NH), 7.35 (d, *J* = 9.0 Hz, 1H, H-7), 7.10 (d, *J* = 2.3 Hz, 1H, H-4), 7.02 (m, 1H, H-3), 6.95 (dd, *J* = 2.3 & 9.0 Hz, 1H, H-6), 4.77 (s, 2H, OCH₂CO), 4.32 (q, *J* = 7.2 Hz, 2H, OCH₂CH₃), 3.46-3.58 (m, 8H, 4x CH₂), 1.32 (t, *J* = 7.2 Hz, 3H, OCH₂CH₃).

Lithium aluminum hydride (0.46 g, 12 mmol) was added 5-(2-morpholin-4-yl-2-oxo-ethoxy)-1*H*-indole-2-carboxylic acid ethyl ester (0.8 g, 2.4 mmol) in tetrahydrofuran (50 mL). The mixture was stirred at room temperature for 16 hours and cooled to 0 °C. It was then quenched with water (4.6 mL), 15% sodium hydroxide (4.6 mL) and water (4.6 mL). The precipitate was filtered off and the filtrate was concentrated to give 0.7 g of [5-(2-morpholin-4-yl-ethoxy)-1*H*-indol-2-yl]-methanol.

¹HNMR (360 MHz, DMSO-*d*₆) δ 10.74 (s, br, 1H, NH), 7.17 (d, *J* = 8.6 Hz, 1H, H-7), 6.95 (m, 1H, H-4), 6.66 (m, 1H, H-6), 6.15 (s, 1H, H-3), 5.11 (d, *J* = 5.4 Hz, 1H, CH₂OH), 4.55 (d, *J* = 5.4 Hz, 2H, CH₂OH), 4.03 (t, *J* = 5.7 Hz, 2H, OCH₂CH₂N), 3.56 (m, 4H, 2x CH₂), 2.67 (t, *J* = 5.7 Hz, 2H, OCH₂CH₂N), 2.47 (m, under DMSO, 2x CH₂).

MS-EI *m/z* 276 [*M*⁺+2].

Manganese (IV) oxide (3.2 g, 36 mmol) was added to [5-(2-morpholin-4-yl-ethoxy)-1*H*-indol-2-yl]-methanol (0.7 g) in dichloromethane (150 mL). The mixture was stirred at room temperature for 16 hours. The precipitate was filtered off and the filtrate was concentrated. The residue was recrystallized from ethyl acetate and hexane to give 0.18 g of 5-(2-morpholin-4-yl-ethoxy)-1*H*-indole-2-carbaldehyde.

¹HNMR (360 MHz, DMSO-*d*₆) δ 11.78 (s, br, 1H, NH), 9.79 (s, 1H, CHO), 7.34 (d, *J* = 9.2 Hz, 1H, H-7), 7.25 (m, 1H, H-3), 7.19 (d, *J* = 1.8 Hz, 1H, H-4), 6.99 (dd, *J* = 1.8 & 9.2 Hz, 1H, H-6), 4.08 (t, *J* = 5.7 Hz, 2H, OCH₂CH₂N), 3.57 (m, 4H, 2x CH₂), 2.70 (t, *J* = 5.7 Hz, 2H, OCH₂CH₂N), 2.48 (m, under DMSO, 2x CH₂).

EXAMPLE 4: Preparation of 5-(2-diethylamino-ethoxy)-1H-indole-2-carbaldehyde

A mixture of 5-hydroxy-1H-indole-2-carboxylic acid ethyl ester (1 g, 5 mmol), 2-chloro-N,N-diethylacetamide (1 mL, 7.5 mmol) and cesium carbonate (5 g, 15 mmol) in dimethylformamide (10 mL) was stirred at room temperature for 6 hours. The reaction was diluted with ethyl acetate (160 mL), washed with water (5x 50 mL) and brine, dried and concentrated. The residue was recrystallized from ethyl acetate and hexane to give 0.52 g of 5-diethylcarbamoylmethoxy-1H-indole-2-carboxylic acid ethyl ester as an off-white solid.

¹HNMR (360 MHz, DMSO-d₆) δ 11.70 (s, br, 1H, NH), 7.34 (d, *J* = 9.0 Hz, 1H, H-7), 7.08 (d, *J* = 2.3 Hz, 1H, H-4), 7.02 (s, 1H, H-3), 6.94 (dd, *J* = 2.3 & 9.0 Hz, 1H, H-6), 4.71 (s, 2H, OCH₂CON(CH₂CH₃)₂), 4.31 (q, *J* = 7.2 Hz, 2H, OCH₂CH₃), 3.25-3.38 (m, 4H, N(CH₂CH₃)₂), 1.32 (t, *J* = 7.2 Hz, 3H, OCH₂CH₃), 1.15 (t, *J* = 7.0 Hz, 3H, NCH₂CH₃), 1.03 (t, *J* = 7.0 Hz, 3H, NCH₂CH₃).

MS-EI *m/z* 318 [M⁺].

Lithium aluminum hydride (3.6 g, 45 mmol) was added to 5-diethylcarbamoylmethoxy-1H-indole-2-carboxylic acid ethyl ester (0.52 g, 1.6 mmol) in tetrahydrofuran (80 mL). The mixture was stirred at room temperature for 16 hours and cooled to 0 °C. It was then quenched with water (3.6 mL), 15% sodium hydroxide (3.6 mL) and water (3.6 mL). The precipitate was filtered off and the filtrate was concentrated to give 0.4 g (43%) of [5-(2-diethylamino-ethoxy)-1H-indol-2-yl]-methanol.

¹HNMR (300 MHz, DMSO-d₆) δ 10.78 (s, br, 1H, NH), 7.16 (d, *J* = 8.5 Hz, 1H, H-7), 6.94 (d, *J* = 2.3 Hz, 1H, H-4), 6.64 (dd, *J* = 2.3 & 8.5 Hz, 1H, H-6), 6.15 (d, *J* = 1.2 Hz, 1H, H-3), 5.17 (t, *J* = 5.6 Hz, 1H, CH₂OH), 4.54 (d, *J* = 5.6 Hz, 2H, CH₂OH), 3.96 (t, *J* = 6.3 Hz, 2H, OCH₂CH₂N), 2.75 (t, *J* = 6.3 Hz, 2H, OCH₂CH₂N), 2.54 (q, *J* = 7.1 Hz, 4H, 2x NCH₂CH₃), 0.96 (t, *J* = 7.1 Hz, 6H, 2x NCH₂CH₃).

MS *m/z* 263.2 [M⁺+1].

Manganese (IV) oxide (1.95 g, 23 mmol) was added to 5-(2-diethylamino-ethoxy)-1H-indol-2-yl]-methanol (0.4 g, 1.5 mmol) in dichloromethane (100 mL). The mixture was stirred at room temperature for 24 hours. The precipitate was

filtered off and the filtrate was concentrated to give 0.31 g (79%) of 5-(2-diethylamino-ethoxy)-1*H*-indole-2-carbaldehyde.

¹HNMR (360 MHz, DMSO-*d*₆) δ 11.78 (s, 1H, NH), 9.78 (s, 1H, CHO), 7.34 (d, *J* = 9.0 Hz, 1H, H-7), 7.25 (d, *J* = 1.4 Hz, 1H, H-3), 7.18 (d, *J* = 2.4 Hz, 1H, H-4), 6.97 (dd, *J* = 2.4 & 9.0 Hz, 1H, H-6), 4.01 (t, *J* = 6.2 Hz, 2H, OCH₂CH₂N), 2.78 (t, *J* = 6.2 Hz, 2H, OCH₂CH₂N), 2.55 (q, *J* = 7.0 Hz, 4H, 2x NCH₂CH₃), 0.97 (t, *J* = 7.0 Hz, 6H, 2x NCH₂CH₃).

MS-EI *m/z* 260 [M⁺].

10 EXAMPLE 5: Preparation of 5-(3-diethylamino-propyl)-1*H*-indole-2-carbaldehyde

A mixture of 5-bromo-1*H*-indole-2-carboxylic acid ethyl ester (5.36 g, 20 mmol) in triethylamine (30 mL) and pyridine (6 mL) was degassed by bubbling nitrogen through for 10 minutes, followed by the addition of diethylpropargyl amine and the degassing was continued for another 5 minutes.

15 Dichlorobis(triphenylphosphine)palladium(II) (702 mg, 1 mmol) and copper (I) iodide (95 mg, 0.5 mmol) was added to the mixture and it was heated at 90 °C under nitrogen for 17 hours. The reaction was concentrated and the residue was partitioned between 1 N HCl (150 mL) and ethyl acetate (300 mL). The layers were separated, the aqueous layer was extracted with another 300 mL of ethyl acetate.

20 The combined organic layers were washed with brine, dried and concentrated to give 3.4 g (57%) of 5-(3-diethylamino-prop-1-ynyl)-1*H*-indole-2-carboxylic acid ethyl ester (hydrochloride salt) as a light brown crystalline solid.

¹HNMR (360 MHz, DMSO-*d*₆) δ 12.14 (s, 1H, NH), 11.23 (s, br, 1H, NH), 7.89 (s, 1H, H-3), 7.49 (d, *J* = 8.4 Hz, 1H, H-7), 7.36 (dd, *J* = 1.4 & 8.4 Hz, 1H, H-6), 7.17 (d, *J* = 1.4 Hz, 1H, H-4), 4.34 (q, *J* = 7.1 Hz, 2H, OCH₂CH₃), 4.29 (s, CH₂), 3.16-3.24 (m, 4H, 2xNCH₂CH₃), 1.28-1.35 (9H, 3xCH₃).

MS-EI *m/z* 298 [M⁺].

5-(3-Diethylamino-prop-1-ynyl)-1*H*-indole-2-carboxylic acid ethyl ester hydrochloride (2g, 6.7 mmol) was hydrogenated over palladium on carbon (10%) (713 mg, 0.67 mmol) under nitrogen for 5 hours. Palladium hydroxide on carbon

30

(20%) (2x350 mg) and acetic acid (4mL) were added to the mixture and continued to stir for another 12 hours. The reaction was filtered through celite and the filtrate was concentrated. The residue was partitioned between 1 N HCl (150 mL) and ethyl acetate (300mL). The aqueous layer was made basic to pH 8, extracted with ethyl acetate, washed with brine, dried and concentrated to give 1.126 g orange solid.

NMR spectrum of the above solid showed desired saturated compound and some alkene material. Therefore, the solid was dissolved in ethanol (10 mL) and acetic acid (7 mL) and it was hydrogenated over palladium hydroxide (175 mg) for 42 hours. Following the work-up as outlined above, 1.2 g (59%) of 5-(3-diethylamino-propyl)-1*H*-indole-2-carboxylic acid ethyl ester (acetate salt) was obtained.

¹HNMR (360 MHz, DMSO-d₆) δ 11.70 (s, br, 1H, NH), 7.42 (s, 1H, H-3), 7.35 (d, *J* = 8.5 Hz, 1H, H-7), 7.11 (dd, *J* = 1.8 & 8.5 Hz, 1H, H-6), 7.04 (d, *J* = 1.8 Hz, 1H, H-4), 4.32 (q, *J* = 7.1 Hz, 2H, OCH₂CH₃), 2.62 (t, *J* = 7.7 Hz, 2H, CH₂), 2.39-2.50 (m, 6H, 2xNCH₂CH₃ & CH₂), 1.68-1.72 (m, 2H, CH₂), 1.32 (t, *J* = 7.1 Hz, 3H, OCH₂CH₃), 0.92 (t, *J* = 6.8 Hz, 6H, 2x NCH₂CH₃).

MS-EI *m/z* 302 [M⁺].

5-(3-Diethylamino-propyl)-1*H*-indole-2-carboxylic acid ethyl ester (1.2 g, 3.97 mmol) in tetrahydrofuran (48 mL) was added dropwise to the lithium aluminum hydride (602 mg, 15.87 mmol) stirred in tetrahydrofuran (30 mL) under nitrogen. After stirring for 2 hours, the reaction was quenched in water (0.6 mL), 15% sodium hydroxide (0.6 mL) and then water (1.8 mL). The mixture was stirred overnight. The precipitate was filtered off and the filtrate was concentrated to give 680 mg (66%) of [5-(3-diethylamino-propyl)-1*H*-indol-2-yl]-methanol.

¹HNMR (300 MHz, DMSO-d₆) δ 10.83 (s, br, 1H, NH), 7.22 (s, 1H, H-3), 7.20 (d, *J* = 8.4 Hz, 1H, H-7), 6.85 (dd, *J* = 1.8 & 8.4 Hz, 1H, H-6), 6.16 (d, *J* = 1.8 Hz, 1H, H-4), 5.18 (t, *J* = 5.4 Hz, 1H, OH), 4.55 (d, *J* = 5.4 Hz, 2H, CH₂OH), 2.58 (t, *J* = 7.5 Hz, 2H, CH₂), 2.33-2.45 (m, 6H, 3xCH₂), 1.64-1.72 (m, 2H, CH₂), 0.90 (t, *J* = 7.0 Hz, 6H, 2x NCH₂CH₃).

MS-EI *m/z* 260 [M⁺].

Manganese oxide (3.75g, 43.1 mmol) was added portionwise to [5-(3-diethylamino-propyl)-1*H*-indol-2-yl]-methanol (660 mg, 2.53 mmol) dissolved in dichloromethane (68 mL) under nitrogen at room temperature. The precipitate was filtered through celite, washing with hot dichloromethane. The filtrate was concentrated to give 563 mg (83%) of 5-(3-diethylamino-propyl)-1*H*-indole-2-carbaldehyde as a waxy brown solid.

¹HNMR (360 MHz, DMSO-d₆) δ 11.80 (s, br, 1H, NH), 9.80 (s, 1H, CHO), 7.50 (s, 1H, H-3), 7.35 (d, *J* = 8.5 Hz, 1H, H-7), 7.28 (s, 1H, H-4), 7.17 (dd, *J* = 1.1 & 8.5 Hz, 1H, H-6), 2.63 (t, *J* = 7.5 Hz, 2H, CH₂), 2.34-2.45 (m, 6H, 3xCH₂), 1.67-1.71 (m, 2H, CH₂), 0.91 (t, *J* = 7.0 Hz, 6H, 2x NCH₂CH₃).

MS-EI *m/z* 258 [M⁺].

EXAMPLE 6: Preparation of 5-(3-pyrrolidin-1-yl-propyl)-1*H*-indole-2-carbaldehyde

A mixture of 5-bromoindole-2-carboxylic acid ethyl ester (2.68 g, 10 mmol), tert-butyl acrylate (4.4 mL, 30 mmol) and dichlorobis(triphenylphosphine)palladium(II) (0.7 g, 1 mmol) and triethylamine (17 mL) in acetic acid (17 mL) was heated at 120 °C in a closed vessel for 6 hrs. The cooled reaction mixture was poured into water and dichloromethane. The biphasic mixture was filtered through a pad of celite and the aqueous layer was extracted with dichloromethane (2x). The combined organic layers were washed with saturated sodium bicarbonate solution, brine, dried and concentrated. The residue was purified by column chromatography (10% of ethyl acetate in hexanes) to give 900 mg (28%) of 5-(2-*tert*-butoxycarbonyl-vinyl)-1*H*-indole-2-carboxylic acid ethyl ester.

¹HNMR (300 MHz, DMSO-d₆) δ 12.10 (s, br, 1H, NH), 7.94 (s, 1H), 7.64 (dd, *J* = 1.35 & 9.15 Hz, 1H), 7.62 (d, *J* = 16.2 Hz, H-vinyl), 7.44 (d, *J* = 9.3 Hz, 1H), 7.16 (s, 1H), 6.41 (d, *J* = 16.2 Hz, 1H, H-vinyl), 4.33 (q, *J* = 7.1 Hz, 2H, OCH₂CH₃), 1.47 (s, 9H, 3xCH₃), 1.33 (t, *J* = 7.1 Hz, 3H, OCH₂CH₃).

MS-EI 315 [M⁺].

A solution of 5-(2-*tert*-butoxycarbonyl-vinyl)-1*H*-indole-2-carboxylic acid ethyl ester (1.6 g, 5 mmol) in ethyl acetate (40 mL) and methanol (80 mL) was hydrogenated over 5% palladium on carbon (0.2 g) at room temperature for overnight. The catalyst was filtered off and the filtrate was concentrated to give 1.6 g (99%) of 5-(2-*tert*-butoxycarbonyl-ethyl)-1*H*-indole-2-carboxylic acid ethyl ester as a white solid. The solid was used in the next step without further purification.

To a solution of 5-(2-*tert*-butoxycarbonyl-ethyl)-1*H*-indole-2-carboxylic acid ethyl ester (1.6 g, 5 mmol) in dichloromethane (12 mL) was added 8 mL of 40% trifluoroacetic acid. The mixture was stirred at room temperature for 1 hour. The reaction was concentrated, dissolved in toluene, concentrate, dissolved in dichloromethane and concentrated to give 1.3 g (99%) of 5-(2-carboxy-ethyl)-1*H*-indole-2-carboxylic acid ethyl ester.

To a solution of 5-(2-carboxy-ethyl)-1*H*-indole-2-carboxylic acid ethyl ester (1.3 g, 5 mmol) in 10 mL of dimethylformamide, was added 1,1'-carbonyldiimidazole (973 mg, 6 mmol). The mixture was stirred under nitrogen at room temperature for 30 mins. To the reaction mixture was added pyrrolidine (1.3 mL, 15 mmol) dropwise. After stirring at room temperature for overnight, the reaction was diluted with dichloromethane (300 mL), washed with 1N hydrochloric acid, saturated sodium bicarbonate solution, brine, dried and concentrated to give 1.5 g (95%) of 5-(3-oxo-3-pyrrolidin-1-yl-propyl)-1*H*-indole-2-carboxylic acid ethyl ester.

¹HNMR (360 MHz, DMSO-d₆) δ 11.69 (s, br, 1H, NH), 7.45 (s, 1H), 7.34 (d, *J* = 8.45 Hz, 1H), 7.14 (dd, *J* = 1.87 & 8.45 Hz, 1H), 7.04 (m, 1H), 4.32 (q, *J* = 7.18 Hz, 2H, OCH₂CH₃), 3.31 (t, *J* = 6.55 Hz, 2H, CH₂), 3.26 (t, *J* = 6.93 Hz, 2H, CH₂), 2.87 (t, *J* = 7.69 Hz, 2H, CH₂), 2.53 (t, *J* = 7.61 Hz, 2H, CH₂), 1.69-1.80 (m, 4H, 2xCH₂), 1.32 (t, *J* = 7.18 Hz, 2H, OCH₂CH₃).

To a solution of 5-(3-oxo-3-pyrrolidin-1-yl-propyl)-1*H*-indole-2-carboxylic acid ethyl ester (1.5 g, 4.8 mmol) in 100 mL of tetrahydrofuran, lithium aluminum hydride powder (728 mg, 19.2 mmol) was added. The mixture was stirred under nitrogen at room temperature for 2 hour. To the reaction mixture was added water (0.75 mL), 15% sodium hydroxide (0.75 mL) and water (0.75 mL). It was then

filtered through a pad of celite and the filtrate was concentrated. The residue was washed with ethyl acetate to give 960 mg (78%) of [5-(3-pyrrolidin-1-yl-propyl)-1*H*-indol-2-yl]-methanol.

¹HNMR (360 MHz, DMSO-*d*₆) was added 10.78 (br s, 1H, NH), 7.22 (s, 1H),
 5 7.20 (d, *J* = 8.05 Hz, 1H), 6.85 (dd, *J* = 1.27 & 8.05 Hz, 1H), 6.16 (d, *J* = 1.27 Hz, 1H), 5.11 (t, *J* = 5.53 Hz, 1H, OH), 4.56 (d, *J* = 5.17 Hz, 2H, CH₂OH), 2.61 (t, *J* = 7.73 Hz, 2H, CH₂), 2.34-2.38 (m, 6H, 3xCH₂), 1.73 (m, 2H, CH₂), 1.65 (m, 4H, 2xCH₂).

Manganese (IV) oxide (4.7 g, 54 mmol) was added to a mixture of [5-(3-pyrrolidin-1-yl-propyl)-1*H*-indol-2-yl]-methanol (940 mg, 3.6 mmol) in
 10 dichloromethane (150 mL) and toluene (150 mL). The mixture was stirred at room temperature for overnight. The reaction was filtered through a pad of celite and the filtrate was concentrated to give 670 mg (72%) of 5-(3-pyrrolidin-1-yl-propyl)-1*H*-indole-2-carbaldehyde.

¹HNMR (300 MHz, DMSO-*d*₆) δ 11.85 (br s, 1H, NH), 9.79 (s, 1H, CHO),
 15 7.50 (s, 1H), 7.35 (d, *J* = 8.6 Hz, 1H), 7.29 (d, *J* = 1.6 Hz, 1H), 7.18 (d, *J* = 1.6 & 8.6 Hz, 1H), 2.65 (t, *J* = 7.8 Hz, 2H, CH₂), 2.33-2.38 (m, 6H, 3xCH₂), 1.74 (m, 2H, CH₂), 1.65 (m, 4H, 2xCH₂).

MS-EI 256 [M⁺].

20

EXAMPLE 7: General condensation of oxindoles and aldehydes

A mixture of 1 eq. of the oxindole, 1 eq. of the aldehyde and 1–3 eq. of piperidine (or pyrrolidine) in ethanol (0.4 M) was heated at 80-100 °C until the reaction was complete. The mixture was cooled to room temperature and the solid
 25 was collected by vacuum filtration, washed with ethanol and dried to give the product. Where solid was not obtained from the reaction mixture, the mixture was concentrated and purified by column chromatography.

EXAMPLE 8: COMPOUND IN-001

3-[5-(3-Diethylamino-propyl)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

A mixture of oxindole (26 mg, 0.19 mmol), 5-(3-diethylamino-propyl)-1*H*-indole-2-carbaldehyde (50 mg, 0.19 mmol) and piperidine(0.1 mL) in ethanol (1 mL) was heated in a sealed tube at 95 °C for 20 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 13 mg (18%) of the title compound as an orange crystalline solid.

¹HNMR (300 MHz, DMSO-*d*₆) δ 12.88 (s, br, 1H, NH), 11.02 (s, br, 1H, NH), 7.90 (s, 1H, H-vinyl), 7.71 (d, *J* = 7.5 Hz, 1H), 7.44-7.49 (m, 2H), 7.20 (m, 1H), 7.13 (dd, *J* = 1.4 & 8.5 Hz, 1H), 7.0-7.04 (m, 2H), 6.89 (d, *J* = 7.8 Hz, 1H), 2.63 (t, *J* = 7.5 Hz, 2H, CH₂), 2.34-2.44 (m, 6H, 3xCH₂), 1.67-1.72 (m, 2H, CH₂), 0.91 (t, *J* = 7.0 Hz, 6H, 2x NCH₂CH₃).

MS-EI *m/z* 373 [M⁺].

EXAMPLE 9: COMPOUND IN-002

5-Bromo-3-[5-(3-diethylamino-propyl)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

A mixture of 5-bromo-2-oxindole (41 mg, 0.19 mmol), 5-(3-diethylamino-propyl)-1*H*-indole-2-carbaldehyde (50 mg, 0.19 mmol) and piperidine(0.1 mL) in ethanol (1 mL) was heated in a sealed tube at 95 °C for 20 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 52 mg (59%) of the title compound as a red-orange solid.

¹HNMR (300 MHz, DMSO-*d*₆) δ 12.80 (s, br, 1H, NH), 11.17 (s, br, 1H, NH), 8.06 (s, 1H, H-vinyl), 7.97 (d, *J* = 1.7 Hz, 1H), 7.47-7.51 (m, 2H), 7.35 (dd, *J* = 1.7 & 8.2 Hz, 1H), 7.15 (m, 1H), 7.05 (s, 1H), 6.85 (d, *J* = 8.2 Hz, 1H), 2.53-2.68 (m, 8H, 4xCH₂), 1.75-1.79 (m, 2H, CH₂), 0.97 (t, *J* = 7.0 Hz, 6H, 2x NCH₂CH₃).

MS-EI *m/z* 451 and 453 [M⁺-1 and M⁺+1].

EXAMPLE 10: COMPOUND IN-0033-[5-(3-Diethylamino-propyl)-1*H*-indol-2-ylmethylene]-6-phenyl-1,3-dihydro-indol-2-one

A mixture of 6-phenyl-2-oxindole (41 mg, 0.19 mmol) 5-(3-diethylamino-propyl)-1*H*-indole-2-carbaldehyde (50 mg, 0.19 mmol) and piperidine(0.1 mL) in ethanol (1 mL) was heated in a sealed tube at 95 °C for 20 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 49 mg (56%) of the title compound as an orange crystalline solid.

¹HNMR (300 MHz, DMSO-d₆) δ 12.85 (s, br, 1H, NH), 11.08 (s, br, 1H, NH), 7.92 (s, 1H, H-vinyl), 7.79 (d, *J* = 1.7 Hz, 1H), 7.65 (d, *J* = 7.3 Hz, 2H), 7.44-7.49 (m, 4H), 7.32-7.38 (m, 2H), 7.13-7.15 (m, 2H), 7.06 (s, 1H), 2.64 (t, *J* = 7.4 Hz, 2H, CH₂), 2.36-2.46 (m, 6H, 3xCH₂), 1.67-1.75 (m, 2H, CH₂), 0.92 (t, *J* = 7.2 Hz, 6H, 2x NCH₂CH₃).

MS-EI *m/z* 449 [M⁺].

EXAMPLE 11: COMPOUND IN-0043-[5-(3-Diethylamino-propyl)-1*H*-indol-2-ylmethylene]-5-phenyl-1,3-dihydro-indol-2-one

A mixture of 5-phenyl-2-oxindole (41 mg, 0.19 mmol), 5-(3-diethylamino-propyl)-1*H*-indole-2-carbaldehyde (50 mg, 0.19 mmol) and piperidine(0.1 mL) in ethanol (1 mL) was heated in a sealed tube at 95 °C for 20 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 10 mg (11%) of the title compound as an orange crystalline solid.

MS-EI *m/z* 449 [M⁺].

EXAMPLE 12: COMOPUND IN-0053-[5-(2-dimethylamino-ethoxy)-1*H*-indol-2-ylmethylene]-5-phenyl-1,3-dihydro-indol-2-one

A mixture of 5-phenyl-2-oxindole (41 mg, 0.2 mmol), 5-(2-dimethylamino-ethoxy)-1*H*-indole-2-carbaldehyde (46 mg, 0.2 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 100 °C for 2 hours. The precipitate was collected by

vacuum filtration, washed with ethanol and dried to give 58 mg (68%) of the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.87 (s, br, 1H, NH), 11.06 (s, br, 1H, NH), 8.08 (s, 2H), 7.70 (d, *J* = 7.2 Hz, 2H), 7.44-7.53 (m, 4H), 7.33 (m, 1H), 7.16 (s, 1H), 6.92-7.0 (m, 3H), 4.06 (t, *J* = 5.8 Hz, 2H, OCH₂CH₂N), 2.65 (t, *J* = 5.8 Hz, 2H, OCH₂CH₂N), 2.23 (s, 6H, N(CH₃)₂.)
MS-EI *m/z* 423 [M⁺].

EXAMPLE 13: COMPOUND IN-006

10 5-Phenyl-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

A mixture of 5-phenyl-2-oxindole (23 mg, 0.11 mmol), 5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indole-2-carbaldehyde (29 mg, 0.11 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 100 °C for 2 hours. The precipitate was collected by
15 vacuum filtration, washed with ethanol and dried to give 30 mg (61%) of the title compound as an orange solid.

¹HNMR (360 MHz, DMSO-d₆) δ 12.87 (s, br, 1H, NH), 11.05 (s, br, 1H, NH), 8.08 (d, *J* = 1.4 Hz, 2H), 7.70 (d, *J* = 7.6 Hz, 2H), 7.44-7.53 (m, 4H), 7.31-7.35 (m, 1H), 7.15 (d, *J* = 1.8 Hz, 1H), 6.91-7.03 (m, 3H), 4.07 (t, *J* = 6.0 Hz, 2H, OCH₂CH₂N), 2.80 (t, *J* = 6.0 Hz, 2H, OCH₂CH₂N), 2.49 (m, under DMSO H-pyrrolidine), 1.68 (m, 4H, H-pyrrolidine).
20

MS-EI *m/z* 451 [M⁺+2].

EXAMPLE 14: COMPOUND IN-007

25 3-[5-(2-Morpholin-4-yl-ethoxy)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

A mixture of 2-oxindole (29 mg, 0.22 mmol), 5-(2-morpholin-4-yl-ethoxy)-1*H*-indole-2-carbaldehyde (60 mg, 0.22 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 100 °C for 2 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 81 mg (95%) of the title compound
30 as a yellow solid.

¹HNMR (360 MHz, DMSO-d₆) δ 12.87 (s, 1H, NH), 10.97 (s br, 1H, NH), 7.88 (s, 1H, H-vinyl), 7.71 (d, *J* = 7.6 Hz, 1H), 7.48 (m, 1H), 7.21 (t, *J* = 7.4 Hz, 1H), 7.13 (m, 1H), 7.03 (m, 2H), 6.89-6.95 (m, 2H), 4.10 (t, *J* = 5.8 Hz, 2H, OCH₂CH₂N), 3.58 (t, *J* = 4.5 Hz, 4H, 2x CH₂), 2.71 (t, *J* = 5.8 Hz, 2H, OCH₂CH₂N), 2.49 (m, under DMSO, 2x CH₂).

MS-EI *m/z* 389 [M⁺].

EXAMPLE 15: COMPOUND IN-008

5-Bromo-3-[5-(2-morpholin-4-yl-ethoxy)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

A mixture of 5-bromo-2-oxindole (46 mg, 0.22 mmol), 5-(2-morpholin-4-yl-ethoxy)-1*H*-indole-2-carbaldehyde (60 mg, 0.22 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 100 °C for 2 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 65 mg (64%) of the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.78 (s, 1H, NH), 11.02 (s br, 1H, NH), 8.03 (s, 1H, H-vinyl), 7.96 (m, 1H), 7.49 (d, *J* = 8.6 Hz, 1H), 7.35 (dd, *J* = 1.4 & 8.6 Hz, 1H), 7.15 (m, 1H), 7.04 (s, 1H), 6.93-6.96 (m, 1H), 6.85 (d, *J* = 7.9 Hz, 1H), 4.09 (t, *J* = 5.6 Hz, 2H, OCH₂CH₂N), 3.58 (t, *J* = 4.3 Hz, 4H, 2x CH₂), 2.71 (t, *J* = 5.6 Hz, 2H, OCH₂CH₂N), 2.49 (m, under DMSO, 2x CH₂).

MS-EI *m/z* 468 [M⁺].

EXAMPLE 16: COMPOUND IN-009

3-[5-(2-Morpholin-4-yl-ethoxy)-1*H*-indol-2-ylmethylene]-6-phenyl-1,3-dihydro-indol-2-one

A mixture of 6-phenyl-2-oxindole (44 mg, 0.21 mmol), 5-(2-morpholin-4-yl-ethoxy)-1*H*-indole-2-carbaldehyde (58 mg, 0.21 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 100 °C for 2 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 70 mg (71%) of the title compound as an orange solid.

¹HNMR (300 MHz, DMSO-d₆) δ 12.86 (s, br, 1H, NH), 11.11 (s, br, 1H, NH), 7.93 (s, 1H, H-vinyl), 7.79 (d, *J* = 8.1 Hz, 1H), 7.65 (d, *J* = 7.2 Hz, 2H), 7.44-7.51 (m, 3H), 7.35 (m, 2H), 7.12-7.15 (m, 2H), 7.04 (s, 1H), 6.93 (dd, *J* = 2.1 & 8.1 Hz, 1H), 4.09 (t, *J* = 5.8 Hz, 2H, OCH₂CH₂N), 3.58 (t, *J* = 4.6 Hz, 4H, 2x CH₂), 2.71 (t, *J* = 5.8 Hz, 2H, OCH₂CH₂N), 2.49 (m, under DMSO, 2x CH₂).

MS-EI *m/z* 465 [M⁺].

EXAMPLE 17: COMPOUND IN-010

3-[5-(2-Dimethylamino-ethoxy)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

A mixture of oxindole (28 mg, 0.216 mmol), 5-(2-dimethylamino-ethoxy)-1*H*-indole-2-carbaldehyde (50 mg, 0.21 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 100 °C for 2 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 59 mg (79%) of the title compound as an orange solid.

¹HNMR (300 MHz, DMSO-d₆) δ 12.88 (s, br, 1H, NH), 11.01 (s, br, 1H, NH), 7.89 (s, 1H, H-vinyl), 7.71 (d, *J* = 7.2 Hz, 1H), 7.48 (d, *J* = 8.7 Hz, 1H), 7.20 (dt, *J* = 0.9 & 7.5 Hz, 1H), 7.13 (d, *J* = 2.1 Hz, 1H), 7.0-7.05 (m, 2H), 6.88-6.93 (m, 2H), 4.05 (t, *J* = 5.8 Hz, 2H, OCH₂CH₂N), 2.63 (t, *J* = 5.8 Hz, 2H, OCH₂CH₂N), 2.22 (s, 6H, N(CH₃)₂).

MS-EI *m/z* 347 [M⁺].

EXAMPLE 18: COMPOUND IN-011

5-Bromo-3-[5-(2-dimethylamino-ethoxy)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

A mixture of 5-bromo-2-oxindole (46 mg, 0.215 mmol), 5-(2-dimethylamino-ethoxy)-1*H*-indole-2-carbaldehyde (50 mg, 0.21 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 100 °C for 2 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 72 mg (78%) of the title compound as a red solid.

¹HNMR (300 MHz, DMSO-d₆) δ 12.79 (s, br, 1H, NH), 11.10 (s, br, 1H, NH), 8.05 (s, 1H, H-vinyl), 7.97 (d, *J* = 1.2 Hz, 1H), 7.50 (d, *J* = 9.3 Hz, 1H), 7.35

(dd, $J = 1.5$ & 8.1 Hz, 1H), 7.15 (d, $J = 2.1$ Hz, 1H), 7.03 (s, 1H), 6.93 (dd, $J = 2.3$ & 8.8 Hz, 1H), 6.85 (d, $J = 8.7$ Hz, 1H), 4.04 (t, $J = 5.9$ Hz, 2H, $\text{OCH}_2\text{CH}_2\text{N}$), 2.63 (t, $J = 5.9$ Hz, 2H, $\text{OCH}_2\text{CH}_2\text{N}$), 2.22 (s, 6H, $\text{N}(\text{CH}_3)_2$).

MS-EI m/z 426 [M^+].

5

EXAMPLE 19: COMPOUND IN-012

3-[5-(2-Dimethylamino-ethoxy)-1H-indol-2-ylmethylene]-6-phenyl-1,3-dihydro-indol-2-one

A mixture of 6-phenyl-2-oxindole (45 mg, 0.21 mmol), 5-(2-dimethylamino-ethoxy)-1H-indole-2-carbaldehyde (50 mg, 0.21 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 100°C for 2 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 68 mg (76%) of the title compound as a solid.

^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 12.86 (s, br, 1H, NH), 11.12 (s, br, 1H, NH), 7.94 (s, 1H, H-vinyl), 7.80 (d, $J = 8.7$ Hz, 1H), 7.66 (d, $J = 7.2$ Hz, 2H), 7.44-7.51 (m, 3H), 7.33-7.39 (m, 2H), 7.12-7.15 (m, 2H), 7.04 (s, 1H), 6.92 (dd, $J = 2.1$ & 8.7 Hz, 1H), 4.05 (t, $J = 5.9$ Hz, 2H, $\text{OCH}_2\text{CH}_2\text{N}$), 2.63 (t, $J = 5.9$ Hz, 2H, $\text{OCH}_2\text{CH}_2\text{N}$), 2.22 (s, 6H, $\text{N}(\text{CH}_3)_2$).

MS-EI m/z 423 [M^+].

20

EXAMPLE 20: COMPOUND IN-013

3-[5-(2-Pyrrolidin-1-yl-ethoxy)-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

A mixture of oxindole (26.6mg, 0.2 mmol), 5-(2-pyrrolidin-1-yl-ethoxy)-1H-indole-2-carbaldehyde (52 mg, 0.2 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 100°C for 2 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 61 mg (81%) of the title compound.

^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 12.88 (s, br, 1H, NH), 11.01 (s, br, 1H, NH), 7.89 (s, 1H), 7.71 (d, $J = 7.5$ Hz, 1H), 7.48 (d, $J = 9.3$ Hz, 1H), 7.20 (dt, $J = 1.2$ & 7.6 Hz, 1H), 7.12 (d, $J = 2.4$ Hz, 1H), 7.0-7.05 (m, 2H), 6.88-6.93 (m, 2H), 4.06 (t, $J = 5.8$ Hz, 2H, $\text{OCH}_2\text{CH}_2\text{N}$), 2.79 (t, $J = 5.8$ Hz, 2H, $\text{OCH}_2\text{CH}_2\text{N}$), 2.52 (m, under DMSO H-pyrrolidine), 1.68 (m, 4H, H-pyrrolidine).

MS-EI m/z 373 [M^+].

EXAMPLE 21: COMPOUND IN-014

5-Bromo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

A mixture of 5-bromo-2-oxindole (42 mg, 0.2 mmol), 5-(2-pyrrolidin-1-yl-ethoxy)-1H-indole-2-carbaldehyde (52 mg, 0.2 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 100 °C for 2 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 72 mg (80%) of the title compound.

^1H NMR (300 MHz, DMSO- d_6) δ 12.79 (s, br, 1H, NH), 11.12 (s, br, 1H, NH), 8.05 (s, 1H), 7.97 (d, J = 1.9 Hz, 1H), 7.50 (d, J = 9.0 Hz, 1H), 7.35 (dd, J = 1.9 & 8.3 Hz, 1H), 7.14 (d, J = 2.1 Hz, 1H), 7.03 (s, 1H), 6.93 (dd, J = 2.1 & 9.0 Hz, 1H), 6.85 (d, J = 8.3 Hz, 1H), 4.06 (t, J = 6.0 Hz, 2H, $\text{OCH}_2\text{CH}_2\text{N}$), 2.79 (t, J = 6.0 Hz, 2H, $\text{OCH}_2\text{CH}_2\text{N}$), 2.50 (m, under DMSO H-pyrrolidine), 1.67 (m, 4H, H-pyrrolidine).

MS-EI m/z 451/453 [M^+].

EXAMPLE 22: COMPOUND IN-015

6-Phenyl-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

A mixture of 6-phenyl-2-oxindole (41 mg, 0.2 mmol), 5-(2-pyrrolidin-1-yl-ethoxy)-1H-indole-2-carbaldehyde (50 mg, 0.2 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 100 °C for 2 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 65 mg (72%) of the title compound as a red solid.

^1H NMR (300 MHz, DMSO- d_6) δ 12.86 (s, br, 1H, NH), 11.12 (s, br, 1H, NH), 7.94 (s, 1H), 7.80 (d, J = 8.6 Hz, 1H), 7.66 (d, J = 7.5 Hz, 2H), 7.44-7.51 (m, 3H), 7.33-7.39 (m, 2H), 7.12-7.14 (m, 2H), 7.05 (s, 1H), 6.92 (dd, J = 1.9 & 8.6 Hz, 1H), 4.07 (t, J = 5.8 Hz, 2H, $\text{OCH}_2\text{CH}_2\text{N}$), 2.80 (t, J = 5.8 Hz, 2H, $\text{OCH}_2\text{CH}_2\text{N}$), 2.50 (m, under DMSO H-pyrrolidine), 1.68 (m, 4H, H-pyrrolidine).

MS-EI m/z 449 [M^+].

EXAMPLE 23: COMPOUND IN-016

3-[5-(2-Diethylamino-ethoxy)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

A mixture of oxindole (26 mg, 0.2 mmol), 5-(2-diethylamino-ethoxy)-1*H*-indole-2-carbaldehyde (50 mg, 0.2 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 100 °C for 2 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 50 mg (67%) of the title compound.

¹HNMR (360 MHz, DMSO- d_6) δ 12.87 (s, br, 1H, NH), 10.98 (s, br, 1H, NH), 7.89 (s, 1H), 7.71 (d, J = 9.3 Hz, 1H), 7.48 (d, J = 8.6 Hz, 1H), 7.21 (m, 1H), 7.13 (d, J = 2.2 Hz, 1H), 7.03 (m, 2H), 6.89-6.92 (m, 2H), 4.02 (t, J = 5.9 Hz, 2H, OCH₂CH₂N), 2.79 (t, J = 5.9 Hz, 2H, OCH₂CH₂N), 2.56 (q, J = 7.2 Hz, 4H, 2x NCH₂CH₃), 0.98 (t, J = 7.2 Hz, 6H, 2x NCH₂CH₃).

MS-EI m/z 375 [M^+].

EXAMPLE 24: COMPOUND IN-017

5-Bromo-3-[5-(2-diethylamino-ethoxy)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

A mixture of 5-bromo-2-oxindole (41 mg, 0.2 mmol), 5-(2-diethylamino-ethoxy)-1*H*-indole-2-carbaldehyde (50 mg, 0.2 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 100 °C for 2 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 60 mg (66%) of the title compound.

¹HNMR (360 MHz, DMSO- d_6) δ 12.78 (s, br, 1H, NH), 11.09 (s, br, 1H, NH), 8.04 (s, 1H), 7.96 (d, J = 1.7 Hz, 1H), 7.49 (d, J = 8.9 Hz, 1H), 7.35 (dd, J = 1.7 & 8.2 Hz, 1H), 7.14 (d, J = 2.2 Hz, 1H), 7.03 (s, 1H), 6.93 (dd, J = 2.2 & 8.9 Hz, 1H), 6.85 (d, J = 8.2 Hz, 1H), 4.02 (t, J = 6.1 Hz, 2H, OCH₂CH₂N), 2.79 (t, J = 6.0 Hz, 2H, OCH₂CH₂N), 2.55 (q, J = 7.2 Hz, 4H, 2x NCH₂CH₃), 0.98 (t, J = 7.2 Hz, 6H, 2x NCH₂CH₃).

EXAMPLE 25: COMPOUND IN-018

3-[5-(2-Diethylamino-ethoxy)-1*H*-indol-2-ylmethylene]-6-phenyl-1,3-dihydro-indol-2-one

A mixture of 6-phenyl-2-oxindole (40.5 mg, 0.2 mmol), 5-(2-diethylamino-ethoxy)-1*H*-indole-2-carbaldehyde (50 mg, 0.2 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 100 °C for 2 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 65 mg (72%) of the title compound.

¹HNMR (360 MHz, DMSO-*d*₆) δ 12.85 (s, br, 1H, NH), 11.08 (s, br, 1H, NH), 7.92 (s, 1H), 7.80 (d, *J* = 7.9 Hz, 1H), 7.66 (d, *J* = 7.9 Hz, 2H), 7.45-7.49 (m, 3H), 7.33-7.36 (m, 2H), 7.13 (m, 2H), 7.05 (s, 1H), 6.92 (dd, 1H), 4.03 (t, *J* = 6.0 Hz, 2H, OCH₂CH₂N), 2.80 (t, *J* = 6.0 Hz, 2H, OCH₂CH₂N), 2.56 (q, *J* = 7.2 Hz, 4H, 2x NCH₂CH₃), 0.98 (t, *J* = 7.2 Hz, 6H, 2x NCH₂CH₃).

MS-EI *m/z* 451 [M⁺].

EXAMPLE 26: COMPOUND IN-019

3-[5-(2-Diethylamino-ethoxy)-1*H*-indol-2-ylmethylene]-5-phenyl-1,3-dihydro-indol-2-one

A mixture of 5-phenyl-2-oxindole (39 mg, 0.18 mmol), 5-(2-diethylamino-ethoxy)-1*H*-indole-2-carbaldehyde (48 mg, 0.18 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 100 °C for 2 hours. The precipitate was collected by vacuum filtration, washed with ethanol and dried to give 44 mg (54%) of the title compound as a red solid.

¹HNMR (360 MHz, DMSO-*d*₆) δ 12.87 (s, br, 1H, NH), 11.05 (s, br, 1H, NH), 8.09 (d, *J* = 2.2 Hz, 2H), 7.70 (d, *J* = 8.3 Hz, 2H), 7.44-7.53 (m, 4H), 7.33-7.35 (m, 1H), 7.15 (s, 1H), 7.03 (s, 1H), 6.98 (d, *J* = 8.3 Hz, 1H), 6.92 (dd, *J* = 2.2 & 9.0 Hz, 1H), 4.03 (t, *J* = 6.1 Hz, 2H, OCH₂CH₂N), 2.80 (t, *J* = 6.1 Hz, 2H, OCH₂CH₂N), 2.56 (q, *J* = 7.2 Hz, 4H, 2x NCH₂CH₃), 0.98 (t, *J* = 7.2 Hz, 6H, 2x NCH₂CH₃).

EXAMPLE 27: COMPOUND IN-020

3-[5-(3-Pyrrolidin-1-yl-propyl)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

A mixture of oxindole (27 mg, 0.2 mmol), 5-(3-pyrrolidin-1-yl-propyl)-1*H*-indole-2-carbaldehyde (51 mg, 0.2 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated in a sealed tube at 80 °C for 3 hours. The precipitate was collected by vacuum filtration, washed with ethanol/ethyl acetate and dried to give 55 mg (74%) of the title compound.

¹H NMR (300 MHz, DMSO-*d*₆) δ 12.89 (br s, 1H, NH), 11.03 (s, 1H, NH), 7.90 (s, 1H), 7.71 (d, *J* = 7.45 Hz, 1H), 7.48 (d, *J* = 8.41 Hz, 1H), 7.44 (s, 1H), 7.21 (t, 1H), 7.13 (d, 1H), 7.0-7.04 (m, 2H), 6.89 (d, *J* = 7.6 Hz, 1H), 2.66 (t, *J* = 7.6 Hz, 2H, CH₂), 2.35-2.40 (m, 6H, 3xCH₂), 1.75 (m, 2H, CH₂), 1.65 (m, 4H, 2xCH₂). MS-EI 371 [M⁺].

EXAMPLE 28: COMPOUND IN-021

2-Oxo-3-[5-(3-pyrrolidin-1-yl-propyl)-1*H*-indol-2-ylmethylene]-2,3-dihydro-1*H*-indole-5-carboxylic acid

A mixture of 5-carboxy-2-oxindole (35 mg, 0.2 mmol), 5-(3-pyrrolidin-1-yl-propyl)-1*H*-indole-2-carbaldehyde (51 mg, 0.2 mmol) and piperidine (0.2 mL) in ethanol (1 mL) was heated in a sealed tube at 80 °C for 6 hours. The reaction mixture was concentrated and the residue was dissolved in methanol, made acidic with 1 N hydrochloric acid until solid was formed. The precipitate was filtered, washed with 1 N hydrochloric acid, water, ethanol and dried to give the title compound.

¹H NMR (360 MHz, DMSO-*d*₆) δ 12.75 (br s, 1H, NH), 11.37 (s, 1H, NH), 10.67 (br s, 1H, COOH), 8.32 (s, 1H), 8.14 (s, 1H), 7.85 (dd, *J* = 1.26 & 8.23 Hz, 1H), 7.51-7.55 (m, 2H), 7.15-7.19 (m, 2H), 7.0 (d, *J* = 8.6 Hz, 1H), 3.50 (m, 2H, CH₂), 3.09 (m, 2H, CH₂), 2.94 (m, 2H, CH₂), 2.73 (t, *J* = 7.6 Hz, 2H, CH₂), 1.84-2.07 (m, 6H, 3xCH₂). MS 416.4 [M⁺+1].

EXAMPLE 29: COMPOUND IN-021

2-Oxo-3-[5-(3-pyrrolidin-1-yl-propyl)-1H-indol-2-ylmethylene]-2,3-dihydro-1H-indole-5-carboxylic acid

A mixture of 5-carboxy-2-oxindole (35 mg, 0.2 mmol), 5-(3-diethylamino-propyl)-1H-indole-2-carbaldehyde (51 mg, 0.2 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 80 °C for 6 hours. The reaction was acidified with 1 N HCl and the precipitate was collected by vacuum filtration, washed with 1 N HCl, water and ethanol, dried to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.75 (s, 1H, NH), 11.37 (s, 1H, NH), 10.67 (br s, 1H, COOH), 8.32 (s, 1H), 8.14 (s, 1H), 7.85 (dd, *J* = 1.26 & 8.23 Hz, 1H), 7.51-7.55 (m, 2H), 7.15-7.19 (m, 2H), 7.0 (d, *J* = 8.6 Hz, 1H), 3.50 (m, 2H), 3.09 (m, 2H), 2.94 (m, 2H), 2.73 (t, *J* = 7.6 Hz, 2H), 1.84-2.07 (m, 6H, 3xCH₂).

MS *m/z* 416 [M⁺+1].

EXAMPLE 30: COMPOUND IN-022

2-Oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1H-indol-2-ylmethylene]-2,3-dihydro-1H-indole-5-carboxylic acid

2-Oxo-2,3-dihydro-1H-indole-5-carboxylic acid was condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1H-indole-2-carbaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.74 (s, 1H, NH), 11.42 (br s, 1H, NH), 8.31 (s, 1H), 8.12 (s, 1H), 7.84 (dd, 1H), 7.53 (d, 1H), 7.2 (3, 1H), 7.14 (m, 1H), 7.0 (m, 2H), 4.26 (m, 2H, CH₂), 2.94 (m, 2H, CH₂), 2.49 (m, under DMSO), 1.67 (m, 4H, 2xCH₂).

MS-Ve APCI *m/z* 416.5 [M⁺-1].

EXAMPLE 31: COMPOUND IN-023

2-Oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1H-indol-2-ylmethylene]-2,3-dihydro-1H-indole-6-carboxylic acid

2-Oxo-2,3-dihydro-1H-indole-6-carboxylic acid was condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1H-indole-2-carbaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.86 (s, 1H, NH), 11.11 (br s, 1H, NH), 8.01 (s, 1H, H-vinyl), 7.78 (d, *J* = 8 Hz, 1H), 7.64 (dd, *J* = 1.5 & 8 Hz, 1H), 7.51 (d, *J* = 9 Hz, 1H), 7.43 (d, 1H), 7.15 (d, *J* = 2 Hz, 1H), 7.10 (s, 1H), 4.11 (t, *J* = 6 Hz, 2H, CH₂), 2.89 (t, *J* = 6 Hz, 2H, CH₂), 2.61 (m, 4H, 2xCH₂), 1.71 (m, 4H, 2xCH₂).

5 MS-Ve APC1 m/z 416.5 [M⁺-1].

EXAMPLE 32: COMPOUND IN-024

4-(2-Hydroxy-ethyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

10 4-(2-Hydroxy-ethyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indole-2-carbaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 13.03 (s, 1H, NH), 10.98 (br s, 1H, NH), 7.76 (s, 1H, H-vinyl), 7.48 (d, *J* = 9 Hz, 1H), 7.12 (m, 3H), 6.92 (dd, *J* = 2.5 & 9 Hz, 1H), 6.86 (d, *J* = 8 Hz, 1H), 6.77 (d, *J* = 8 Hz, 1H), 4.83 (t, *J* = 5.4 Hz, 2H, OH),
15 4.07 (t, *J* = 6 Hz, 2H, CH₂), 3.74 (m, 2H, CH₂), 3.12 (m, 2H, CH₂), 2.80 (t, *J* = 6 Hz, 2H, CH₂), 2.52 (m, 4H, 2xCH₂), 1.68 (m, 4H, 2xCH₂).

MS-Ve APCI 416.6.[M⁺-1].

EXAMPLE 33: COMPOUND IN-025

20 6-Pyridin-3-yl-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

6-Pyridin-3-yl-1,3-dihydro-indol-2-one was condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indole-2-carbaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.85 (s, 1H, NH), 11.13 (s, 1H, NH), 8.88
25 (d, *J* = 2 Hz, 1H), 8.56 (dd, 1H), 8.06 (m, 1H), 7.95 (s, 1H, H-vinyl), 7.83 (d, *J* = 8 Hz, 1H), 7.48 (m, 2H), 7.40 (dd, 1H), 7.17 (br s, 1H), 7.13 (d, 1H), 7.06 (br s, 1H), 6.93 (dd, 1H), 4.07 (t, *J* = 6 Hz, 2H, CH₂), 2.80 (t, *J* = 6 Hz, 2H, CH₂), 2.50 (m, 4H, 2xCH₂), 1.68 (m, 4H, 2xCH₂).

MS-Ve APCI m/z 449.5 [M⁺-1]

30

EXAMPLE 34: COMPOUND IN-0266-(4-Methoxy-phenyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

6-(4-Methoxy-phenyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indole-2-carbaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.86 (s, 1H, NH), 11.03 (br s, 1H, NH), 7.85 (s, 1H, H-vinyl), 7.73 (d, *J* = 8 Hz, 1H), 7.59 (m, 2H), 7.48 (d, *J* = 9 Hz, 1H), 7.28 (dd, *J* = 1.5 & 8 Hz, 1H), 7.12 (d, *J* = 2 Hz, 1H), 7.08 (d, *J* = 1 Hz, 1H), 7.02 (m, 3H), 6.92 (dd, *J* = 2 & 9 Hz, 1H), 4.07 (t, *J* = 6 Hz, 2H, CH₂), 3.80 (s, 3H, OCH₃), 2.80 (t, *J* = 6 Hz, 2H, CH₂), 2.53 (m, 4H, 2xCH₂), 1.68 (m, 4H, 2xCH₂).

MS-Ve APCI m/z 478.5 [M⁺-1]

EXAMPLE 35: COMPOUND IN-027

6-(3-Methoxy-phenyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

6-(3-Methoxy-phenyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indole-2-carbaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.85 (s, 1H, NH), 11.05 (s, 1H, NH), 7.91 (s, 1H, H-vinyl), 7.78 (d, *J* = 8 Hz, 1H), 7.49 (d, *J* = 9 Hz, 1H), 7.33-7.39 (m, 2H), 7.21 (br d, *J* = 8 Hz, 1H), 7.17 (m, 1H), 7.13 (br s, 2H), 7.04 (s, 1H), 6.91-6.95 (m, 2H), 4.07 (t, *J* = 6 Hz, 2H, CH₂), 3.82 (s, 3H, CH₃), 2.80 (t, *J* = 6 Hz, 2H, CH₂), 2.51 (m, 4H, 2xCH₂), 1.68 (m, 4H, 2xCH₂).

MS-Ve APCI m/z 478.7 [M⁺-1].

EXAMPLE 36: COMPOUND IN-028

6-(2-Methoxy-phenyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

6-(2-Methoxy-phenyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indole-2-carbaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.88 (s, 1H, NH), 10.97 (s, 1H, NH), 7.88 (s, 1H, H-vinyl), 7.73 (d, *J* = 8 Hz, 1H), 7.48 (d, *J* = 9 Hz, 1H), 7.30-7.34 (m, 2H),

7.10-7.14 (m, 3H), 7.05 (m, 3H), 6.93 (dd, $J = 2$ & 9 Hz, 1H), 4.08 (t, $J = 6$ Hz, 2H, CH₂), 3.78 (s, 3H, OCH₃), 2.80 (t, $J = 6$ Hz, 2H, CH₂), 2.51 (m, 4H, 2xCH₂), 1.68 (m, 4H, 2xCH₂).

MS-Ve APCI m/z 478.7 [$M^+ - 1$].

5

EXAMPLE 37: COMPOUND IN-029

6-(4-Fluoro-phenyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

6-(4-Fluoro-phenyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indole-2-carbaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.84 (s, 1H, NH), 11.08 (s, 1H, NH), 7.91 (s, 1H, H-vinyl), 7.78 (d, $J = 8$ Hz, 1H), 7.69 (m, 2H), 7.49 (d, $J = 9$ Hz, 1H), 7.26-7.33 (m, 3H), 7.13 (d, $J = 2$ Hz, 1H), 7.10 (d, $J = 1.4$ Hz, 1H), 7.04 (s, 1H), 6.93 (dd, $J = 2$ & 9 Hz, 1H), 4.07 (t, $J = 6$ Hz, 2H, CH₂), 2.80 (t, $J = 6$ Hz, 2H, CH₂), 2.53 (m, 4H, 2xCH₂), 1.68 (m, 4H, 2xCH₂).

MS-Ve APCI m/z 466.6 [$M^+ - 1$].

EXAMPLE 38: COMPOUND IN-030

3-[5-(2-Morpholin-4-yl-ethoxy)-1*H*-indol-2-ylmethylene]-2-oxo-2,3-dihydro-1*H*-indole-5-carboxylic acid

2-Oxo-2,3-dihydro-1*H*-indole-5-carboxylic acid was condensed with 5-(2-morpholin-4-yl-ethoxy)-1*H*-indole-2-carbaldehyde to give the title compound.

EXAMPLE 39: COMPOUND IN-031

3-[5-(2-Morpholin-4-yl-ethoxy)-1*H*-indol-2-ylmethylene]-2-oxo-2,3-dihydro-1*H*-indole-6-carboxylic acid

2-Oxo-2,3-dihydro-1*H*-indole-6-carboxylic acid was condensed with 5-(2-morpholin-4-yl-ethoxy)-1*H*-indole-2-carbaldehyde to give the title compound.

EXAMPLE 40: COMPOUND IN-032

3-[5-(2-Morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-5-phenyl-1,3-dihydro-indol-2-one

5-Phenyl-1,3-dihydro-indol-2-one was condensed with 5-(2-morpholin-4-yl-ethoxy)-1*HH*-indole-2-carbaldehyde to give the title compound.

EXAMPLE 41: COMPOUND IN-033

4-(2-Hydroxy-ethyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

4-(2-Hydroxy-ethyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-morpholin-4-yl-ethoxy)-1*HH*-indole-2-carbaldehyde to give the title compound.

EXAMPLE 42: COMPOUND IN-034

3-[5-(2-Morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-6-pyridin-3-yl-1,3-dihydro-indol-2-one

6-Pyridin-3-yl-1,3-dihydro-indol-2-one was condensed with 5-(2-morpholin-4-yl-ethoxy)-1*HH*-indole-2-carbaldehyde to give the title compound.

EXAMPLE 43: COMPOUND IN-035

6-(4-Methoxy-phenyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

6-(4-Methoxy-phenyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-morpholin-4-yl-ethoxy)-1*HH*-indole-2-carbaldehyde to give the title compound.

EXAMPLE 44: COMPOUND IN-036

6-(3-Methoxy-phenyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

6-(3-Methoxy-phenyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-morpholin-4-yl-ethoxy)-1*HH*-indole-2-carbaldehyde to give the title compound.

EXAMPLE 45: COMPOUND IN-037

6-(2-Methoxy-phenyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

6-(2-Methoxy-phenyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-morpholin-4-yl-ethoxy)-1H-indole-2-carbaldehyde to give the title compound.

EXAMPLE 46: COMPOUND IN-038

6-(4-Fluoro-phenyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

6-(4-Fluoro-phenyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-morpholin-4-yl-ethoxy)-1H-indole-2-carbaldehyde to give the title compound.

EXAMPLE 47: COMPOUND IN-039

2-Oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1H-indol-2-ylmethylene]-2,3-dihydro-1H-indole-5-sulfonic acid amide

A mixture of 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid amide (28 mg, 0.13 mmol), 5-(2-pyrrolidin-1-yl-ethoxy)-1H-indole-2-carbaldehyde (34 mg, 0.13 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 90 °C for 2 hours. The reaction was cooled at 0 °C for overnight. The precipitate was collected by vacuum filtration, washed with cold ethanol and dried to give 47 mg (80%) of the title compound as an orange solid.

¹HNMR (360 MHz, DMSO-d₆) δ 12.72 (s, 1H, NH), 11.15 (br s, 1H, NH), 8.17 (br s, 1H), 8.07 (s, 1H, H-vinyl), 7.69 (br dd, 1H), 7.51 (d, *J* = 8.84 Hz, 1H), 7.15 (m, 4H), 7.05 (d, *J* = 8.45 Hz, 1H), 6.95 (br dd, 1H), 4.08 (t, *J* = 5.8 Hz, 2H, CH₂), 2.80 (t, *J* = 5.8 Hz, 2H, CH₂), 2.53 (m, 4H, 2xCH₂), 1.68 (m, 4H, 2xCH₂).

MS-EI *m/z* 452 [M⁺].

EXAMPLE 48: COMPOUND IN-0403-[5-(2-Morpholin-4-yl-ethoxy)-1*H*-indol-2-ylmethylene]-2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid amide

A mixture of 2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid amide (27 mg, 0.12 mmol), 5-(2-morpholin-4-yl-ethoxy)-1*H*-indole-2-carbaldehyde (30 mg, 0.11 mmol) and piperidine (0.1 mL) in ethanol (1 mL) was heated at 90 °C for 2 hours. The reaction was cooled at 0 °C for overnight. The precipitate was collected by vacuum filtration, washed with cold ethanol and dried to give 38 mg (62%) of the title compound as an orange solid.

¹HNMR (360 MHz, DMSO-d₆) δ 12.72 (s, 1H, NH), 11.31 (br s, 1H, NH), 8.17 (br s, 1H), 8.07 (s, 1H, H-vinyl), 7.69 (br dd, 1H), 7.51 (d, *J* = 8.9 Hz, 1H), 7.16 (m, 4H), 7.05 (d, *J* = 8.4 Hz, 1H), 6.95 (br dd, 1H), 4.1 (t, *J* = 5.7 Hz, 2H, CH₂), 3.58 (m, 4H, 2xCH₂), 2.71 (t, *J* = 5.7 Hz, 2H, CH₂), 2.49 (m, 4H, 2xCH₂).

MS-Ve APCI m/z 465.7 [M⁺1].

EXAMPLE 49: COMPOUND IN-0412-Oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indol-2-ylmethylene]-2,3-dihydro-1*H*-indole-5-sulfonic acid methylamide

2-Oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid methylamide was condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indole-2-carbaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.71 (s, 1H, NH), 11.37 (br s, 1H, NH), 8.14 (br s, 2H), 7.63 (dd, *J* = 2 & 8 Hz, 1H), 7.51 (d, *J* = 9 Hz, 1H), 7.22 (m, 1H, CH₃NH), 7.15 (br s, 2H), 7.08 (d, *J* = 8 Hz, 1H), 6.95 (dd, *J* = 2 & 9 Hz, 1H), 4.08 (t, *J* = 6 Hz, 2H, CH₂), 2.80 (t, *J* = 6 Hz, 2H, CH₂), 2.55 (m, 4H, 2xCH₂), 2.44 (d, *J* = 5 Hz, 3H, CH₃), 1.67 (m, 4H, 2xCH₂).

MS-Ve APCI m/z 465.7 [M⁺-1].

EXAMPLE 50: COMPOUND IN-0422-Oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1H-indol-2-ylmethylene]-2,3-dihydro-1H-indole-5-sulfonic acid dimethylamide

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid dimethylamide was condensed
 5 with 5-(2-pyrrolidin-1-yl-ethoxy)-1H-indole-2-carbaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.72 (s, 1H, NH), 11.41 (br s, 1H, NH), 8.25 (s, 1H, H-vinyl), 8.14 (d, *J* = 2 Hz, 1H), 7.59 (dd, *J* = 2 & 8 Hz, 1H), 7.52 (d, *J* = 9 Hz, 1H), 7.17 (d, *J* = 2 Hz, 1H), 7.13 (br s, 1H), 7.11 (d, *J* = 8 Hz, 1H), 6.96 (dd, *J* = 2 & 9 Hz, 1H), 4.08 (t, *J* = 6 Hz, 2H, CH₂), 2.81 (t, *J* = 6 Hz, 2H, CH₂), 2.55 (m, 4H, 2xCH₂), 1.68 (m, 4H, 2xCH₂).

MS-Ve APCI *m/z* 479.8 [M⁺-1].

EXAMPLE 51: COMPOUND IN-043

15 2-Oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1H-indol-2-ylmethylene]-2,3-dihydro-1H-indole-5-sulfonic acid isopropylamide

Oxo-2,3-dihydro-1H-indole-5-sulfonic acid isopropylamide was condensed
 with 5-(2-pyrrolidin-1-yl-ethoxy)- 2-1H-indole-2-carbaldehyde to give the title compound.

20 ¹HNMR (360 MHz, DMSO-d₆) δ 12.72 (s, 1H, NH), 11.35 (br s, 1H, NH), 8.15 (d, *J* = 2 Hz, 1H), 7.66 (dd, *J* = 2 & 8 Hz, 1H), 7.51 (d, *J* = 9 Hz, 1H), 7.34 (d, *J* = 7 Hz, 1H, (CH₃)₂CHNH), 7.15 (m, 2H), 7.06 (d, *J* = 8 Hz, 1H), 6.95 (dd, *J* = 2 & 9 Hz, 1H), 4.08 (t, *J* = 6 Hz, 2H, CH₂), 3.27 (m, 1H, (CH₃)₂CH), 2.80 (t, *J* = 6 Hz, 2H, CH₂), 2.55 (m, 4H, 2xCH₂), 1.68 (m, 4H, 2xCH₂), 0.97 (d, *J* = 6.5 Hz, 6H, 2xCH₃).

25 MS-Ve APCI *m/z* 493.8 [M⁺-1].

EXAMPLE 52: COMPOUND IN-044

2-Oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1H-indol-2-ylmethylene]-2,3-dihydro-1H-indole-5-sulfonic acid phenylamide

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid phenylamide was condensed
5 with 5-(2-pyrrolidin-1-yl-ethoxy)-1H-indole-2-carbaldehyde to give the title
compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.68 (s, 1H, NH), 11.36 (br s, 1H, NH),
10.09 (br s, 1H, NH), 8.14 (d, *J* = 2 Hz, 1H), 8.08 (s, 1H, H-vinyl), 7.60 (dd, *J* = 2 &
8.5 Hz, 1H), 7.50 (d, *J* = 9 Hz, 1H), 7.21 (t, 2H), 7.11-7.15 (m, 4H), 7.01 (m, 2H),
10 6.95 (dd, *J* = 2 & 9 Hz, 1H), 4.08 (t, *J* = 6 Hz, 2H, CH₂), 2.81 (t, *J* = 6 Hz, 2H, CH₂),
2.54 (m, 4H, 2xCH₂), 1.68 (m, 4H, 2xCH₂).

EXAMPLE 53: COMPOUND IN-045

15 2-Oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1H-indol-2-ylmethylene]-2,3-dihydro-1H-indole-5-sulfonic acid pyridin-3-ylamide

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid pyridin-3-ylamide was
condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1H-indole-2-carbaldehyde to give the
title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.68 (s, 1H, NH), 11.38 (br s, 1H, NH),
20 8.30 (d, *J* = 2.5 Hz, 1H), 8.19 (dd, *J* = 1.4 & 5 Hz, 1H), 8.16 (d, *J* = 2 Hz, 1H), 8.11
(s, 1H, H-vinyl), 7.60 (dd, *J* = 2 & 8 Hz, 1H), 7.50 (m, 2H), 7.25 (dd, *J* = 5 & 8 Hz,
1H), 7.16 (br s, 2H), 7.02 (d, *J* = 8 Hz, 1H), 6.96 (dd, *J* = 2.5 & 9 Hz, 1H), 4.09 (t, *J*
= 6 Hz, 2H, CH₂), 2.85 (t, *J* = 6 Hz, 2H, CH₂), 2.57 (m, 4H, 2xCH₂), 1.70 (m, 4H,
2xCH₂).

25 MS-Ve APCI m/z 528.8 [M⁺-1].

EXAMPLE 54: COMPOUND IN-046

5-(2,3-Dihydro-indole-1-sulfonyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

5 5-(2,3-Dihydro-indole-1-sulfonyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indole-2-carbaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.67 (s, 1H, NH), 11.36 (br s, 1H, NH), 8.25 (d, *J* = 2 Hz, 1H), 8.21 (s, 1H, H-vinyl), 7.62 (dd, *J* = 2 & 8 Hz, 1H), 7.51 (dd, *J* = 3 & 8 Hz, 2H), 7.12-7.21 (m, 4H), 6.92-7.01 (m, 3H), 6.92-7.01 (m, 3H), 4.08 (t, *J* = 6 Hz, 2H, CH₂), 3.96 (t, *J* = 8 Hz, 2H, CH₂), 2.93 (t, *J* = 8 Hz, 2H, CH₂), 2.81 (t, *J* = 6 Hz, 2H, CH₂), 2.54 (m, 4H, 2xCH₂), 1.68 (m, 4H, 2xCH₂).

MS -Ve APCI m/z 553.8 [M⁺-1].

EXAMPLE 55: COMPOUND IN-047

15 2-Oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indol-2-ylmethylene]-2,3-dihydro-1*H*-indole-5-sulfonic acid (3-chloro-phenyl)-amide

2-Oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid (3-chloro-phenyl)-amide was condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1*H*-indole-2-carbaldehyde to give the title compound.

20 ¹HNMR (360 MHz, DMSO-d₆) δ 12.67 (s, 1H, NH), 11.39 (br s, 1H, NH), 8.18 (d, *J* = 2 Hz, 1H), 8.12 (s, 1H, H-vinyl), 7.62 (dd, *J* = 2 & 8 Hz, 1H), 7.51 (d, *J* = 8 Hz, 1H), 7.24 (t, *J* = 8 Hz, 1H), 7.16 (m, 2H), 7.13 (m, 1H), 7.08 (m, 1H), 7.01-7.03 (m, 2H), 6.96 (dd, *J* = 2 & 9 Hz, 1H), 4.08 (t, *J* = 6 Hz, 2H, CH₂), 2.83 (t, *J* = 6 Hz, 2H, CH₂), 2.56 (m, 4H, 2xCH₂), 1.69 (m, 4H, 2xCH₂).

25 MS-Ve APCI m/z 562 [M⁺-1].

EXAMPLE 56: COMPOUND IN-048

2-Oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1H-indol-2-ylmethylene]-2,3-dihydro-1H-indole-5-sulfonic acid (3-chloro-phenyl)-methyl-amide

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid (3-chloro-phenyl)-methyl-
5 amide was condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1H-indole-2-carbaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.69 (s, 1H, NH), 11.43 (br s, 1H, NH), 8.18 (s, 1H, H-vinyl), 8.03 (d, *J* = 2 Hz, 1H), 7.52 (d, *J* = 9 Hz, 1H), 7.35 (m, 2H), 7.24-7.29 (m, 2H), 7.11-7.17 (m, 3H), 7.02 (d, *J* = 8 Hz, 1H), 6.96 (dd, *J* = 2.5 & 9 Hz, 1H), 4.08 (t, *J* = 6 Hz, 2H, CH₂), 3.27 (s, 3H, CH₃), 2.81 (t, *J* = 6 Hz, 2H, CH₂), 2.53 (m, 4H, 2xCH₂), 1.67 (m, 4H, 2xCH₂).

MS -Ve APCI m/z 576 [M⁺-1].

EXAMPLE 57: COMPOUND IN-049

15 2-Oxo-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1H-indol-2-ylmethylene]-2,3-dihydro-1H-indole-5-sulfonic acid (4-chloro-2-fluoro-phenyl)-amide

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid (4-chloro-2-fluoro-phenyl)-amide was condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1H-indole-2-carbaldehyde to give the title compound.

20 ¹HNMR (360 MHz, DMSO-d₆) δ 12.70 (s, 1H, NH), 11.36 (br s, 1H, NH), 8.09 (d, *J* = 2 Hz, 1H), 8.07 (s, 1H, H-vinyl), 7.57 (dd, *J* = 2 & 8 Hz, 1H), 7.51 (d, *J* = 9 Hz, 1H), 7.31 (dd, *J* = 2 & 10 Hz, 1H), 7.24 (d, *J* = 9 Hz, 1H), 7.14 (m, 3H), 7.01 (d, *J* = 8 Hz, 2H), 6.96 (dd, *J* = 2 & 9 Hz, 1H), 4.11 (t, *J* = 6 Hz, 2H, CH₂), 2.92 (t, *J* = 6 Hz, 2H, CH₂), 2.66 (m, 4H, 2xCH₂), 1.72 (m, 4H, 2xCH₂).

25 MS-Ve APCI m/z 567.7 [M⁺-1].

EXAMPLE 58: COMPOUND IN-0505-(3,4-Dihydro-2H-quinoline-1-sulfonyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

5 5-(3,4-Dihydro-2H-quinoline-1-sulfonyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1H-indole-2-carbaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.67 (s, 1H, NH), 11.38 (br s, 1H, NH), 8.14 (s, 1H, H-vinyl), 8.11 (d, *J* = 2 Hz, 1H), 7.63 (d, *J* = 8 Hz, 1H), 7.51 (d, *J* = 9 Hz, 1H), 7.37 (dd, *J* = 2 & 8 Hz, 1H), 7.13-7.19 (m, 3H), 7.06 (m, 2H), 6.94-6.99 (m, 2H), 4.08 (t, *J* = 6 Hz, 2H, CH₂), 3.80 (t, *J* = 6 Hz, 2H, CH₂), 3.25 (m, 2H, CH₂), 2.81 (t, *J* = 6 Hz, 2H, CH₂), 2.52 (m, 4H, 2xCH₂), 2.48 (m, 2H, CH₂), 1.67 (m, 4H, 2xCH₂).

MS-Ve APCI m/z 580 [M⁺-1].

15 EXAMPLE 59: COMPOUND IN-0515-(3,4-Dihydro-1H-isoquinoline-2-sulfonyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

5-(3,4-Dihydro-1H-isoquinoline-2-sulfonyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1H-indole-2-carbaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.71 (s, 1H, NH), 11.39 (br s, 1H, NH), 8.22 (s, 1H, H-vinyl), 8.20 (d, *J* = 1.4 Hz, 1H), 7.67 (dd, *J* = 1.4 & 8 Hz, 1H), 7.51 (d, *J* = 9 Hz, 1H), 7.07-7.17 (m, 7H), 6.95 (dd, *J* = 2.5 & 9 Hz, 1H), 4.23 (s, 2H, CH₂), 4.08 (t, *J* = 6 Hz, 2H, CH₂), 3.33 (t, *J* = 6 Hz, 2H, CH₂), 2.86 (t, *J* = 6 Hz, 2H, CH₂), 2.81 (t, *J* = 6 Hz, 2H, CH₂), 2.52 (m, 4H, 2xCH₂), 1.68 (m, 4H, 2xCH₂).

MS-Ve APCI m/z 567.7 [M⁺-1].

EXAMPLE 60: COMPOUND IN-052

5-(5-Bromo-2,3-dihydro-indole-1-sulfonyl)-3-[5-(2-pyrrolidin-1-yl-ethoxy)-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

5 5-(5-Bromo-2,3-dihydro-indole-1-sulfonyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-pyrrolidin-1-yl-ethoxy)-1H-indole-2-carbaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.67 (s, 1H, NH), 11.41 (br s, 1H, NH), 8.25 (d, *J* = 2 Hz, 1H), 8.23 (s, 1H, H-vinyl), 7.63 (dd, *J* = 2 & 9 Hz, 1H), 7.51 (d, *J* = 9 Hz, 1H), 7.45 (d, *J* = 9 Hz, 1H), 7.36 (d, *J* = 2 Hz, 1H), 7.33 (s, 1H), 7.17 (d, *J* = 2.5 Hz, 1H), 7.14 (s, 1H), 7.02 (d, *J* = 8 Hz, 1H), 6.96 (dd, *J* = 3 & 9 Hz, 1H), 4.08 (t, *J* = 6 Hz, 2H, CH₂), 3.97 (t, *J* = 9 Hz, 2H, CH₂), 2.95 (t, *J* = 9 Hz, 2H, CH₂), 2.81 (t, *J* = 6 Hz, 2H, CH₂), 2.53 (m, 4H, 2xCH₂), 1.68 (m, 4H, 2xCH₂).

MS-Ve APCI 632.0 and 634.2.

15 EXAMPLE 61: COMPOUND IN-053

3-[5-(2-Morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid methylamide

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid methylamide was condensed with 5-(2-morpholin-4-yl-ethoxy)-1H-indole-2-carbaldehyde to give the title compound.

EXAMPLE 62: COMPOUND IN-054

3-[5-(2-Morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid dimethylamide

25 2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid dimethylamide was condensed with 5-(2-morpholin-4-yl-ethoxy)-1H-indole-2-carbaldehyde to give the title compound.

EXAMPLE 63: COMPOUND IN-055

3-[5-(2-Morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid isopropylamide

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid isopropylamide was condensed with 5-(2-morpholin-4-yl-ethoxy)-1*HH*-indole-2-carbaldehyde to give the title compound.

EXAMPLE 64: COMPOUND IN-056

3-[5-(2-Morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid phenylamide

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid phenylamide was condensed with 5-(2-morpholin-4-yl-ethoxy)-1*HH*-indole-2-carbaldehyde to give the title compound.

EXAMPLE 65: COMPOUND IN-057

3-[5-(2-Morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid pyridin-3-ylamide

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid pyridin-3-ylamide was condensed with 5-(2-morpholin-4-yl-ethoxy)-1*HH*-indole-2-carbaldehyde to give the title compound.

EXAMPLE 66: COMPOUND IN-058

5-(2,3-Dihydro-indole-1-sulfonyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

5-(2,3-Dihydro-indole-1-sulfonyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-morpholin-4-yl-ethoxy)-1*HH*-indole-2-carbaldehyde to give the title compound.

EXAMPLE 67: COMPOUND IN-059

3-[5-(2-Morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid (3-chloro-phenyl)-amide

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid (3-chloro-phenyl)-amide was condensed with 5-(2-morpholin-4-yl-ethoxy)-1*HH*-indole-2-carbaldehyde to give the title compound.

5 EXAMPLE 68: COMPOUND IN-060

3-[5-(2-Morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid (3-chloro-phenyl)-methyl-amide

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid (3-chloro-phenyl)-methyl-amide was condensed with 5-(2-morpholin-4-yl-ethoxy)-1*HH*-indole-2-carbaldehyde to give the title compound.

EXAMPLE 69: COMPOUND IN-061

3-[5-(2-Morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid (4-chloro-2-fluoro-phenyl)-amide

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid (4-chloro-2-fluoro-phenyl)-amide was condensed with 5-(2-morpholin-4-yl-ethoxy)-1*HH*-indole-2-carbaldehyde to give the title compound.

EXAMPLE 70: COMPOUND IN-062

5-(3,4-Dihydro-2H-quinoline-1-sulfonyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

5-(3,4-Dihydro-2H-quinoline-1-sulfonyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-morpholin-4-yl-ethoxy)-1*HH*-indole-2-carbaldehyde to give the title compound.

EXAMPLE 71: COMPOUND IN-063

5-(3,4-Dihydro-1H-isoquinoline-2-sulfonyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

5-(3,4-Dihydro-1H-isoquinoline-2-sulfonyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-morpholin-4-yl-ethoxy)-1*HH*-indole-2-carbaldehyde to give the title compound.

EXAMPLE 72: COMPOUND IN-064

5-(5-Bromo-2,3-dihydro-indole-1-sulfonyl)-3-[5-(2-morpholin-4-yl-ethoxy)-1H-indol-2-ylmethylene]-1,3-dihydro-indol-2-one

5-(5-Bromo-2,3-dihydro-indole-1-sulfonyl)-1,3-dihydro-indol-2-one was condensed with 5-(2-morpholin-4-yl-ethoxy)-1H-indole-2-carbaldehyde to give the title compound.

EXAMPLE 73: COMPOUND IN-065

3-(1H-Indol-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid amide

10 5-Aminosulfonyl-2-oxindole was condensed with indole-3-carboxaldehyde to give the title compound.

¹HNMR (300 MHz, DMSO-d₆) δ 12.03 (br s, 1H, NH), 10.80 (br s, 1H, NH), 9.37 (s, 1H), 8.09-8.24 (m, 3H), 7.51 (m, 1H), 7.41 (m, 1H), 7.13 (m, 2H), 7.03 (s, 2H, NH₂), 6.85 (d, *J* = 7.8 Hz, 1H).

15 MS-EI *m/z* 339 [M⁺].

EXAMPLE 74: COMPOUND IN-066

3-(2-Methyl-1H-indol-3-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid amide

20 5-Aminosulfonyl-2-oxindole was condensed with 2-methylindole-3-carboxaldehyde to give the title compound.

MS-EI 353 [M⁺].

EXAMPLE 75: COMPOUND IN-067

25 3-(1H-Indol-5-ylmethylene)-2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid amide

5-Aminosulfonyl-2-oxindole was condensed with indole-5-carbaldehyde to give the title compound.

MS-EI 339 [M⁺].

09871700-060401

EXAMPLE 76: COMPOUND IN-068

3-(1*H*-Indol-3-ylmethylene)-2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid
methanamide

5 5-Methanaminosulfonyl-2-oxindole was condensed with indole-3-
carboxaldehyde to give the title compound.

¹HNMR (300 MHz, DMSO-*d*₆) δ 12.03 (br s, 1H, NH), 10.84 (br s, 1H, NH),
9.39 (s, 1H), 8.25 (s, 1H, H-vinyl), 8.22 (d, *J* = 1.5 Hz, 1H), 8.16 (m, 1H), 7.40-7.46
(m, 2H), 7.06-7.15 (m, 2H), 7.04 (m, 1H, CH₃NH), 6.88 (d, *J* = 7.8 Hz, 1H), 2.32 (d,
J = 5.1 Hz, 3H, CH₃).

10 MS-EI 353 [M⁺].

EXAMPLE 77: COMPOUND IN-069

3-(2-Methyl-1*H*-indol-3-ylmethylene)-2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid
methanamide

15 5-Methanaminosulfonyl-2-oxindole was condensed with 2-methylindole-3-
carboxaldehyde to give the title compound.

EXAMPLE 78: COMPOUND IN-070

3-(1*H*-Indol-5-ylmethylene)-2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid
methanamide

20 5-Methanaminosulfonyl-2-oxindole was condensed with indole-5-
carbaldehyde to give the title compound.

¹HNMR (300 MHz, DMSO-*d*₆) δ 11.38 (br s, 1H, NH), 10.85 (br s, 1H, NH),
8.11 (m, 1H), 8.01 (m, 1H), 7.90 (m, 1H), 7.78 (m, 1H), 7.37-7.53 (m, 3H), 7.10 (m,
25 1H), 6.95 (m, 1H), 6.4 (m, 1H), 2.28 (s, 3H, CH₃).

MS-EI 353 [M⁺].

EXAMPLE 79: COMPOUND IN-071

3-(1*H*-Indol-2-ylmethylene)-2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid
methanamide

5-Methylaminosulfonyl-2-oxindole was condensed with indole-2-
5 carbaldehyde to give the title compound.

¹HNMR (300 MHz, DMSO-d₆) δ 12.75 (br s, 1H, NH), 11.40 (br s, 1H, NH),
8.15 (s, 1H, H-vinyl), 8.10 (d, *J* = 1.5 Hz, 1H), 7.54-7.65 (m, 3H), 7.20-7.27 (m,
3H), 7.02-7.06 (m, 2H), 2.38 (d, *J* = 4.8 Hz, 3H, CH₃).

MS-EI 353 [M⁺].

EXAMPLE 80: COMPOUND IN-072

3-(1*H*-Indol-3-ylmethylene)-2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid
dimethanamide

5-Dimethylaminosulfonyl-2-oxindole was condensed with indole-3-
15 carboxaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.11 (br s, 1H, NH), 10.94 (br s, 1H, NH),
9.50 (s, 1H), 8.44 (s, 1H), 8.33 (m, 2H), 7.51-7.53 (m, 2H), 7.25 (m, 2H), 7.03 (d, *J*
= 7.9 Hz, 1H), 2.63 (s, 6H, 2xCH₃).

MS-EI 367 [M⁺].

EXAMPLE 81: COMPOUND IN-073

3-(2-Methyl-1*H*-indol-3-ylmethylene)-2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid
dimethanamide

5-Dimethylaminosulfonyl-2-oxindole was condensed with 2-methylindole-3-
25 carboxaldehyde to give the title compound.

¹HNMR (300 MHz, DMSO-d₆) δ 11.95 (br s, 1H, NH), 10.86 (br s, 1H, NH),
7.84 (s, 1H, H-vinyl), 7.41 (dd, *J* = 1.8 & 8.4 Hz, 1H), 7.33 (d, *J* = 8.4 Hz, 1H), 7.04
(m, 1H), 6.85-6.95 (m, 4H).

MS-EI 381 [M⁺].

3-(1*H*-Indol-5-ylmethylene)-2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid
dimethylamide

5-Dimethylaminosulfonyl-2-oxindole was condensed with indole-5-
5 carbaldehyde to give the title compound.

MS-EI 367 [M⁺].

EXAMPLE 83: COMPOUND IN-075

10 3-(1*H*-Indol-2-ylmethylene)-2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid
dimethylamide

5-Dimethylaminosulfonyl-2-oxindole was condensed with indole-2-carbaldehyde to give the title compound.

¹HNMR (360 MHz, DMSO-d₆) δ 12.80 (br s, 1H, NH), 11.44 (br s, 1H, NH), 8.30 (s, 1H, H-vinyl), 8.15 (d, *J* = 2.2 Hz, 1H), 7.70 (d, *J* = 7.2 Hz, 1H), 7.60 (m, 2H), 7.30 (m, 1H), 7.23 (s, 1H), 7.07-7.12 (m, 2H), 2.63 (s, 6H, 2xCH₃).

MS-EI 367 [M⁺].

EXAMPLE 84: COMPOUND IN-076

20 3-(4-Methoxy-1*H*-indol-2-ylmethylene)-2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic
acid methylamide

A mixture of 2-oxo-2,3-dihydro-1*H*-indole-5-sulfonic acid methylamide (194 mg, 0.86 mmol), 4-methoxy-1*H*-indole-2-carbaldehyde (150 mg, 0.86 mmol) and piperidine (36 mg, 0.43 mmol) in ethanol (0.2 M) was stirred at room temperature for a total of 5 days. The reaction was concentrated to ¼ of its volume and the precipitate was collected by vacuum filtration to give the title compound as a pale orangish-red solid.

¹H NMR (360 MHz, DMSO-d₆) δ 12.82 (s, 1H, NH), 11.36 (br s, 1H, NH), 8.11 (s, 1H-H-vinyl), 8.11 (d, *J* = 1.3 Hz, 1H), 7.63 (dd, *J* = 1.3 & 8.5 Hz, 1H), 7.15-7.27 (m, 4H), 7.08 (d, *J* = 7.96 Hz, 1H), 6.56 (d, *J* = 7.7 Hz, 1H), 3.92 (s, 3H, OCH₃), 2.43 (d, *J* = 5.02 Hz, 3H, NCH₃).

MS m/z 384 $[M^++1]$.

Oxindole Preparation (Oxindoles Used in the Combinatorial Library)EXAMPLE 85:5-(Pyrrolidine-1-sulfonyl)-1,3-dihydro-indol-2-one

5 A suspension of 5-chlorosulfonyl-2-oxindole (1.62 g, 7 mmol), pyrrolidine (0.701 mL, 8.4 mmol) and pyridine (1 mL) in dichloromethane (20 mL) was stirred at room temperature for 4 hours. The reaction mixture was diluted with ethyl acetate (300 mL) and made acidic with 1 N hydrochloric acid (16 mL). The organic layer was then washed with sodium bicarbonate and brine, dried and concentrated. The
 10 residue was washed with ethanol (3 mL). It was then purified by chromatography on silica gel eluting with methanol:dichloromethane 1:9 to give 0.432 g (27%) 5-(pyrrolidine-1-sulfonyl)-1,3-dihydro-indol-2-one.

¹H NMR (360 MHz, DMSO-d₆) δ 10.77 (s, 1H, NH), 7.6-7.64 (m, 2H), 6.97 (d, *J* = 8 Hz, 1H), 3.58 (s, 2H, H-3), 3.07-3.11 (m, 4H, c-pentyl), 1.63-1.66 (m, 4H, c-pentyl). MS *m/z* 266 [M]⁺
 15

EXAMPLE 86:2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid benzylamide

A suspension of 5-chlorosulfonyl-2-oxindole (1.62 g, 7 mmol), benzylamine
 20 (0.918 mL, 8.4 mmol) and pyridine (1 mL) in dichloromethane (20 mL) was stirred at room temperature for 4 hours. The reaction mixture was diluted with ethyl acetate (300 mL) and acidified with 1 N hydrochloric acid (16 mL). The organic layer was then washed with sodium bicarbonate and brine, dried and concentrated. The residue was washed with ethanol (3 mL) and was then purified by chromatography
 25 on silica gel eluting with methanol:dichloromethane 1:9 to give 1.4 g (66%) of 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid benzylamide.

¹H NMR (360 MHz, DMSO-d₆) δ 0.70 (s, 1H, NH-1), 7.91 (t, *J* = 6 Hz, 1H, NH₂SO₂-), 6.63 (dd, *J* = 2 and 8 Hz, 1H, Ar-H), 7.57 (s, 1H), 7.20-7.29 (m, 5H, Ar-H), 6.92 (d, *J* = 8.5 Hz, 1H), 3.94 (d, *J* = 6 Hz, 2H, CH₂NSO₂-), 3.54 (s, 2H, H-3).
 30

EXAMPLE 87:

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid 4-fluoro-benzylamide

A suspension of 5-chlorosulfonyl-2-oxindole (5.0g), 4-fluorobenzylamine (3.0 mL) and pyridine (3.5 mL) in dichloromethane (30 mL) was stirred at room temperature for 4 hours. The precipitate was filtered, washed with dichloromethane and dried to give 5.8 g (84%) of 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid 4-fluoro-benzylamide.

¹H NMR (360 MHz, DMSO-d₆) δ 10.74 (s, 1H, NH-1), 7.50-7.55 (m, 2H, H-4, 6), 7.21-7.26 (m, 4H, Ar-H), 7.04-7.09 (m, 1H, SO₂NH), 6.90-6.92 (d, 1H, H-7), 3.91-3.93 (d, 2H, NHCH₂), 3.53 (s, 2H, H-3). MS *m/z* 320

EXAMPLE 88:

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid (3-chloro-phenyl)-amide

A suspension of 5-chlorosulfonyl-2-oxindole (5 g), 3-chloroaniline (2.74 mL) and pyridine (3.5 mL) in dichloromethane (30 mL) was stirred at room temperature for overnight. The precipitate was filtered, washed with dichloromethane and dried to give 5.2 g (74%) 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid (3-chloro-phenyl)-amide.

¹H NMR (360 MHz, DMSO-d₆) δ 10.74 (s, br, 1H, NH), 10.41 (s, 1H, SO₂NH), 7.59-7.64 (m, 2H, H-4,6), 7.22-7.26 (m, 1H, Ar-H), 7.10-7.11 (m, 1H, Ar-H), 7.03-7.07 (m, 2H, Ar-H), 6.9-6.92 (d, 1H, H-7), 3.54 (s, 3H, CH₃). MS *m/z* 322 (M⁺).

EXAMPLE 89:

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid (2-methoxy-phenyl)-amide

A solution of 5-chlorosulfonyl-2-oxindole (3 g), o-anisidine (1.8 mL) and pyridine (2.1 mL) in dichloromethane (20 mL) was stirred at room temperature for overnight at which time the red color solid was present. The solid was filtered, washed with ethanol and dried under vacuum to yield 1.5 g (37%) of 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid (2-methoxy-phenyl)-amide.

¹H NMR (360 MHz, DMSO-d₆) δ 10.71 (s, 1H, NH-1), 9.19 (s, 1H, SO₂NH), 7.51-7.54 (m, 2H, H-4, 6), 7.18-7.20 (m, 1H, Ar-H), 7.04-7.09 (m, 1H, Ar-H), 6.78-6.89 (m, 3H, H-7, Ar-H), 3.44-3.56 (m, 5H, H-3, OCH₃). MS *m/z* (APCI-) 317.2

5 EXAMPLE 90:

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid pyridin-3-ylamide

A solution of 5-chlorosulfonyl-2-oxindole (3 g) and 3-aminopyridine (1.46g) in pyridine (15 mL) was stirred at room temperature for overnight at which time the brown color solid was present. The precipitate was filtered, washed with ethanol
10 and dried under vacuum to yield 1.4 g (38%) of 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid pyridin-3-ylamide.

¹H NMR (360 MHz, DMSO-d₆) δ 10.74 (s, 1H, NH-1), 10.39 (s, 1H, SO₂NH), 8.27-8.28 (d, 1H, Ar-H), 8.21-8.23 (m, 1H, Ar-H), 7.59-7.62 (m, 2H, H-4, 6), 7.44-7.68 (m, 1H, Ar-H), 7.24-7.28 (m, 1H, Ar-H), 6.69-6.71 (d, 1H, H-7), 3.54
15 (s, 2H, H-3). MS *m/z* (APCI+) 290.2

EXAMPLE 91:

2-Oxo-2,3-dihydro-1H-indole-5-sulfonic acid (2-methoxy-ethyl)-amide

A solution of 5-chlorosulfonyl-2-oxindole (5 g), 2-methoxyethylamine (2.25 mL) and pyridine (7 mL) in dichloromethane (30 mL) was stirred at room
20 temperature for 4 hours. The reaction was concentrated and dichloromethane (15 mL) was added. The precipitate was filtered, washed with dichloromethane and dried to give 1.9 g (33%) of 2-oxo-2,3-dihydro-1H-indole-5-sulfonic acid (2-methoxy-ethyl)-amide.

¹H NMR (360 MHz, DMSO-d₆) δ 10.72 (s, br, 1H, NH), 7.63 (dd, *J* = 2 and 8 Hz, 1H), 7.59 (s, 1H), 7.48 (t, *J* = 6 Hz, 1H, NH), 6.93 (d, *J* = 8 Hz, 1H), 3.57 (s, 2H), 3.29 (t, *J* = 6 Hz, 2H, CH₂), 3.16 (s, 3H, OCH₃), 2.82-2.87 (q, *J* = 6 Hz, 2H, CH₂). MS *m/z* 270 M⁺
25

EXAMPLE 92:5-Aminosulfonyl-2-oxindole

To a 100 mL flask charged with 27 mL of chlorosulfonic acid, 13.3 g of 2-oxindole was added slowly. The reaction temperature was maintained below 30 °C during the addition. After the addition, the reaction mixture was stirred at room temperature for 1.5 hour, heated to 68 °C for 1 hour, cooled, and poured into water. The precipitate was washed with water and dried in a vacuum oven to give 11.0 g of 5-chlorosulfonyl-2-oxindole (50% yield) which was used without further purification.

5-Chlorosulfonyl-2-oxindole (2.1 g) was added to 10 mL of ammonium hydroxide in 10 mL of ethanol and stirred at room temperature overnight. The mixture was concentrated and the solid collected by vacuum filtration to give 0.4 g (20% yield) of 5-aminosulfonyl-2-oxindole as an off-white solid.

¹H NMR (360 MHz, DMSO-d₆) δ 10.67 (s, 1H, NH-1), 7.63 - 7.66 (m, 2H, H-4,6), 7.13 (s, 2H, 5-SO₂NH₂), 6.91 (d, *J* = 8 Hz, 1H, H-7), and 3.56 (s, 2H, CH₂-3). MS *m/z* 211 [M-1]⁺.

EXAMPLE 93:5-Methylaminosulfonyl-2-oxindole

A suspension of 3.38 g of 5-chlorosulfonyl-2-oxindole in 10 mL of 2 M methylamine in tetrahydrofuran was stirred at room temperature for 4 hours at which time a white solid was present. The precipitate was collected by vacuum filtration, washed twice with 5 mL of water each time and dried under vacuum at 40 °C overnight to give 3.0 g (88 % yield) of 5-methylaminosulfonyl-2-oxindole.

¹H NMR (300 MHz, DMSO-d₆) δ 10.87 (s, br, 1H, NH-1), 7.86 (s, br, 1H, 5-SO₂NHCH₃), 7.61 (d, *J* = 8 Hz 1H, H-6), 7.32 (d, *J* = 5 Hz, 1H, H-4), 6.97 (d, *J* = 8 Hz, 1H, H-7), 2.53 (s, 2H, CH₂-3), and 2.36 (s, 3H, 5-SO₂NHCH₃). MS *m/z* 226.

EXAMPLE 94:

5-Dimethylaminosulfonyl-2-oxindole

A suspension of 2.3 g of 5-chlorosulfonyl-2-oxindole in 10 mL of 2 M dimethylamine in methanol was stirred at room temperature for 4 hours at which time a white solid was present. The precipitate was collected by vacuum filtration, washed with 5 mL of 1 N sodium hydroxide and 5 mL of 1 N hydrochloric acid and dried under vacuum at 40 °C overnight to give 1.9 g (79 % yield) of 5-dimethylaminosulfonyl-2-oxindole.

¹H NMR (300 MHz, DMSO-d₆) δ 10.87 (s, br, 1H, NH), 7.73 (d, *J* = 1 Hz, 1H, H-4), 7.58 (dd, *J* = 1 and 8 Hz, 1H, H-6), 7.02 (d, *J* = 8 Hz, 1H, H-7), 2.59 (s, 3H, CH₃), 2.54 (s, 2H, H-3), 2.36 (s, 3H, CH₃).

EXAMPLE 95:

5-Isopropylaminosulfonyl-2-oxindole

A suspension of 3 g of 5-chlorosulfonyl-2-oxindole, 1.15 g of isopropylamine and 1.2 mL of pyridine in 50 mL of dichloromethane was stirred at room temperature for 4 hours at which time a white solid was present. The precipitate was collected by vacuum filtration. The solids were slurry-washed with hot ethanol, cooled, collected by vacuum filtration and dried under vacuum at 40 °C overnight to give 1.5 g (45% yield) of 5-isopropylaminosulfonyl-2-oxindole.

¹H NMR (360 MHz, DMSO-d₆) δ 10.69 (s, br, 1H, NH), 7.63 (dd, *J* = 2 and 8 Hz, 1H, H-6), 7.59 (d, *J* = 2 Hz, 1H, H-4), 7.32 (d, *J* = 7 Hz, 1H, NH-SO₂-), 6.93 (d, *J* = 8 Hz, 1H, H-7), 3.57 (s, 2H, H-3), 3.14-3.23 (m, 1H, CH-(CH₃)₂), 0.94 (d, *J* = 7 Hz, 6H, 2xCH₃).

EXAMPLE 96:

5-Phenyl-1,3-dihydro-indol-2-one

5-Bromo-2-oxindole (5 g, 23.5 mmol) was dissolved in 110 mL of toluene and 110 mL of ethanol with stirring and a little heating.

Tetrakis(triphenylphosphine)-palladium(0) (1.9 g, 1.6 mmol) was added followed by a 2 M aq. solution of sodium carbonate (40 mL, 80 mmol). To this mixture benzene

boronic acid (3.7 g, 30.6 mmol) was added and the mixture was heated in a 100 °C oil bath. After 12 h, the reaction was diluted with ethyl acetate (500 mL), washed with saturated aq. sodium bicarbonate (200 mL), water (200 mL), 1 N HCl (200 mL) and brine (200 mL). The organic layer was dried with magnesium sulfate and concentrated to afford a brown solid. Trituration with dichloromethane and filtering afforded 3.8 g (77%) of a tan solid.

¹H NMR (360 MHz, DMSO-d₆) 10.4 (br s, 1H, NH), 7.57 (dd, *J* = 1.8 and 7.2 Hz, 1H, Ar-H), 7.5 to 7.35 (m, 5H, Ar-H), 7.29 (m, 1H, Ar-H), 6.89 (d, *J* = 8.2 Hz, 1H, Ar-H), 3.51 (s, 2H, CH₂CO). MS *m/z* (relative intensity %, ion) found 209 (100, M⁺); calc. 209.2.

EXAMPLE 97:

6-Pyridin-3-yl-1,3-dihydro-indol-2-one

To a solution of 6-bromo-2-oxindole (4 g, 26.3 mmol) dissolved in 60 mL of toluene and 60 mL of ethanol with stirring and a little heating tetrakis(triphenylphosphine)palladium(0) (2.3 g, 1.9 mmol) was added followed by a 2 M aqueous solution of sodium carbonate (50 mL, 100 mmol) and pyridine-3-boronic acid propane diol (5 g, 30.7 mmol). The mixture was heated in a 100 °C oil bath for 12 hours. The cooled reaction was diluted with ethyl acetate (500 mL), washed with saturated aqueous sodium bicarbonate (200 mL), water (200 mL) and brine (200 mL). The organic layer was dried with magnesium sulfate and concentrated to afford a brown solid. The residue was triturated with methylene chloride/diethyl ether to give 2.32 g (42%) of 6-pyridin-3-yl-1,3-dihydro-indol-2-one as a brown solid.

¹H NMR (360 MHz, DMSO-d₆) δ 10.51 (s, 1H, NH), 8.81 (d, *J* = 2.5 Hz, 1H, Ar-H), 8.55 (dd, *J* = 1.8 and 5.7 Hz, 1H, Ar-H), 8 (m, 1H, Ar-H), 7.45 (dd, *J* = 5.7 and 9.3 Hz, 1H, Ar-H), 7.3 (m, 2H, Ar-H), 7.05 (s, 1H, Ar-H), 3.51 (s, 2H, CH₂CO). MS *m/z* 210 [M]⁺.

EXAMPLE 98:

6-Phenyl-2-oxindole

5 Tetrakis(triphenylphosphine)palladium (0.8 g) was added to a mixture of benzeneboronic acid (3.1 g), 5-bromo-2-fluoronitrobenzene (5 g) and 22 mL of 2 M sodium carbonate solution in toluene (50 mL) and ethanol (50 mL). The mixture was heated to reflux for 2 hours, concentrated, and the residue was extracted twice with ethyl acetate. The ethyl acetate layer was washed with water, brine, dried, and concentrated to give a yellow oil. The oil was purified by column chromatography on silica gel in 5% ethyl acetate in hexane to give 4.75 g (96% yield) of 4-fluoro-3-nitrobiphenyl as a yellow oil.

10 Dimethyl malonate (10 mL) in 25 mL of dimethylsulfoxide was added dropwise to 3.5 g of sodium hydride suspended in 25 mL of dimethylsulfoxide and the mixture heated at 100 °C for 10 minutes. The mixture was cooled to room temperature and 4.7 g of 4-fluoro-3-nitrobiphenyl in 25 mL of dimethylsulfoxide was added. The mixture was heated at 100 °C for 2 hours, cooled and quenched with 300 mL of saturated ammonium chloride solution. The mixture was extracted three times with ethyl acetate and the combined organic layers were washed with water and brine and evaporated to give crude dimethyl-3-nitrobiphenyl-4-malonate as a yellow oil.

20 Crude dimethyl-3-nitrobiphenyl-4-malonate was heated to reflux in 30 mL of 6 N hydrochloric acid for 24 hours. The precipitate was collected by filtration, washed with water and dried to give 4.5 g (80% based on 4-fluoro-3-nitrobiphenyl) of 3-nitrobiphenyl-4-acetic acid as a cream colored solid.

25 Iron chips (2.6 g) were added all at once to 4.5 g of 3-nitrobiphenyl-4-acetic acid in 40 mL of acetic acid. The mixture was heated to reflux for 2 hours, concentrated to dryness and taken up in ethyl acetate. The solids were removed by filtration and the filtrate was washed twice with 1 N hydrochloric acid and brine and dried over anhydrous sodium sulfate. The filtrate was concentrated to give 3.4 g (93% yield) of 6-phenyl-2-oxindole as a light brown solid.

¹H NMR (360 MHz, DMSO-d₆) δ 10.4 (s, br, 1H, NH-1), 7.57-7.6 (m, 2H), 7.42-7.46 (m, 2H), 7.32-7.37 (m, 1H), 7.27 (d, *J* = 8 Hz, 1H, H-4), 7.19 (dd, *J* = 2, 8 Hz, 1H, H-5), 7.01 (d, *J* = 2 Hz, 1H, H-7), 3.49 (s, 2H, CH₂). MS *m/z* 210 [M+1]⁺.

5 EXAMPLE 99:

6-(2-Methoxyphenyl)-2-oxindole

Tetrakis(triphenylphosphine)palladium (1 g) was added to a mixture of 2-methoxyphenylboronic acid (5 g), 5-bromo-2-fluoronitrobenzene (6.6 g) and 30 mL of 2 M sodium carbonate solution in toluene (50 mL) and ethanol (50 mL). The mixture was heated to reflux for 2 hours, concentrated, and the residue was extracted twice with ethyl acetate. The ethyl acetate layer was washed with water, brine, dried, and concentrated to give a dark green oil which solidified on standing, crude 4-fluoro-2'-methoxy-3-nitrobiphenyl.

Dimethyl malonate (14 mL) was added dropwise to 2.9 g of sodium hydride suspended in 50 mL of dimethylsulfoxide. The mixture was heated at 100 °C for 15 minutes and cooled to room temperature. Crude 4-fluoro-2'-methoxy-3-nitrobiphenyl in 60 mL of dimethylsulfoxide was added and the mixture was heated at 100 °C for 2 hours. The reaction mixture was cooled and quenched with 300 mL of saturated ammonium chloride solution and extracted twice with ethyl acetate. The extracts were combined, washed with saturated ammonium chloride, water, and brine, dried over anhydrous sodium sulfate and concentrated to give crude dimethyl 2'-methoxy-3-nitrobiphenyl-4-malonate as a yellow oil.

Crude 2'-methoxy-3-nitrobiphenyl-4-malonate was heated at 100 °C in 50 mL of 6 N hydrochloric acid for 24 hours and cooled. The precipitate was collected by filtration, washed with water and hexane, and dried to give 9.8 of 2'-methoxy-2-nitrobiphenyl-4-acetic acid as a light tan solid.

Iron chips (5 g) were added in one portion to 9.8 g of 2'-methoxy-3-nitrobiphenyl-4-acetic acid in 50 mL of glacial acetic acid and heated to 100 °C for 3 hours. The reaction mixture was concentrated to dryness, sonicated in ethyl acetate and filtered to remove the insolubles. The filtrate was washed twice with 1 N hydrochloric acid, water, brine, dried over anhydrous sodium sulfate and

concentrated. The residue was purified by column chromatography on silica gel in ethyl acetate:hexane 1:2 to give 5.4 g (69 % yield based on 5-bromo-2-fluoronitrobenzene) of 6-(2-methoxyphenyl)-2-oxindole as a rose colored solid.

¹H NMR (360 MHz, DMSO-d₆) δ 10.32 (s, br, 1H, NH), 7.29- 7.34 (m, 1H), 7.19-7.25 (m, 2H), 7.08 (d, *J* = 8 Hz, 1H, H-4), 6.97-7.02 (m, 2H), 6.91 (d, *J* = 1 Hz, 1H, H-7), 3.8 (s, 3H, OCH₃), 3.47 (s, 2H, CH₂). MS *m/z* 239.8 [M+1]⁺.

EXAMPLE 100:

6-(3-Methoxyphenyl)-2-oxindole

10 Tetrakis(triphenylphosphine)palladium (0.7 g) was added to a mixture of 3.8 g of 3-methoxyphenylboronic acid, 5 g of 5-bromo-2-fluoronitrobenzene and 11 mL of 2 M sodium carbonate solution in 100 mL of toluene. The mixture was heated to reflux for 2 hours, diluted with water and extracted with ethyl acetate. The ethyl acetate was washed with saturated sodium bicarbonate, brine, dried, and
15 concentrated to give an oily solid. The solid was purified by column chromatography on silica gel in ethyl acetate:hexane 1:6 to give 4.3 g (77% yield) of 4-fluoro-3'-methoxy-3-nitrobiphenyl.

Dimethyl malonate (9.7 mL) was added dropwise to 2.0 g of sodium hydride suspended in 50 mL of dimethylsulfoxide. The mixture was heated to 100 °C for 35
20 minutes and cooled to room temperature. 4-Fluoro-2'-methoxy-3-nitrobiphenyl (4.2 g) in 50 mL of dimethylsulfoxide was added and the mixture was heated at 100 °C for 1 hour. The reaction mixture was cooled and quenched with 300 mL of saturated ammonium chloride solution and extracted twice with ethyl acetate. The extracts were combined, washed with brine, dried over anhydrous sodium sulfate and
25 concentrated to give crude dimethyl 3'-methoxy-3-nitrobiphenyl-4-malonate as a pale yellow solid.

Crude 3'-methoxy-3-nitro-biphenyl-4-malonate was heated at 110 °C in 45 mL of 6 N hydrochloric acid for 4 days and cooled. The precipitate was collected by filtration, washed with water and hexane, and dried to give 5.3 g of 3'-methoxy-2-
30 nitrobiphenyl-4-acetic acid as a light tan solid.

3'-Methoxy-3-nitrobiphenyl-4-acetic acid (5.2 g) was dissolved in methanol and hydrogenated over 0.8 g of 10% palladium on carbon for 3 hours at room temperature. The catalyst was removed by filtration, washed with methanol and the filtrates combined and concentrated to give a brown solid. The solid was purified by column chromatography on silica gel in ethyl acetate:hexane:acetic acid 33:66:1 to give 3.0 g (75% yield based on 4-fluoro-3'-methoxy-3-nitrobiphenyl) of 6-(3-methoxyphenyl)-2-oxindole as a pink solid.

¹H NMR (360 MHz, DMSO-d₆) δ 10.39 (s, br, 1H, NH), 7.35 (t, *J* = 8 Hz, 1H), 7.26 (d, *J* = 8 Hz, 1H), 7.19 (dd, *J* = 1, 8 Hz, 1H), 7.13-7.16 (m, 1H), 7.09-7.1 (m, 1H), 7.01 (d, *J* = 1 Hz, 1H), 6.90-6.93 (m, 1H), 3.8 (s, 3H, OCH₃), 3.49 (s, 2H, CH₂). MS *m/z* 240.0 [M+1]⁺.

EXAMPLE 101:

6-(4-Methoxyphenyl)-2-oxindole

Tetrakis(triphenylphosphine)palladium (1 g) was added to a mixture of 5 g of 4-methoxyphenylboronic acid, 6.6 g of 5-bromo-2-fluoronitrobenzene and 30 mL of 2 M sodium carbonate solution in 50 mL of toluene and 50 mL of ethanol. The mixture was heated to reflux for 2 hours, concentrated, and the residue extracted twice with ethyl acetate. The ethyl acetate layer was washed with water, brine, dried, and concentrated to give a brown oily solid. The solid was purified by column chromatography on silica gel in 5 % ethyl acetate in hexane to give crude 4-fluoro-4'-methoxy-3-nitrobiphenyl as a pale yellow solid.

Dimethyl malonate (10 mL) was added dropwise to 2.0 g of sodium hydride suspended in 60 mL of dimethylsulfoxide. The mixture was heated to 100 °C for 10 minutes and cooled to room temperature. Crude 4-fluoro-2'-methoxy-3-nitrobiphenyl (5.2 g) in 50 mL of dimethylsulfoxide was added and the mixture was heated at 100 °C for 2 hours. The reaction mixture was cooled and quenched with 300 mL of saturated ammonium chloride solution and extracted three times with ethyl acetate. The extracts were combined, washed with saturated ammonium chloride, water and brine, dried over anhydrous sodium sulfate and concentrated to give crude dimethyl 4'-methoxy-3-nitrobiphenyl-4-malonate as a yellow oil.

Crude 4'-methoxy-3-nitro-biphenyl-4-malonate was heated at 100 °C in 60 mL of 6 N hydrochloric acid for 15 hours and cooled. The precipitate was collected by filtration, washed with water and hexane, and dried to give 7.2 g of crude 4'-methoxy-3-nitrobiphenyl-4-acetic acid as a light tan solid.

5 Iron chips (3.6 g) were added in one portion to 7.2 g of 4'-methoxy-3-nitrobiphenyl-4-acetic acid in 50 mL of glacial acetic acid and heated at 100 °C overnight. The reaction mixture was concentrated to dryness, sonicated in ethyl acetate and filtered to remove the insolubles. The filtrate was washed twice with 1 N hydrochloric acid, brine, dried over anhydrous sodium sulfate and concentrated to
10 give 2.7 g (54% yield based on 5-bromo-2-fluoronitrobenzene) of 6-(4-methoxyphenyl)-2-oxindole as a rose colored solid.

¹H NMR (360 MHz, DMSO-d₆) δ 10.38 (s, br, 1H, NH-1), 7.52 (d, *J* = 9 Hz, 2H), 7.23 (d, *J* = 7 Hz, 1H, H-4), 7.14 (dd, *J* = 1, 7 Hz, 1H, H-5), 7.0 (d, *J* = 9 Hz, 2H), 6.96 (d, *J* = 1 Hz, 1H, H-7), 3.78 (s, 3H, OCH₃), 3.47 (s, 2H, CH₂). MS *m/z*
15 214.0 [M+1]⁺.

EXAMPLE 102:

2-Oxo-2,3-dihydro-1H-indole-4-carboxylic acid (3-chloro-4-methoxy-phenyl)-amide

To a solution of 2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid (200 mg, 1.13 mmol) and 3-chloro-4-methoxy-phenylamine (178 mg, 1.13 mmol) in dimethylformamide (15 mL) at room temperature benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (BOP reagent 997 mg, 2.26 mmol) was added followed by 4-dimethylaminopyridine (206 mg, 1.69 mmol). The mixture was stirred at room temperature for 72 hours. The reaction was then
25 diluted with ethyl acetate (300 mL), washed with saturated sodium bicarbonate solution (100 mL), water, 2 N hydrochloric acid (100 mL), water (3x200 mL) and brine, dried (over magnesium sulfate) and concentrated. The residue was triturated with ethyl acetate to give 2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid (3-chloro-4-methoxy-phenyl)-amide as a tarnish-pink solid.

30 ¹H NMR (360 MHz, DMSO-d₆) δ 10.50 (s, br, 1H, NH), 10.12 (s, br, 1H, NH), 7.9 (s, *J* = 2.5 Hz, 1H), 7.62 (dd, *J* = 2.5 & 9 Hz, 1H), 7.38 (d, *J* = 7.6 Hz, 1H),

7.32 (t, $J = 7.6$ Hz, 1H), 7.13 (d, $J = 9$ Hz, 1H), 6.98 (d, $J = 7.6$ Hz, 1H), 3.83 (s, 3H, OCH₃), 3.69 (s, 2H, CH₂). MS-EI m/z 316 [M]⁺.

EXAMPLE 103:

5 2-Oxo-2,3-dihydro-1H-indole-4-carboxylic acid ethylamide

Ethylamine (2.1 mL of 2.0 M solution in tetrahydrofuran, 4.2 mmol) was added dropwise to a suspension of 4-carboxy-2-oxindole (380 mg, 2.1 mmol) in dimethylformamide (8 mL), followed by 4-dimethylaminopyridine (385 mg, 3.15 mmol) and PyBOP (2.185 g, 4.2 mmol). The mixture was stirred at room
10 temperature for 6 hrs. The reaction was concentrated. The residue was dissolved in dichloromethane, washed with saturated sodium bicarbonate and brine, dried, concentrated and recrystallized to give 280 mg (65%) of 2-oxo-2,3-dihydro-1H-indole-4-carboxylic acid ethylamide.

¹H NMR (300 MHz, DMSO-d₆) δ 10.44 (br s, 1H, NH), 8.24 (m, 1H, CONHCH₂CH₃), 7.24 (m, 2H), 6.90 (dd, $J = 2.1$ & 6.4 Hz, 1H), 3.63 (s, 2H, CH₂),
15 3.24 (m, 2H, NCH₂CH₃), 1.10 (t, $J = 7.1$ Hz, 3H, NCH₂CH₃). MS-EI 204 [M]⁺.

EXAMPLE 104:

General Synthetic Method for the Combinatorial Libraries

20 The preparation of the compounds of the invention that were prepared using the combinatorial library method followed one of the following general schemes.


Solution Phase:

0.5M solutions in DMSO of oxindoles and aldehydes were prepared.
25 Piperidine (20 μ L) was added to each solution to aid in solvation. 220 μ L each of the appropriately substituted oxindole and the appropriately substituted aldehyde were added to each well, followed by piperidine (30 μ L). The mixtures were heated at 70°C for 15 hours. After cooling down, DMSO was added to each well to make the total volume to 1 μ L. The solutions were then transferred to 96 mother plate and
30 used for biological testing.

0.5M Solutions in DMSO (2.5 μ L) of oxindoles and aldehydes were prepared. Piperidine (20 μ L) was added to each solution to aid in solvation. 220 μ L each of the appropriately substituted oxindole and the appropriately substituted aldehyde were added to each well, followed by piperidine (30 μ L). The mixtures were heated at 70°C for 15 hours. After cooling down, DMSO was added to each well to make the total volume to 1 mL. Piperidine was removed under high vacuum. The solutions were then transferred to 96 mother-well plate and used for biological testing.


10 Solid Phase:

1. Attachment of resin to oxindoles



15 Starting oxindole (1.584 g, 1.2 eq) was dissolved in 50 mL Dichloromethane and 5.2 mL (4 eq) DIPEA under nitrogen. DMF (5 mL) was added to aid in solvation. Resin (5 g, 7.45 mmol, 1 eq) was added and stirred gently overnight. Filtered resin and washed with 300 mL of 17:2:1 DCM/MeOH/DIPEA, 200 mL of DMF, and 200 mL of DCM.

20 The solvent was removed under high vacuum. 6.12g of resin was collected.




Starting oxindole (1.7 g, 2 eq) was dissolved in 68 mL DMF under nitrogen. Diisopropylcarbodiimide (DIC) (1.5 mL, 2eq) was added followed by 234 mg (0.4

eq) of DMAP. Then resin (6.85g, 4.798mmol, 1eq) was added and stirred gently overnight. Resin was filtered and washed with DMF, MeOH and DCM.

The solvent was remove under high vacuum. 7.45g of resin was collected.

2. Condensation of the resin with aldehydes



Resin (125 mg, 1 eq) was treated with aldehyde (1 mL of 0.2 M solution in DMSO, 2 eq) and 1 mL of 50% piperidine in DMSO at 60 °C for 14 h. Resin was filtered and washed with DMSO, DMF, and MeOH. It was then treated to 1.5 mL of 10% TFA/DCM for 1 h. Resin was filtered again and washed with DMSO.

Volatiles were removed from filtrate in vacuum.

ASSAY PROCEDURES

The following *in vitro* assays may be used to determine the level of activity and effect of the different compounds of the present invention on one or more of the PKs. Similar assays can be designed along the same lines for any PK using techniques well known in the art.

Three general types of assays are useful for evaluating compounds: cellular/catalytic, cellular/biological and *in vivo*. The object of the cellular/catalytic assays is to determine the effect of a compound on the ability of a TK to phosphorylate tyrosines on a known substrate in a cell. The object of the cellular/biological assays is to determine the effect of a compound on the biological

response stimulated by a TK in a cell. The object of the *in vivo* assays is to determine the effect of a compound in an animal model of a particular disorder such as cancer.

5 The cellular/catalytic assays described herein are performed in an ELISA format. The general procedure is as follows: a compound is introduced to cells expressing the test kinase, either naturally or recombinantly, for some period of time after which, if the test kinase is a receptor, a ligand known to activate the receptor is added. The cells are lysed and the lysate is transferred to the wells of an ELISA plate previously coated with a specific antibody recognizing the substrate of the enzymatic phosphorylation reaction. Non-substrate components of the cell lysate are washed away and the amount of phosphorylation on the substrate is detected with an antibody specifically recognizing phosphotyrosine compared with control cells that were not contacted with a test compound.

15 The cellular/biologic assays described herein measure the amount of DNA made in response to activation of a test kinase, which is a general measure of a proliferative response. The general procedure for this assay is as follows: a compound is introduced to cells expressing the test kinase, either naturally or recombinantly, for some period of time after which, if the test kinase is a receptor, a ligand known to activate the receptor is added. After incubation at least overnight, a DNA labeling reagent such as Bromodeoxy-uridine (BrdU) or 3H-thymidine is added. The amount of labeled DNA is detected with either an anti-BrdU antibody or by measuring radioactivity and is compared to control cells not contacted with a test compound.

25 Cellular/Catalytic Assays

Enzyme linked immunosorbent assays (ELISA) may be used to detect and measure the presence of PK activity. The ELISA may be conducted according to known protocols which are described in, for example, Voller, et al., 1980, "Enzyme-Linked Immunosorbent Assay," In: Manual of Clinical Immunology, 2d ed., edited by Rose and Friedman, pp 359-371 Am. Soc. Of Microbiology, Washington, D.C.

The disclosed protocol may be adapted for determining activity with respect to a specific PK. For example, the preferred protocols for conducting the ELISA experiments for specific PKs is provided below. Adaptation of these protocols for determining a compound's activity for other members of the RTK family, as well as for CTKs and STKs, is well within the scope of knowledge of those skilled in the art.

EXAMPLE 105: FLK-1

An ELISA assay was conducted to measure the kinase activity of the FLK-1 receptor and more specifically, the inhibition or activation of TK activity on the FLK-1 receptor. Specifically, the following assay was conducted to measure kinase activity of the FLK-1 receptor in cells genetically engineered to express Flk-1.

Materials and Methods.

Materials.

- The following reagents and supplies were used:
- a. Corning 96-well ELISA plates (Corning Catalog No. 25805-96);
 - b. Cappel goat anti-rabbit IgG (catalog no. 55641);
 - c. PBS (Gibco Catalog No. 450-1300EB);
 - d. TBSW Buffer (50 mM Tris (pH 7.2), 150 mM NaCl and 0.1% Tween-20);
 - e. Ethanolamine stock (10% ethanolamine (pH 7.0), stored at 4 °C);
 - f. HNTG buffer (20 mM HEPES buffer (pH 7.5), 150 mM NaCl, 0.2% Triton X-100, and 10% glycerol);
 - g. EDTA (0.5 M (pH 7.0) as a 100X stock);
 - h. Sodium orthovanadate (0.5 M as a 100X stock);
 - i. Sodium pyrophosphate (0.2 M as a 100X stock);
 - j. NUNC 96 well V bottom polypropylene plates (Applied Scientific Catalog No. AS-72092);
 - k. NIH3T3 C7#3 Cells (FLK-1 expressing cells);
 - l. DMEM with 1X high glucose L-Glutamine (catalog No. 11965-050);
 - m. FBS, Gibco (catalog no. 16000-028);

- n. L-glutamine, Gibco (catalog no. 25030-016);
- o. VEGF, PeproTech, Inc. (catalog no. 100-20)(kept as 1 µg/100 µL stock in Milli-Q dH₂O and stored at -20 °C;
- p. Affinity purified anti-FLK-1 antiserum;
- 5 q. UB40 monoclonal antibody specific for phosphotyrosine (*see*, Fendley, *et al.*, 1990, *Cancer Research* 50:1550-1558);
- r. EIA grade Goat anti-mouse IgG-POD (BioRad catalog no. 172-1011);
- s. 2,2-azino-bis(3-ethylbenz-thiazoline-6-sulfonic acid (ABTS) solution
- 10 (100 mM citric acid (anhydrous), 250 mM Na₂HPO₄ (pH 4.0), 0.5 mg/mL ABTS (Sigma catalog no. A-1888)), solution should be stored in dark at 4 °C until ready for use;
- t. H₂O₂ (30% solution) (Fisher catalog no. H325);
- u. ABTS/H₂O₂ (15 mL ABTS solution, 2 µL H₂O₂) prepared 5 minutes
- 15 before use and left at room temperature;
- v. 0.2 M HCl stock in H₂O;
- w. dimethylsulfoxide (100%) (Sigma Catalog No. D-8418); and
- y. Trypsin-EDTA (Gibco BRL Catalog No. 25200-049).

20 Protocol

The following protocol was used for conducting the assay:

1. Coat Corning 96-well ELISA plates with 1.0 µg per well Cappel Anti-rabbit IgG antibody in 0.1 M Na₂CO₃ pH 9.6. Bring final volume to 150 µL per well. Coat plates overnight at 4 °C. Plates can be kept up to two weeks when stored
- 25 at 4 °C.
2. Grow cells in Growth media (DMEM, supplemented with 2.0mM L-Glutamine, 10% FBS) in suitable culture dishes until confluent at 37 °C, 5% CO₂.
3. Harvest cells by trypsinization and seed in Corning 25850 polystyrene 96-well round bottom cell plates, 25,000 cells/well in 200 µL of growth
- 30 media.
4. Grow cells at least one day at 37 °C, 5% CO₂.

5. Wash cells with D-PBS 1X.
6. Add 200 μ L/well of starvation media (DMEM, 2.0 mM l-Glutamine, 0.1% FBS). Incubate overnight at 37 °C, 5% CO₂.
7. Dilute Compounds 1:20 in polypropylene 96 well plates using starvation media. Dilute dimethylsulfoxide 1:20 for use in control wells.
8. Remove starvation media from 96 well cell culture plates and add 162 μ L of fresh starvation media to each well.
9. Add 18 μ L of 1:20 diluted Compound dilution (from step 7) to each well plus the 1:20 dimethylsulfoxide dilution to the control wells (+/- VEGF), for a final dilution of 1:200 after cell stimulation. Final dimethylsulfoxide is 0.5%. Incubate the plate at 37 °C, 5% CO₂ for two hours.
10. Remove unbound antibody from ELISA plates by inverting plate to remove liquid. Wash 3 times with TBSW + 0.5% ethanolamine, pH 7.0. Pat the plate on a paper towel to remove excess liquid and bubbles.
11. Block plates with TBSW + 0.5% Ethanolamine, pH 7.0, 150 μ L per well. Incubate plate thirty minutes while shaking on a microtiter plate shaker.
12. Wash plate 3 times as described in step 10.
13. Add 0.5 μ g/well affinity purified anti-FLU-1 polyclonal rabbit antiserum. Bring final volume to 150 μ L/well with TBSW + 0.5% ethanolamine pH 7.0. Incubate plate for thirty minutes while shaking.
14. Add 180 μ L starvation medium to the cells and stimulate cells with 20 μ L/well 10.0 mM sodium ortho vanadate and 500 ng/mL VEGF (resulting in a final concentration of 1.0 mM sodium ortho vanadate and 50 ng/mL VEGF per well) for eight minutes at 37 °C, 5% CO₂. Negative control wells receive only starvation medium.
15. After eight minutes, media should be removed from the cells and washed one time with 200 μ L/well PBS.
16. Lyse cells in 150 μ L/well HNTG while shaking at room temperature for five minutes. HNTG formulation includes sodium ortho vanadate, sodium pyrophosphate and EDTA.
17. Wash ELISA plate three times as described in step 10.

18. Transfer cell lysates from the cell plate to ELISA plate and incubate while shaking for two hours. To transfer cell lysate pipette up and down while scrapping the wells.

19. Wash plate three times as described in step 10.

5 20. Incubate ELISA plate with 0.02 µg/well UB40 in TBSW + 05% ethanolamine. Bring final volume to 150 µL/well. Incubate while shaking for 30 minutes.

21. Wash plate three times as described in step 10.

10 22. Incubate ELISA plate with 1:10,000 diluted EIA grade goat anti-mouse IgG conjugated horseradish peroxidase in TBSW + 0.5% ethanolamine, pH 7.0. Bring final volume to 150 µL/well. Incubate while shaking for thirty minutes.

23. Wash plate as described in step 10.

24. Add 100 µL of ABTS/H₂O₂ solution to well. Incubate ten minutes while shaking.

15 25. Add 100 µL of 0.2 M HCl for 0.1 M HCl final to stop the color development reaction. Shake 1 minute at room temperature. Remove bubbles with slow stream of air and read the ELISA plate in an ELISA plate reader at 410 nm.

EXAMPLE 106: GST-FLK-1 BIOASSAY

20 This assay analyzes the tyrosine kinase activity of GST-Flk1 on poly glu tyr peptides.

Materials and Reagents:

1. Corning 96-well Elisa plates (Corning Catalog No. 25805-96).
- 25 2. poly glu tyr 4:1, lyophilizate (Sigma Catalog # P0275). Prepare 1 mg/mL poly glu tyr in sterile PBS and store in 1ml aliquots at -20 °C.
3. Preparation of poly glu tyr (pEY) coated assay plates: Coat 2 µg/well of poly glu tyr (pEY) in 100 µL PBS at room temperature for 2 hours or at +4 °C overnight. Cover plates well to prevent evaporation.
- 30 4. PBS Buffer: To make 1 liter of a 1x working solution, mix .02 g KH₂PO₄, 1.15 g Na₂HPO₄, 0.2 g KCl and 8 g NaCl in approx. 900 mL dH₂O.

When all reagents have dissolved, adjust the pH to 7.2 with HCl. Bring total volume to 1 L with dH₂O.

5. PBS-Tw Buffer: To 1 L of PBS Buffer, add 1/0 mL Tween-20. Stir until dissolved.
- 5 6. TBB - Blocking Buffer: To make one liter of a 1x working solution, mix 1.21 g TRIS, 8.77 g NaCl, 1 mL TWEEN-20 in approximately 900 mL dH₂O. Adjust pH to 7.2 with HCl. Add 10 g BSA, stir to dissolve. Bring total volume to 1 L with dH₂O. Filter to remove particulate matter.
7. 1% BSA in PBS: To make a 1x working solution, add 10 g BSA to 10 approx. 990 mL PBS buffer, stir to dissolve. Adjust total volume to 1 L with PBS buffer, filter to remove particulate matter.
8. 50 mM Hepes pH 7.5.
9. GST-Flk1cd purified from sf9 recombinant baculovirus transformation.
- 15 10. 4% DMSO in dH₂O.
11. 10 mM ATP in dH₂O.
12. 40 mM MnCl₂
13. Kinase Dilution Buffer (KDB): Mix 10 mL Hepes (pH 7.5), 1 mL of 5M NaCl, 40 µL of 100 mM NaVO₄ and 0.4 mL of 5% BSA in dH₂O with 88.56 mL 20 of dH₂O.
14. NUNC 96-well V bottom polypropylene plates Applied Scientific Catalog # AS-72092
15. EDTA: Mix 14.12 g ethylenediaminetetraacetic acid (EDTA) to approx. 70 mL dH₂O. Add 10 N NaOH until EDTA dissolves. Adjust pH to 8.0. 25 Adjust total volume to 100 mL with dH₂O.
16. 1° Antibody Dilution Buffer: Mix 10 mL of 5% BSA in PBS buffer with 89.5 mL TBSTw.
17. Anti-phosphotyrosine monoclonal conjugated to horseradish peroxidase (PY99 HRP, Santa Cruz Biotech).
- 30 18. 2,2'-Azino-bis(3-ethylbenzthiazoline-6-sulfonic acid (ABTS, Moss, Cat. No. ABST).

19. 10% SDS.

PROCEDURE:

1. Coat Corning 96 well ELISA plates with 2 µg of polyEY peptide in
5 sterile PBS as described in step 3 of Materials and Reagents.
2. Remove unbound liquid from wells by inverting plate. Wash once
with TBSTw. Pat the plate on a paper towel to remove excess liquid.
3. Add 100 µL of 1% BSA in PBS to each well. Incubate, with shaking,
for 1 hr. at room temperature.
- 10 4. Repeat step 2.
5. Soak wells with 50 mM Hepes pH7.5 (150 µL/well).
6. Dilute test compound with dH₂O/4% DMSO to 4 times the desired
final assay concentration in 96-well polypropylene plates.
7. Add 25 µL diluted test compound to ELISA plate. To control wells
15 (wells which do not receive any test compound), add 25 µL of dH₂O/4% DMSO.
8. Add 25 µL of 40 mM MnCl₂ with 4x ATP (2 µM) to all wells.
9. Add 25 µL 0.5M EDTA to negative control wells.
10. Dilute GST-Flk1 0.005 µg (5 ng)/well in KDB. For 50 ml KDB add
100 µL of 0.050 mg/mL GST-Flk1 enzyme.
- 20 11. Add 50 µL of diluted enzyme to each well.
12. Incubate, with shaking, for 15 minutes at room temperature.
13. Stop reaction by adding 50 µL of 250 mM EDTA (pH 8.0).
14. Wash 3X with TBSTw and pat plate on paper towel to remove excess
liquid.
- 25 15. Add 100 µL per well anti-phosphotyrosine HRP conjugate, 1:5,000
dilution in antibody dilution buffer. Incubate, with shaking for 90 min. at room
temperature.
16. Wash as described above in step 14.
17. Add 100 µL of room temperature ABTS solution to each well.
- 30 18. Incubate, with shaking, for 10 to 15 minutes. Remove any bubbles.
19. Stop reaction by adding 20 µL of 10% SDS.

20. Read assay on Dynatech MR7000 ELISA reader: test filter at 410 nM; reference filter at 630 nM.

EXAMPLE 107: PYK2 BIOASSAY

5 This assay is used to measure the in vitro kinase activity of HA epitope tagged full length pyk2 (FL.pyk2-HA) in an ELISA assay.

Materials and reagents:

1. Corning 96-well Elisa plates (Corning Catalog # 25805-96).
- 10 2. 12CA5 monoclonal anti-HA antibody
3. PBS (Dulbecco's Phosphate-Buffered Saline, Gibco Catalog # 450-1300EB)
4. TBST Buffer: Mix 8.766 g NaCl, 6.057 g TRIS and 1 ml of 0.1% Triton X-100 in approx. 900 mL dH₂O. Adjust pH to 7.2, bring volume to 1 L.
- 15 5. Blocking Buffer: Mix 100 g of 10% BSA, 12.1 g of 100 mM TRIS, 58.44 g of 1 M NaCl and 10 mL of 1% TWEEN-20.
6. FL.pyk2-HA from sf9 cell lysates.
7. 4% DMSO in MilliQue H₂O.
8. 10 mM ATP in dH₂O.
- 20 9. 1 M MnCl₂.
10. 1 M MgCl₂.
11. 1 M Dithiothreitol (DTT).
12. 10X Kinase buffer phosphorylation mix: Mix 5.0 mL 1 M Hepes (pH 7.5), 0.2 mL 1 M MnCl₂, 1.0 mL MgCl₂, 1.0 mL 10% Triton X-100 in 2.8 ml dH₂O.
- 25 Just prior to use, add 0.1 mL 1 M DTT.
13. NUNC 96-well V bottom polypropylene plates (Applied Scientific Catalog # AS-72092).
14. EDTA
15. Biotin conjugated anti-phosphotyrosine mab (Upstate Biotechnology
- 30 Inc., clone 4G10 cat.# 16-103, ser. # 14495).

16. Vectastain Elite ABC reagent (Avidin peroxidase conjugate, Vector Laboratories (PK-6100).

17. ABTS Solution: Mix 19.21 g citric acid and 35.49 g Na_2HPO_4 in approx. 900 mL dH_2O . Adjust pH to 4.0 with phosphoric acid. Add 5 or 10 g

5 ABST. When all dissolved, filter.

18. Hydrogen peroxide 30% solution.

19. ABTS/ H_2O_2 : Mix 15 mL ABTS solution with 3 μL 30% H_2O_2 5 min. before use.

20. 0.2 M HCl.

10

Procedure:

1. Coat Corning 96 well ELISA plates with 0.5 μg per well 12CA5 anti-HA antibody in 100 μL PBS. Store overnight at 4 °c.

2. Remove unbound HA antibody from wells by inverting plate. Pat the
15 plate on a paper towel to remove excess liquid.

3. Add 150 μL Blocking Buffer to each well. Incubate, with shaking, for 1 hr at room temperature.

4. Wash plates with TBS-T.

5. Dilute lysate in PBS (1.5 μg lysate/100 μL PBS).

20 6. Add 100 μL of diluted lysate to each well. Shake at room temperature for 1 hr.

7. Wash as in step 4.

8. Add 50 μL of 2X kinase Buffer to ELISA plate containing captured pyk2-HA.

25 9. Add 25 μL of 40 μM test compound in 4% DMSO or 4% DMSO alone (control) to plate.

10. Add 25 μL of 0.5 M EDTA to negative control wells.

11. Add 25 μL of 20 μM ATP to all wells. Incubate, with shaking, for 10 minutes.

30 12. Stop reaction by adding 25 μL 500 mM EDTA (pH 8.0) to all wells.

13. Wash as in step 4.

14. Add 100 μ L biotin conjugated anti-phosphotyrosine mab (1:5000 dilution in Blocking Buffer) to each well. Incubate, with shaking for 30 min. at room temperature.

15. Make up Vectastain ABC reagent. Allow 30 min. for complete coupling of the avidin with the biotinylated HRP. Add 1 drop (or 50 μ L) reagent A to 15 mL Blocking Buffer. Mix by inverting tube several times. Add 1 drop (or 50 μ L) reagent B and mix again. Allow ABC reagent to mix at room temperature while the biotin-4G10 anti-phosphotyrosine is incubating in the assay plate.

16. Wash as in step 4.

17. Add 100 μ L per well of prepared Vectastain peroxidase conjugate. Incubate, with shaking, for 30 min. at room temperature.

18. Wash as in step 4, then was once with PBS.

19. Add 100 μ L of ABTS/H₂O₂ solution to each well.

20. Incubate, with shaking, for 10 to 15 minutes. Remove any bubbles.

21. If necessary, stop reaction with the addition of 100 μ L of 0.2 M HCl per well.

22. Read assay on Dynatech MR7000 ELISA reader with test filter at 410 nM and reference filter at 630 nM.

EXAMPLE 108: FGFR1 BIOASSAY

This assay is used to measure the in vitro kinase activity of FGF1-R in an ELISA assay.

Materials and Reagents:

1. Costar 96-well Elisa plates (Corning Catalog # 3369).
2. Poly(Glu,Tyr) (Sigma Catalog # PO275).
3. PBS (Gibco Catalog # 450-1300EB)
4. 50 mM Hepes Buffer Solution.
5. Blocking Buffer (5% BSA/PBS).
6. Purified GST-FGFR1.

7. Kinase Dilution Buffer: Mix 500 μ L 1 M Hepes (GIBCO), 20 μ L 5% BSA/PBS, 10 μ L 100 mM sodium orthovanadate and 50 μ L 5 M NaCl.
8. 10 mM ATP
9. 1 M MnCl_2
- 5 10. ATP/ MnCl_2 phosphorylation mix: Mix 20 μ L ATP, 400 μ L MnCl_2 and 9.56 mL dH_2O .
11. NUNC 96-well V bottom polypropylene plates (Applied Scientific Catalog # AS-72092).
12. 0.5 M EDTA.
- 10 13. 0.05% TBST: Add 500 μ L TWEEN to 1 liter TBS.
14. Rabbit polyclonal anti-phosphotyrosine serum.
15. Goat anti-rabbit IgG peroxidase conjugate (Biosource, Catalog # ALI0404).
16. ABTS Solution
- 15 17. 30% Hydrogen peroxide.
18. ABTS/ H_2O_2

Procedure:

1. Coat Costar 96 well ELISA plates with 1 μ g per well Poly(Glu,Tyr)
- 20 in 100 μ L PBS. Store overnight at 4 °C.
2. Wash coated plates once with PBS.
3. Add 150 μ L of 5%BSA/PBS Blocking Buffer to each well. Incubate, with shaking, for 1 hr.room temperature.
4. Wash plate 2x with PBS, then once with 50 mM Hepes. Pat plates on
- 25 a paper towel to remove excess liquid and bubbles.
5. Add 25 μ L of 0.4 mM test compound in 4% DMSO or 4% DMSO alone (controls) to plate.
6. Dilute purified GST-FGFR1 in Kinase Dilution Buffer (5 ng kinase/50 μ L KDB/well).
- 30 7. Add 50 μ L of diluted kinase to each well.

8. Start kinase reaction by adding 25 μL /well of freshly prepared ATP/ Mn^{++} (0.4 mL 1 M MnCl_2 , 40 μL 10 mM ATP, 9.56 mL dH_2O), freshly prepared).
9. This is a fast kinase reaction and must be stopped with 25 μL of 0.5M EDTA in a manner similar to the addition of ATP.
10. Wash plate 4x with fresh TBST.
11. Make up Antibody Dilution Buffer: Per 50 mL: Mix 5 ml of 5% BSA, 250 μL of 5% milk and 50 μL of 100mM sodium vanadate, bring to final volume with 0.05% TBST.
12. Add 100 μL per well of anti-phosphotyrosine (1:10000 dilution in ADB). Incubate, with shaking for 1 hr. at room temperature.
13. Wash as in step 10.
14. Add 100 μL per well of Biosource Goat anti-rabbit IgG peroxidase conjugate (1:6000 dilution in ADB). Incubate, with shaking for 1 hr. at room temperature.
15. Wash as in step 10 and then with PBS to remove bubbles and excess TWEEN.
16. Add 100 μL of ABTS/ H_2O_2 solution to each well.
17. Incubate, with shaking, for 10 to 20 minutes. Remove any bubbles.
18. Read assay on Dynatech MR7000 ELISA reader: test filter at 410 nM, reference filter at 630 nM.

EXAMPLE 109: CELLULAR HER-2 KINASE ASSAY

This assay is used to measure HER-2 kinase activity in whole cells in an ELISA format.

Materials and Reagents:

1. DMEM (GIBCO Catalog #11965-092).
2. Fetal Bovine Serum (FBS, GIBCO Catalog #16000-044), heat inactivated in a water bath for 30 min. at 56 $^{\circ}\text{C}$.
3. Trypsin (GIBCO Catalog #25200-056).

4. L-Glutamine (GIBCO Catalog #25030-081)
5. HEPES (GIBCO Catalog #15630-080).
6. Growth Media: Mix 500 mL DMEM, 55 mL heat inactivated FBS, 10 mL HEPES and 5.5 ml L-Glutamine.
- 5 7. Starve Media: Mix 500 mL DMEM, 2.5 ml heat inactivated FBS, 10 mL HEPES and 5.5 mL L-Glutamine.
8. PBS.
9. Flat Bottom 96-well Tissue Culture Micro Titer Plates (Corning Catalog # 25860).
- 10 10. 15 cm Tissue Culture Dishes (Corning Catalog #08757148).
11. Corning 96-well ELISA Plates.
12. NUNC 96-well V bottom polypropylene plates.
13. Costar Transfer Cartidges for the Transtar 96 (Costar Catalog #7610).
14. SUMO 1: monoclonal anti-EGFR antibody.
- 15 15. TBST Buffer
16. Blocking Buffer : 5% Carnation Instant Milk® in PBS.
17. EGF Ligand: EGF-201, Shinko American, Japan. Suspend powder in 100 µL of 10 mM HCl. Add 100 µL 10mM NaOH. Add 800 µL PBS and transfer to an Eppendorf tube for storage at -20 °C.
- 20 18. HNTG Lysis Buffer:
For Stock 5X HNTG: Mix 23.83 g Hepes, 43.83 g NaCl, 500 mL glycerol and 100 mL Triton X-100 and enough dH₂O to make 1 L of total solution.
For 1X HNTG*: Mix 2 mL HNTG, 100 µL 0.1 M Na₃VO₄, 250 µL 0.2M Na₄P₂O₇ and 100 µL EDTA.
- 25 19. EDTA
20. Na₃VO₄:
To make stock solution: Mix 1.84 g Na₃VO₄ with 90 mL dH₂O. Adjust pH to 10. Boil in microwave for one minute (solution becomes clear). Cool to room temperature. Adjust pH to 10. Repeat heating/cooling cycle until pH remains at 10.
- 30 21. 200 mM Na₄P₂O₇.

- 5

10

- 15

- 20

- 25

- 30

then transferred to the cell plate at a 1:10 dilution (10 μ L of sample and media into 90 μ L of starve media). The final DMSO concentration will be 1% or lower.

Secondary screening :

Ten samples are put into wells 2-11 of row A of a polypropylene plate.

- 5 These wells contain straight starve-DMEM. For a 1:10 dilution, use 10 μ L of test compound solution in 90 μ L of media. The rest of the wells (including control) will have a DMSO/media mixture. The percentage of DMSO in this mixture is determined by the first dilution factor, e.g., in this example, 1:10. The DMSO concentration is therefore 10%. An equal amount of drug and media from row A is
- 10 put into row B, containing DMSO and media. The same amount is then taken out and put into row C, etc. These are 1:2 dilutions. All wells are then transferred to the cell plate at 1:10 dilution (10 μ L of sample and media into 90 μ L of starve media). The final DMSO concentration will be 1% or lower.

9. Incubate under 5% CO₂ at 37 °C for 2 hours.
- 15 10. Prepare EGF ligand by diluting stock EGF (16.5 μ M) in warm DMEM to 150 nM.
11. Prepare fresh HNTG* sufficient for 100 μ L per well; place on ice.
12. After 2 hour incubation with test compound, add prepared EGF ligand to cells, 50 μ L per well, for a final concentration of 50 nM. Positive control
- 20 wells receive the same amount of EGF. Negative controls do not receive EGF. Incubate at 37 °C for 10 min.
13. Remove test compound, EGF, and DMEM. Wash cells once with PBS.
14. Transfer HNTG* to cells, 100 μ L per well. Place on ice for 5
- 25 minutes. Meanwhile, remove blocking buffer from ELISA plate and wash.
15. With a pipette tip securely fitted to a micropipettor, scrape cells from plate and homogenize cell material by repeatedly aspirating and dispensing the HNTG* lysis buffer. Transfer lysate to a coated, blocked, washed ELISA plate. Alternatively, one may use a Costar transfer cartridge to transfer lysate to the ELISA
- 30 plate.
16. Incubate, with shaking, at room temperature for one hr.

17. Remove lysate, wash. Transfer freshly diluted anti-Ptyr antibody (1:3000 in TBST) to ELISA plate, 100 μ L per well.
18. Incubate, with shaking, at room temperature, for 30 min.
19. Remove anti-Ptyr antibody, wash. Transfer freshly diluted
- 5 BIOSOURCE antibody to ELISA plate(1:8000 in TBST, 100 μ L per well).
20. Incubate, with shaking, at room temperature for 30 min.
21. Remove BIOSOURCE antibody, wash. Transfer freshly prepared ABTS/H₂O₂ solution to ELISA plate, 100 μ L per well.
22. Incubate, with shaking, for 5-10 minutes. Remove any bubbles.
- 10 23. If necessary, stop reaction with the addition of μ L of 0.2 M HCl per well.
24. Read assay on Dynatech MR7000 ELISA reader: test filter set at 410 nM; reference filter at 630 nM.

15 EXAMPLE 110: CDK2/CYCLIN A ASSAY

The following protocol describes the procedures used to analyze protein serine/threonine kinase activity of cdk2/cyclin A in an SPA. The procedure also describes the protocol for the initial screening of drugs for inhibition or activation of the kinase activity.

20

Materials and Reagents:

1. Wallac 96-well polyethylene terephthalate (flexi) plates (Wallac Catalog # 1450-401).
2. Amersham Redivue [γ^{33} P] ATP (Amersham catalog #AH 9968).
- 25 3. Amersham streptavidin coated polyvinyltoluene SPA beads (Amersham catalog #RPNQ0007). Reconstitute beads in PBS without magnesium or calcium, at 20 mg/mL. Store reconstituted beads at 4 °C.
4. Activated cdk2/cyclin A enzyme complex purified from Sf9 cells, -80 °C, 200 μ L aliquots

5. Biotinylated peptide substrate (deb-tide). Peptide biotin-X-PKTPKKAKKL dissolved in dH₂O at a concentration of 5 mg/mL. Stored at -80 °C in 100 µL aliquots.

6. Peptide/ATP Mixture:

5

Reagent	Stock Concentration	2.5 X Working Concentration	Amount per 10 mL	Final Well Concentration
dH ₂ O		10 mL	9.979 mL	---
Cold ATP	10 mM	1.25 µM	0.00125 mL	0.5 µM
Debtide	5 mg/mL	0.005 mg/mL	0.010 mL	0.1 µg/well
γ ³³ P ATP	10 µCi/µL	10 µCi/mL	0.010 mL	0.2 µCi/well

7. 2.5 X kinase buffer

Reagent	Stock solution	Amount per 10 mL	Working Concentration	Final Well Concentration
dH ₂ O	55.5 M	8.85 mL		-----
Tris pH7.4	1 M	0.625 mL	62.5 mM	25 mM TRIS
MgCl ₂	1 M	0.25 mL	25 mM	10 mM MgCl ₂
NP40	10%	0.25 mL	0.25%	0.1 % NP40
*DTT add fresh	1 M	0.025 mL	2.5 mM	1 mM DTT

8. 10 mM ATP (Sigma Catalog # A-5394).

10

9. 1 M Tris, pH 7.4

10. 1 M MgCl₂

11. 1 M DTT

12. PBS (Dulbecco's Phosphate-Buffered Saline) without magnesium or calcium (Gibco Catalog # 14190-144)

15

13. EDTA (14.12 g per 100 mL).

14. Stop solution:

Reagent	Stock solution	Amount per 10 mL	Working Concentration
PBS		9.25 mL	
ATP	100 mM	0.005 mL	50 uM
EDTA	0.5 M	0.1 mL	5 mM
Triton X-100	10%	0.1 mL	0.1%
SPA beads	20 mg/mL	1.25 mL	0.5 mg/well (200 µL)

Procedure:

1. Prepare solutions of inhibitors at 5x the desired final concentration in 5% DMSO. Add 10 μ L to each well. For negative controls, add 10 μ L 5% DMSO.
2. Dilute 5 μ L of cdk2/cyclin A solution into 2.1 mL 2x kinase buffer (per plate).
3. Add 20 μ L enzyme per well. This can be added using a hand pipette or by using the Titertek Multidrop.
4. Add 10 μ L of 0.5 M EDTA to the neagive control wells.
5. To start kinase reaction, add 20 μ L of peptide/ATP mixture using either a hand pipette or the Titertek Multidrop. Let sit on benchtop behind reactive shield for 1 hr.
6. Add 200 μ L stop solution per well using either the Titertek Multidrop or hand pipette.
7. Let stand at least 10 min.
8. Spin plate approx. 2300 rpm 3-5 min.
9. Count plate on Trilux reader using protocol #28 (Brian's SPA assay).

EXAMPLE 111: PDGF-R ELISA

20 All cell culture media, glutamine, and fetal bovine serum were purchased from Gibco Life Technologies (Grand Island, NY) unless otherwise specified. All cells were grown in a humid atmosphere of 90-95% air and 5-10% CO₂ at 37 °C. All cell lines were routinely subcultured twice a week and were negative for mycoplasma as determined by the Mycotect method (Gibco).

25 For ELISA assays, cells (U1242, obtained from Joseph Schlessinger, NYU) were grown to 80-90% confluency in growth medium (MEM with 10% FBS, NEAA, 1 mM NaPyr and 2 mM GLN) and seeded in 96-well tissue culture plates in 0.5% serum at 25,000 to 30,000 cells per well. After overnight incubation in 0.5% serum-containing medium, cells were changed to serum-free medium and treated

30 with test compound for 2 hr in a 5% CO₂, 37 °C incubator. Cells were then stimulated with ligand for 5-10 minute followed by lysis with HNTG (20 mM

Hepes, 150 mM NaCl, 10% glycerol, 5 mM EDTA, 5 mM Na₃VO₄, 0.2% Triton X-100, and 2 mM NaPyr). Cell lysates (0.5 mg/well in PBS) were transferred to ELISA plates previously coated with receptor-specific antibody and which had been blocked with 5% milk in TBST (50 mM Tris-HCl pH 7.2, 150 mM NaCl and 0.1% Triton X-100) at room temperature for 30 min. Lysates were incubated with shaking for 1 hour at room temperature. The plates were washed with TBST four times and then incubated with polyclonal anti-phosphotyrosine antibody at room temperature for 30 minutes. Excess anti-phosphotyrosine antibody was removed by rinsing the plate with TBST four times. Goat anti-rabbit IgG antibody was added to the ELISA plate for 30 min at room temperature followed by rinsing with TBST four more times. ABTS (100 mM citric acid, 250 mM Na₂HPO₄ and 0.5 mg/mL 2,2'-azino-bis(3-ethylbenzthiazoline-6-sulfonic acid)) plus H₂O₂ (1.2 mL 30% H₂O₂ to 10 mL ABTS) was added to the ELISA plates to start color development. Absorbance at 410 nm with a reference wavelength of 630 nm was recorded about 15 to 30 min after ABTS addition.

EXAMPLE 112: IGF-I RECEPTOR ELISA

The following protocol may be used to measure phosphotyrosine level on IGF-I receptor, which indicates IGF-I receptor tyrosine kinase activity.

Materials and Reagents

The following materials and reagents were used:

- a. The cell line used in this assay is 3T3/IGF-1R, a cell line genetically engineered to overexpresses IGF-1 receptor.
- b. NIH3T3/IGF-1R is grown in an incubator with 5% CO₂ at 37 °C. The growth media is DMEM + 10% FBS (heat inactivated)+ 2 mM L-glutamine.
- c. Affinity purified anti-IGF-1R antibody 17-69.
- d. D-PBS:

	KH ₂ PO ₄	0.20 g/L
	K ₂ HPO ₄	2.16 g/L
	KCl	0.20 g/L

NaCl 8.00 g/L(pH 7.2)

e. Blocking Buffer: TBST plus 5% Milk (Carnation Instant Non-Fat Dry Milk).

f. TBST buffer:

5 Tris-HCl 50 mM
NaCl 150 mM (pH 7.2/HCl 10 N)
Triton X-100 0.1%

Stock solution of TBS (10X) is prepared, and Triton X-100 is added to the buffer during dilution.

10

g. HNTG buffer:

HEPES 20 mM
NaCl 150 mM (pH 7.2/HCl 1N)
Glycerol 10%
15 Triton X-100 0.2%

Stock solution (5X) is prepared and kept at 4 °C.

h. EDTA/HCl: 0.5 M pH 7.0 (NaOH) as 100X stock.

i. Na_3VO_4 : 0.5 M as 100X stock and aliquots are kept in -80 °C.

20 j. $\text{Na}_4\text{P}_2\text{O}_7$: 0.2 M as 100X stock.

k. Insulin-like growth factor-1 from Promega (Cat# G5111).

l. Rabbit polyclonal anti-phosphotyrosine antiserum.

m. Goat anti-rabbit IgG, POD conjugate (detection antibody), Tago (Cat. No. 4520, Lot No. 1802): Tago, Inc., Burlingame, CA.

25 n. ABTS (2,2'-azinobis(3-ethylbenzthiazolinesulfonic acid)) solution:

Citric acid 100 mM
 Na_2HPO_4 250 mM (pH 4.0/1 N HCl)
ABTS 0.5 mg/mL

30 ABTS solution should be kept in dark and 4°C. The solution should be discarded when it turns green.

- o. Hydrogen Peroxide: 30% solution is kept in the dark and at 4 °C.

Procedure

All the following steps are conducted at room temperature unless it is specifically indicated. All ELISA plate washings are performed by rinsing the plate with tap water three times, followed by one TBST rinse. Pat plate dry with paper towels.

A. Cell Seeding:

1. The cells, grown in tissue culture dish (Corning 25020-100) to 80-90% confluence, are harvested with Trypsin-EDTA (0.25%, 0.5 mL/D-100, GIBCO).
2. Resuspend the cells in fresh DMEM + 10% FBS + 2 mM L-Glutamine, and transfer to 96-well tissue culture plate (Corning, 25806-96) at 20,000 cells/well (100 µL/well). Incubate for 1 day then replace medium to serum-free medium (90/µL) and incubate in 5% CO₂ and 37 °C overnight.

B. ELISA Plate Coating and Blocking:

1. Coat the ELISA plate (Corning 25805-96) with Anti-IGF-1R Antibody at 0.5 µg/well in 100 µL PBS at least 2 hours.
2. Remove the coating solution, and replace with 100 µL Blocking Buffer, and shake for 30 minutes. Remove the blocking buffer and wash the plate just before adding lysate.

C. Assay Procedures:

1. The drugs are tested in serum-free condition.
2. Dilute drug stock (in 100% DMSO) 1:10 with DMEM in 96-well poly-propylene plate, and transfer 10 µL/well of this solution to the cells to achieve final drug dilution 1:100, and final DMSO concentration of 1.0%. Incubate the cells in 5% CO₂ at 37 °C for 2 hours.
3. Prepare fresh cell lysis buffer (HNTG*)

	HNTG	2 mL
	EDTA	0.1 mL
	Na ₃ VO ₄	0.1 mL

Na₄(P₂O₇) 0.1 mL

H₂O 7.3 mL

4. After drug incubation for two hours, transfer 10 μ l/well of 200nM IGF-1 Ligand in PBS to the cells (Final Conc. = 20 nM), and incubate at 5% CO₂ at 37 °C for 10 minutes.

5. Remove media and add 100 μ L/well HNTG* and shake for 10 minutes. Look at cells under microscope to see if they are adequately lysed.

6. Use a 12-channel pipette to scrape the cells from the plate, and homogenize the lysate by repeated aspiration and dispensing. Transfer all the lysate to the antibody coated ELISA plate, and shake for 1 hour.

7. Remove the lysate, wash the plate, transfer anti-pTyr (1:3,000 with TBST) 100 μ L/well, and shake for 30 minutes.

8. Remove anti-pTyr, wash the plate, transfer TAGO (1:3,000 with TBST) 100 μ L/well, and shake for 30 minutes.

9. Remove detection antibody, wash the plate, and transfer fresh ABTS/H₂O₂ (1.2 μ L H₂O₂ to 10 ml ABTS) 100 μ L/well to the plate to start color development.

10. Measure OD at 410 nm with a reference wavelength of 630 nm in Dynatec MR5000.

EXAMPLE 113: EGF Receptor ELISA

EGF Receptor kinase activity in cells genetically engineered to express human EGF-R was measured as described below:

Materials and Reagents

The following materials and reagents were used:

a. EGF Ligand: stock concentration = 16.5 μ M; EGF 201, TOYOBO, Co., Ltd. Japan.

b. 05-101 (UBI) (a monoclonal antibody recognizing an EGFR extracellular domain).

c. Anti-phosphotyrosine antibody (anti-Ptyr) (polyclonal).

d. Detection antibody: Goat anti-rabbit IgG horse radish peroxidase conjugate, TAGO, Inc., Burlingame, CA.

e. TBST buffer:

	Tris-HCl, pH 7	50 mM
5	NaCl	150 mM
	Triton X-100	0.1

f. HNTG 5X stock:

	HEPES	0.1 M
	NaCl	0.75 M
10	Glycerol	50
	Triton X-100	1.0%

g. ABTS stock:

	Citric Acid	100 mM
	Na ₂ HPO ₄	250 mM
15	HCl, conc.	4.0 pH
	ABTS*	0.5 mg/mL

Keep solution in dark at 4 °C until used.

h. Stock reagents of:

	EDTA 100 mM pH 7.0
20	Na ₃ VO ₄ 0.5 M
	Na ₄ (P ₂ O ₇) 0.2 M

Procedure

The following protocol was used:

25 A. Pre-coat ELISA Plate

1. Coat ELISA plates (Corning, 96 well, Cat. #25805-96) with 05-101 antibody at 0.5 µg per well in PBS, 150 µL final volume/well, and store overnight at 4 °C. Coated plates are good for up to 10 days when stored at 4 °C.
 2. On day of use, remove coating buffer and replace with blocking buffer (5% Carnation Instant Non-Fat Dry Milk in PBS). Incubate the plate, shaking,
- 30

B. Seeding Cells

2. Choose dishes having 80-90% confluence for the experiment.

10 3. Resuspend cells in seeding medium (DMEM, 0.5% bovine serum),
and count the cells using trypan blue. Viability above 90% is acceptable. Seed cells
in DMEM medium (0.5% bovine serum) at a density of 10,000 cells per well, 100
μL per well, in a 96 well microtiter plate. Incubate seeded cells in 5% CO₂ at 37 °C
for about 40 hours.

1. Check seeded cells for contamination using an inverted microscope.

20 2. Prepare EGF ligand: dilute stock EGF in DMEM so that upon
transfer of 10 μ L dilute EGF (1:12 dilution), 25 nM final concentration is attained.

- 25 4. Place on ice.

- Remove drug, EGF, and DMEM. Wash cells twice with PBS.

- 137

remove blocking buffer from other ELISA plate and wash with TBST as described above.

7. With a pipette tip securely fitted to a micropipettor, scrape cells from plate and homogenize cell material by repeatedly aspirating and dispensing the HNTG* lysis buffer. Transfer lysate to a coated, blocked, and washed ELISA plate. Incubate shaking at room temperature for one hour.

8. Remove lysate and wash 4 times with TBST. Transfer freshly diluted anti-Ptyr antibody to ELISA plate at 100 μ L per well. Incubate shaking at room temperature for 30 minutes in the presence of the anti-Ptyr antiserum (1:3000 dilution in TBST).

9. Remove the anti-Ptyr antibody and wash 4 times with TBST. Transfer the freshly diluted TAGO 30 anti-rabbit IgG antibody to the ELISA plate at 100 μ L per well. Incubate shaking at room temperature for 30 minutes (anti-rabbit IgG antibody: 1:3000 dilution in TBST).

10. Remove detection antibody and wash 4 times with TBST. Transfer freshly prepared ABTS/H₂O₂ solution to ELISA plate, 100 μ L per well. Incubate at room temperature for 20 minutes. ABTS/H₂O₂ solution: 1.2 μ L 30% H₂O₂ in 10 mL ABTS stock.

11. Stop reaction by adding 50 μ L 5N H₂SO₄ (optional), and determine O.D. at 410 nm.

12. The maximal phosphotyrosine signal is determined by subtracting the value of the negative controls from the positive controls. The percent inhibition of phosphotyrosine content for extract-containing wells is then calculated, after subtraction of the negative controls.

EXAMPLE 114: Met Autophosphorylation Assay - ELISA

This assay determines Met tyrosine kinase activity by analyzing Met protein tyrosine kinase levels on the Met receptor.

Materials and Reagents

The following materials and reagents were used:

- a. HNTG (5X stock solution): Dissolve 23.83 g HEPES and 43.83 g NaCl in about 350 mL dH₂O. Adjust pH to 7.2 with HCl or NaOH, add 500 mL glycerol and 10 mL Triton X-100, mix, add dH₂O to 1 L total volume. To make 1 L of 1X working solution add 200 mL 5X stock solution to 800 mL dH₂O, check and
5 adjust pH as necessary, store at 4 °C.
- b. PBS (Dulbecco's Phosphate-Buffered Saline), Gibco Cat. # 450-1300EB (1X solution).
- c. Blocking Buffer: in 500 ml dH₂O place 100 g BSA, 12.1 g Tris-pH7.5, 58.44 g NaCl and 10 mL Tween-20, dilute to 1 L total volume.
- 10 d. Kinase Buffer: To 500 mL dH₂O add 12.1 g TRIS pH7.2, 58.4 g NaCl, 40.7 g MgCl₂ and 1.9 g EGTA; bring to 1 L total volume with dH₂O.
- e. PMSF (Phenylmethylsulfonyl fluoride), Sigma Cat. # P-7626, to 435.5 mg, add 100% ethanol to 25 mL total volume, vortex.
- f. ATP (Bacterial Source), Sigma Cat. # A-7699, store powder at -20
15 °C; to make up solution for use, dissolve 3.31 mg in 1 mL dH₂O.
- g. RC-20H HRPO Conjugated Anti-Phosphotyrosine, Transduction Laboratories Cat. # E120H.
- h. Pierce 1-Step (TM) Turbo TMB-ELISA (3,3',5,5'-tetramethylbenzidine, Pierce Cat. # 34022.
- 20 i. H₂SO₄, add 1 mL conc. (18 N) to 35 mL dH₂O.
- j. TRIS HCL, Fischer Cat. # BP152-5; to 121.14 g of material, add 600 mL MilliQ H₂O, adjust pH to 7.5 (or 7.2) with HCl , bring volume to 1 L with MilliQ H₂O.
- k. NaCl, Fischer Cat. # S271-10, make up 5 M solution.
- 25 l. Tween-20, Fischer Cat. # S337-500.
- m. Na₃VO₄, Fischer Cat. # S454-50, to 1.8 g material add 80 ml MilliQ H₂O, adjust pH to 10.0 with HCl or NaOH, boil in microwave, cool, check pH, repeat procedure until pH stable at 10.0, add MilliQ H₂O to 100 ml total volume, make 1 mL aliquots and store at -80 °C.
- 30 n. MgCl₂, Fischer Cat. # M33-500, make up 1 M solution.

o. HEPES, Fischer Cat. # BP310-500, to 200 ml MilliQ H₂O, add 59.6 g material, adjust pH to 7.5, bring volume to 250 mL total, sterile filter.

p. Albumin, Bovine (BSA), Sigma Cat. # A-4503, to 30 grams material add sterile distilled water to make total volume of 300 mL, store at 4 °C.

5 q. TBST Buffer: to approx. 900 mL dH₂O in a 1 L graduated cylinder add 6.057 g TRIS and 8.766 g NaCl, when dissolved, adjust pH to 7.2 with HCl, add 1.0 mL Triton X-100 and bring to 1 L total volume with dH₂O.

r. Goat Affinity purified antibody Rabbit IgG (whole molecule), Cappel Cat. # 55641.

10 s. Anti h-Met (C-28) rabbit polyclonal IgG antibody, Santa Cruz Chemical Cat. # SC-161.

t. Transiently Transfected EGFR/Met chimeric cells (EMR) (Komada, et al., Oncogene, 8:2381-2390 (1993).

15 u. Sodium Carbonate Buffer, (Na₂CO₄, Fischer Cat. # S495): to 10.6 g material add 800 mL MilliQ H₂O, when dissolved adjust pH to 9.6 with NaOH, bring up to 1 L total volume with MilliQ H₂O, filter, store at 4 °C.

Procedure

20 All of the following steps are conducted at room temperature unless it is specifically indicated otherwise. All ELISA plate washing is by rinsing 4X with TBST.

A. EMR Lysis

25 This procedure can be performed the night before or immediately prior to the start of receptor capture.

1. Quick thaw lysates in a 37 °C waterbath with a swirling motion until the last crystals disappear.

2. Lyse cell pellet with 1X HNTG containing 1 mM PMSF. Use 3 ml of HNTG per 15 cm dish of cells. Add ½ the calculated HNTG volume, vortex the
30 tube for 1 min., add the remaining amount of HNTG, vortex for another min.

3. Balance tubes, centrifuge at 10,000x g for 10 min at 4 °C.

4. Pool supernatants, remove an aliquot for protein determination.
 5. Quick freeze pooled sample in dry ice/ethanol bath. This step is performed regardless of whether lysate will be stored overnight or used immediately following protein determination.

5 6. Perform protein determination using standard bicinchoninic acid (BCA) method (BCA Assay Reagent Kit from Pierce Chemical Cat. # 23225).

B. ELISA Procedure

1. Coat Corning 96 well ELISA plates with 5 µg per well Goat anti-Rabbit antibody in Carbonate Buffer for a total well volume of 50 µL. Store overnight at 4 °C.

2. Remove unbound Goat anti-rabbit antibody by inverting plate to remove liquid.

3. Add 150 µL of Blocking Buffer to each well. Incubate for 30 min. at room temperature with shaking.

4. Wash 4X with TBST. Pat plate on a paper towel to remove excess liquid and bubbles.

5. Add 1 µg per well of Rabbit anti-Met antibody diluted in TBST for a total well volume of 100 µL.

6. Dilute lysate in HNTG (90 µg lysate/100 µL)

7. Add 100 µL of diluted lysate to each well. Shake at room temperature for 60 min.

8. Wash 4X with TBST. Pat on paper towel to remove excess liquid and bubbles.

9. Add 50 µL of 1X lysate buffer per well.

10. Dilute compounds/extracts 1:10 in 1X Kinase Buffer in a polypropylene 96 well plate.

11. Transfer 5.5 µL of diluted drug to ELISA plate wells. Incubate at room temperature with shaking for 20 min.

12. Add 5.5 µL of 60 µM ATP solution per well. Negative controls do not receive any ATP. Incubate at room temperature for 90 min., with shaking.

13. Wash 4X with TBST. Pat plate on paper towel to remove excess liquid and bubbles.

14. Add 100 μ L per well of RC20 (1:3000 dilution in Blocking Buffer). Incubate 30 min. at room temperature with shaking.

5 15. Wash 4X with TBST. Pat plate on paper towel to remove excess liquid and bubbles.

16. Add 100 μ L per well of Turbo-TMB. Incubate with shaking for 30-60 min.

17. Add 100 μ L per well of 1 M H_2SO_4 to stop reaction.

10 18. Read assay on Dynatech MR7000 ELISA reader. Test Filter = 450 nm, reference filter = 410 nm.

EXAMPLE 115: Biochemical src assay - ELISA

15 This assay is used to determine src protein kinase activity measuring phosphorylation of a biotinylated peptide as the readout.

Materials and Reagents

The following materials and reagents were used:

- a. Yeast transformed with.
- 20 b. Cell lysates: Yeast cells expressing src are pelleted, washed once with water, re-pelleted and stored at -80 °C until use.
- c. N-terminus biotinylated EEEYEEYEEYEEYEEY is prepared by standard procedures well known to those skilled in the art.
- d. DMSO: Sigma, St. Louis, MO.
- 25 e. 96 Well ELISA Plate: Corning 96 Well Easy Wash, Modified flat Bottom Plate, Corning Cat. #25805-96.
- f. NUNC 96-well V-bottom polypropylene plates for dilution of compounds: Applied Scientific Cat. # A-72092.
- g. Vecastain ELITE ABC reagent: Vector, Burlingame, CA.
- 30 h. Anti-src (327) mab: Schizosaccharomyces Pombe was used to express recombinant Src (Superti-Furga, et al., EMBO J., 12:2625-2634; Superti-

Furga, et al., Nature Biochem., 14:600-605). S. Pombe strain SP200 (h-s leu1.32 ura4 ade210) was grown as described and transformations were pRSP expression plasmids were done by the lithium acetate method (Superti-Furga, supra). Cells were grown in the presence of 1 μ M thiamine to repress expression from the nmt1 promoter or in the absence of thiamine to induce expression.

i. Monoclonal anti-phosphotyrosine, UBI 05-321 (UB40 may be used instead).

j. Turbo TMB-ELISA peroxidase substrate: Pierce Chemical.

10 Buffer Solutions:

a. PBS (Dulbecco's Phosphate-Buffered Saline): GIBCO PBS, GIBCO Cat. # 450-1300EB.

b. Blocking Buffer: 5% Non-fat milk (Carnation) in PBS.

c. Carbonate Buffer: Na_2CO_4 from Fischer, Cat. # S495, make up 100 mM stock solution.

d. Kinase Buffer: 1.0 mL (from 1 M stock solution) MgCl_2 ; 0.2 mL (from a 1 M stock solution) MnCl_2 ; 0.2 mL (from a 1 M stock solution) DTT; 5.0 mL (from a 1 M stock solution) HEPES; 0.1 mL TX-100; bring to 10 mL total volume with MilliQ H_2O .

e. Lysis Buffer: 5.0 HEPES (from 1 M stock solution.); 2.74 mL NaCl (from 5 M stock solution); 10 mL glycerol; 1.0 mL TX-100; 0.4 mL EDTA (from a 100 mM stock solution); 1.0 mL PMSF (from a 100 mM stock solution); 0.1 mL Na_3VO_4 (from a 0.1 M stock solution); bring to 100 mL total volume with MilliQ H_2O .

f. ATP: Sigma Cat. # A-7699, make up 10 mM stock solution (5.51 mg/mL).

g. TRIS-HCl: Fischer Cat. # BP 152-5, to 600 mL MilliQ H_2O add 121.14 g material, adjust pH to 7.5 with HCl, bring to 1 L total volume with MilliQ H_2O .

h. NaCl: Fischer Cat. # S271-10, Make up 5M stock solution with MilliQ H_2O .

- i. Na_3VO_4 : Fischer Cat. # S454-50; to 80 mL MilliQ H_2O , add 1.8 g material; adjust pH to 10.0 with HCl or NaOH; boil in a microwave; cool; check pH, repeat pH adjustment until pH remains stable after heating/cooling cycle; bring to 100 mL total volume with MilliQ H_2O ; make 1 mL aliquots and store at -80°C .
- 5 j. MgCl_2 : Fischer Cat. # M33-500, make up 1M stock solution with MilliQ H_2O .
- k. HEPES: Fischer Cat. # BP 310-500; to 200 mL MilliQ H_2O , add 59.6 g material, adjust pH to 7.5, bring to 250 ml total volume with MilliQ H_2O , sterile filter (1 M stock solution).
- 10 l. TBST Buffer: TBST Buffer: To 900 mL dH_2O add 6.057 g TRIS and 8.766 g NaCl; adjust pH to 7.2 with HCl, add 1.0 mL Triton-X100; bring to 1 L total volume with dH_2O .
- m. MnCl_2 : Fischer Cat. # M87-100, make up 1 M stock solution with MilliQ H_2O .
- 15 n. DTT; Fischer Cat. # BP172-5.
- o. TBS (TRIS Buffered Saline): to 900 mL MilliQ H_2O add 6.057 g TRIS and 8.777 g NaCl; bring to 1 L total volume with MilliQ H_2O .
- p. Kinase Reaction Mixture: Amount per assay plate (100 wells): 1.0 mL Kinase Buffer, 200 μg GST- ζ , bring to final volume of 8.0 mL with MilliQ H_2O .
- 20 q. Biotin labeled EEEYEEYEEEEEEEEEEY: Make peptide stock solution (1 mM, 2.98 mg/ml) in water fresh just before use.
- r. Vectastain ELITE ABC reagent: To prepare 14 mL of working reagent, add 1 drop of reagent A to 15 mL TBST and invert tube several times to mix. Then add 1 drop of reagent B. Put tube on orbital shaker at room temperature
- 25 and mix for 30 minutes.

Procedures

- a. Preparation of src coated ELISA plate.
 1. Coat ELISA plate with 0.5 μg /well anti-src mab in 100 μL of pH 9.6 sodium carbonate buffer at 4°C overnight.
 2. Wash wells once with PBS.
- 30

- 30

EXAMPLE 116: Biochemical lck Assay - ELISA

This assay is used to determine lck protein kinase activities measuring phosphorylation of GST- ζ as the readout.

5 Materials and Reagents

The following materials and reagents were used:

- a. Yeast transformed with lck. Schizosaccharomyces Pombe was used to express recombinant Lck (Superti-Furga, et al., EMBO J, 12:2625-2634; Superti-Furga, et al., Nature Biotech., 14:600-605). S. Pombe strain SP200 (h-s leul.32 ura4 ade210) was grown as described and transformations with pRSP expression plasmids were done by the lithium acetate method (Superti-Furga, supra). Cells were grown in the presence of 1 μ M thiamine to induce expression.
- b. Cell lysates: Yeast cells expressing lck are pelleted, washed once in water, re-pelleted and stored frozen at -80 °C until use.
- 15 c. GST- ζ : DNA encoding for GST- ζ fusion protein for expression in bacteria obtained from Arthur Weiss of the Howard Hughes Medical Institute at the University of California, San Francisco. Transformed bacteria were grown overnight while shaking at 25 °C. GST- ζ was purified by glutathione affinity chromatography, Pharmacia, Alameda, CA.
- 20 d. DMSO: Sigma, St. Louis, MO.
- e. 96-Well ELISA plate: Corning 96 Well Easy Wash, Modified Flat Bottom Plate, Corning Cat. #25805-96.
- f. NUNC 96-well V-bottom polypropylene plates for dilution of compounds: Applied Scientific Cat. # AS-72092.
- 25 g. Purified Rabbit anti-GST antiserum: Amrad Corporation (Australia) Cat. #90001605.
- h. Goat anti-Rabbit-IgG-HRP: Amersham Cat. # V010301
- i. Sheep ant-mouse IgG (H+L): Jackson Labs Cat. # 5215-005-003.
- j. Anti-Lck (3A5) mab: Santa Cruz Biotechnology Cat # sc-433.
- 30 k. Monoclonal anti-phosphotyrosine UBI 05-321 (UB40 may be used instead).

Buffer solutions:

- a. PBS (Dulbecco's Phosphate-Buffered Saline) 1X solution: GIBCO PBS, GIBCO Cat. # 450-1300EB.
- b. Blocking Buffer: 100 g. BSA, 12.1 g. TRIS-pH7.5, 58.44 g NaCl, 10 mL Tween-20, bring up to 1 L total volume with MilliQ H₂O.
- c. Carbonate Buffer: Na₂CO₄ from Fischer, Cat. # S495; make up 100 mM solution with MilliQ H₂O.
- d. Kinase Buffer: 1.0 mL (from 1 M stock solution) MgCl₂; 0.2 mL (from a 1 M stock solution) MnCl₂; 0.2 mL (from a 1 M stock solution) DTT; 5.0 mL (from a 1 M stock solution) HEPES; 0.1 mL TX-100; bring to 10 mL total volume with MilliQ H₂O.
- e. Lysis Buffer: 5.0 HEPES (from 1 M stock solution.); 2.74 mL NaCl (from 5 M stock solution); 10 mL glycerol; 1.0 mL TX-100; 0.4 mL EDTA (from a 100 mM stock solution); 1.0 mL PMSF (from a 100 mM stock solution); 0.1 mL Na₃VO₄ (from a 0.1 M stock solution); bring to 100 mL total volume with MilliQ H₂O.
- f. ATP: Sigma Cat. # A-7699, make up 10 mM stock solution (5.51 mg/mL).
- g. TRIS-HCl: Fischer Cat. # BP 152-5, to 600 mL MilliQ H₂O add 121.14 g material, adjust pH to 7.5 with HCl, bring to 1 L total volume with MilliQ H₂O.
- h. NaCl: Fischer Cat. # S271-10, Make up 5 M stock solution with MilliQ H₂O.
- i. Na₃VO₄: Fischer Cat. # S454-50; to 80 mL MilliQ H₂O, add 1.8 g material; adjust pH to 10.0 with HCl or NaOH; boil in a microwave; cool; check pH, repeat pH adjustment until pH remains stable after heating/cooling cycle; bring to 100 ml total volume with MilliQ H₂O; make 1 ml aliquots and store at -80 °C.
- j. MgCl₂: Fischer Cat. # M33-500, make up 1M stock solution with MilliQ H₂O.

k. HEPES: Fischer Cat. # BP 310-500; to 200 mL MilliQ H₂O, add 59.6 g material, adjust pH to 7.5, bring to 250 mL total volume with MilliQ H₂O, sterile filter (1M stock solution).

l. Albumin, Bovine (BSA), Sigma Cat. # A4503; to 150 mL MilliQ H₂O add 30 g material, bring 300 mL total volume with MilliQ H₂O, filter through 0.22 µm filter, store at 4 °C.

m. TBST Buffer: To 900 mL dH₂O add 6.057 g TRIS and 8.766 g NaCl; adjust pH to 7.2 with HCl, add 1.0 mL Triton-X100; bring to 1 L total volume with dH₂O.

n. MnCl₂: Fischer Cat. # M87-100, make up 1 M stock solution with MilliQ H₂O.

o. DTT; Fischer Cat. # BP172-5.

p. TBS (TRIS Buffered Saline): to 900 mL MilliQ H₂O add 6.057 g TRIS and 8.777 g NaCl; bring to 1 L total volume with MilliQ H₂O.

q. Kinase Reaction Mixture: Amount per assay plate (100 wells): 1.0 mL Kinase Buffer, 200 µg GST-ζ, bring to final volume of 8.0 mL with MilliQ H₂O.

Procedures

a. Preparation of Lck coated ELISA plate.

1. Coat 2.0 µg/well Sheep anti-mouse IgG in 100 µL of pH 9.6 sodium carbonate buffer at 4 °C overnight.

2. Wash well once with PBS.

3. Block plate with 0.15 mL of blocking Buffer for 30 min. at room temp.

4. Wash plate 5X with PBS.

5. Add 0.5 µg/well of anti-lck (mab 3A5) in 0.1 mL PBS at room temperature for 1-2 hours.

6. Wash plate 5X with PBS.

7. Add 20 µg/well of lck transformed yeast lysates diluted in Lysis Buffer (0.1 mL total volume per well). (Amount of lysate may vary between batches) Shake plate at 4 °C overnight to prevent loss of activity.

- b. Preparation of phosphotyrosine antibody-coated ELISA plate.
 1. UB40 plate: 1.0 µg/well UB40 in 100 µL of PBS overnight at 4 °C and block with 150 µL of Blocking Buffer for at least 1 hour.
- c. Kinase assay procedure.
 - 5 1. Remove unbound proteins from step 1-7, above, and wash plates 5X with PBS.
 2. Add 0.08 mL Kinase Reaction Mixture per well (containing 10 µL of 10X Kinase Buffer and 2 µg GST-ζ per well diluted with water).
 3. Add 10 µL of compound diluted in water containing 10% DMSO and pre-incubate for 15 minutes at room temperature.
 - 10 4. Start kinase reaction by adding 10 µL/well of 0.1 mM ATP in water (10 µM ATP final).
 5. Shake ELISA plate for 60 min. at room temperature.
 6. Stop kinase reaction by adding 10 µL of 0.5 M EDTA per well.
 - 15 7. Transfer 90 µL supernatant to a blocked 4G10 coated ELISA plate from section B, above.
 8. Incubate while shaking for 30 min. at room temperature.
 9. Wash plate 5X with TBST.
 10. Incubate with Rabbit anti-GST antibody at 1:5000 dilution in 100 µL TBST for 30 min. at room temperature.
 - 20 11. Wash the wells 5X with TBST.
 12. Incubate with Goat anti-Rabbit-IgG-HRP at 1:20,000 dilution in 100 µL of TBST for 30 min. at room temperature.
 13. Wash the wells 5X with TBST.
 - 25 14. Develop with Turbo TMB.

EXAMPLE 117: ASSAY MEASURING PHOSPHORYLATING FUNCTION OF RAF

The following assay reports the amount of RAF-catalyzed phosphorylation of its target protein MEK as well as MEK's target MAPK. The RAF gene sequence is described in Bonner et al., 1985, Molec. Cell. Biol. 5: 1400-

5 Materials and Reagents

- 150

14. Wallac Cellulose Phosphate Filter mats; Wallac, Turku, Finland.
15. Filter wash solution: 1 % phosphoric acid, Fisher, Pittsburgh, PA.
16. Tomtec plate harvester, Wallac, Turku, Finland.
17. Wallac beta plate reader # 1205, Wallac, Turku, Finland.
- 5 18. NUNC 96-well V bottom polypropylene plates for compounds
Applied Scientific Catalog # AS-72092.

Procedure

10 All of the following steps were conducted at room temperature unless specifically indicated.

1. ELISA plate coating: ELISA wells are coated with 100 mL of Sheep anti mouse affinity purified antiserum (1 mg/100 mL coating buffer) over night at 4 °C. ELISA plates can be used for two weeks when stored at 4 °C.
2. Invert the plate and remove liquid. Add 100 mL of blocking solution
15 and incubate for 30 min.
3. Remove blocking solution and wash four times with wash buffer. Pat the plate on a paper towel to remove excess liquid.
4. Add 1 mg of antibody specific for RAF-1 to each well and incubate for 1 hour. Wash as described in step 3.
- 20 5. Thaw lysates from RAS/RAF infected Sf9 cells and dilute with TBST to 10 mg/100 mL. Add 10 mg of diluted lysate to the wells and incubate for 1 hour. Shake the plate during incubation. Negative controls receive no lysate. Lysates from RAS/RAF infected Sf9 insect cells are prepared after cells are infected with recombinant baculoviruses at a MOI of 5 for each virus, and harvested 48 hours
25 later. The cells are washed once with PBS and lysed in RIPA buffer. Insoluble material is removed by centrifugation (5 min at 10 000 x g). Aliquots of lysates are frozen in dry ice/ethanol and stored at -80 °C until use.
6. Remove non-bound material and wash as outlined above (step 3).
7. Add 2 mg of T-MEK and 2 mg of His-MAEPK per well and adjust
30 the volume to 40 mL with kinase buffer. Methods for purifying T-MEK and MAPK from cell extracts are provided herein by example.

8. Pre-dilute compounds (stock solution 10 mg/mL DMSO) or extracts 20 fold in TBST plus 1% DMSO. Add 5 mL of the pre-diluted compounds/extracts to the wells described in step 6. Incubate for 20 min. Controls receive no drug.

9. Start the kinase reaction by addition of 5 mL ATPmix; Shake the plates on an ELISA plate shaker during incubation.

10. Stop the kinase reaction after 60 min by addition of 30 mL stop solution to each well.

11. Place the phosphocellulose mat and the ELISA plate in the Tomtec plate harvester. Harvest and wash the filter with the filter wash solution according to the manufacturers recommendation. Dry the filter mats. Seal the filter mats and place them in the holder. Insert the holder into radioactive detection apparatus and quantify the radioactive phosphorous on the filter mats.

Alternatively, 40 mL aliquots from individual wells of the assay plate can be transferred to the corresponding positions on the phosphocellulose filter mat.

15 After air drying the filters, put the filters in a tray. Gently rock the tray, changing the wash solution at 15 min intervals for 1 hour. Air-dry the filter mats. Seal the filter mats and place them in a holder suitable for measuring the radioactive phosphorous in the samples. Insert the holder into a detection device and quantify the radioactive phosphorous on the filter mats.

20

EXAMPLE 118: FLK-1/KDR ASSAY

EXAMPLE 119: ZAP-70 ASSAY

25 Materials and Reagents

1. Corning 96-well Elisa plates Corning Catalog # 25805-96

2.

Reagent	Vendor	Catalog #	Order quantity
Poly Glu-Tyr (4:1)	Sigma	P 0275	100 mg

3. PBS (Dulbecco's Phosphate-Buffered Saline) Gibco Catalog # 450-1300EB

Reagent	Molecular Weight	10x Stock Concentration	Amt. per L	1x Working Concentration
KCl	74.56	27 mM	2.013 g	2.7 mM
KH ₂ PO ₄	174.18	11 mM	1.916 g	1.1 mM
MgCl ₂ •6H ₂ O (anhydrous)	203.31	5 mM	1.017 g	0.5 mM
NaCl	58.44	1.38M	80.65 g	138 mM
Na ₂ HPO ₄	141.96	81 mM	11.50 g	8.1 mM

5 To make 1 liter of 10x stock solution:

- 1) To a 1 liter graduated cylinder add ~900ml dH₂O
- 2) Add all reagents except the MgCl₂
- 3) When all reagents have dissolved, pH to 7.2 with HCl
- 4) Add MgCl₂
- 5) Bring volume to 1 liter dH₂O

10

One does not necessarily have to make this buffer up. There are two sources of stock PBS:

1) Sterile GIBCO PBS (1x) in 500ml bottles found in the media refrigerator (This is the buffer of choice)

15

2) Sterile 10x and 1x PBS found in the glass cabinets. If this PBS is used the pH must be adjusted to 7.2 with HCl.

To make a 1x working solution from either of the above 10x stocks:

Reagent	Amount per L
10x stock	100 ml
dH ₂ O	900 ml

It is advisable to check the pH after diluting the 10x stock.

*Note-PBS can be left at room temperature, but 4°C is the preferred storage temp.

5

4. 50 mM HEPES

Dilute Gibco Tissue Culture Grade 1 M HEPES to a final concentration of 50 mM HEPES using MilliQue H₂O.

Reagent	M.W.	1xWorking Concentration	Amount per 1 L
1M HEPES		50 mM	50 mL
MilliQue H ₂ O	NA	NA	950 mL

10

5. Blocking Buffer

Reagent	M.W.	10x Stock Concentration	Amount per L	1xWorking Concentration
BSA	NA	10%	100g	10g
TRIS-pH7.5	121.14	100mM	12.1 g	10mM
NaCl	58.44	1 M	58.44 g	100mM
Tween-20	NA	1%	10mL	0.1%

15

6. Purified GST fusion protein containing the Zap70 kinase domain

Biochemistry Lab, SUGEN, Inc., -80°C.

Batch #917 p88, concentration 0.18 mg/mL

7. TBS-W Buffer

Reagent	M.W.	1xWorking Concentration	Amount per L
Tris	121.14	50 mM	6.057 g
NaCl	58.44	150 mM	8.766 g
Tween-20		0.05%	0.5 ml

To make 1 liter of a 1x working solution:

- 1) To a 1 liter graduated cylinder add ~900ml dH₂O
- 2) Add all reagents
- 3) When all reagents have dissolved, pH to 7.6 with HCl
- 4) Bring volume to 1 liter dH₂O
- 5) Do not keep a 10% stock solution of Tween20. Add 100%

Tween20 to the buffer.

A 10x stock solution can be made by multiplying the amounts by 10 (but keeping the final volume of 1 liter). This stock is then diluted 10 fold with dH₂O and re-pHed to 7.2.

1X and 10X TBS (pH 7.6) is supplied by the Media Preparation Department.

8. MilliQue H₂O + 4% DMSO

Reagent	M.W.	1xWorking Concentration	Amount per 1 L
MilliQue H ₂ O	NA	NA	960 mL
DMSO	NA	4%	40 ml

9. 1 mM ATP

Reagent	Vendor	Catalog #	Order quantity	M.W.	Amount for 5 ml
Adenosine-5'-triphosphate (from Equine muscle)	Sigma	A-5394	5g	551.1	2.75 mg
dH ₂ O					5 ml

To make 1 mM Stock solution:

- 1) Add 5 ml of dH₂O to 2.75 mg ATP
- 2) Vortex

5 *Note any mg amount of ATP can be used provided it is kept in the same ATP to dH₂O ratio.

 *Note this reagent can be stored at -20 in small aliquots to be taken out just prior to use and kept on ice. Do not freeze/thaw aliquots; discard any unused portion.

10. 1M MnCl₂

10

Reagent	M.W.	Amount per 100 ml	Stock Concentration
MnCl ₂	197.91	19.79 g	1M

12. Reduced Form Glutathione

Reagent	Vendor	Catalog #	Order quantity	M.W.	Amount for 1 ml	Stock Concentration
Reduced Form Glutathione	Sigma	G-4251	25 g	307.3	30.73	100 mM

15

- **Not Stable in resuspension. Prepare fresh stock for every experiment!**

13. 2X Kinase Dilution Buffer

Reagent	Stock solution	Amount per 100 ml	Working Concentration
dH ₂ O	NA	88.4 mL	
Hepes pH7.5	1 M	10 ml	100 mM
BSA/PBS	5%	0.4 mL	0.02%
Na - orthovanadate	0.1 M	0.2 mL	0.2 mM
NaCl	5M	3 mL	150 mM
Reduced Form Glutathione	100 mM	1 mL	1 mM

14. 4X ATP Reaction Mixture

Reagent	Stock solution	Amount per 10 ml	Working Concentration
dH ₂ O	NA	9.56 mL	
MnCl ₂	1 M	0.4 mL	40 mM
ATP	0.01 M	0.008 mL	0.8 μM

15. EDTA

Reagent	M.W.	Stock solution	Amount per 100ml	Working Solution
ethylenediamine-tetraacetic acid	292.25	500 mM	14.12 g	500mM

To make stock solution:

- 1) Add ~70 ml dH₂O to a 250ml beaker
- 2) Add EDTA
- 3) With pH probe in beaker, add 10N NaOH dropwise EDTA

will not dissolve until pH is around 7.0 as EDTA dissolves the pH will fall, add

more NaOH

- 4) When all EDTA is dissolved, adjust the pH to 8.0
- 5) Transfer to 100 ml graduated cylinder, bring volume to 100

ml with dH₂O.

Stock Solution also available from the Media Preparation

5 Department.

16. Antibody Dilution Buffer

Reagent	Stock solution	Amount per 100 ml	Working Concentration
TBS		88	NA
BSA/PBS	5%	1 ml	0.05%
Tween-20	10%	1 mL	0.1 %

10 17. HRP-Conjugated Anti-Ptyr

Reagent	Vendor	Catalog #	Order quantity
HRP-Conjugated Anti-Ptyr (PY99)	Sant Cruz Biotechnology	SC-7020 (lot G219)	100 µL

15 18. Stable ABTS Solution

Reagent	Vendor	Catalog #	Order quantity
2,2'-azino-bis(3-ethylbenzthiazoline-6-sulfonic acid) (ABTS)	Moss	ABTS-2000	20 L

19. NUNC 96-well V bottom polypropylene plates Applied Scientific
Catalog # AS-72092.

20. 10% SDS (from in House Stock in Common Reagent Room).

5

Procedure

1. Coat ELISA plates with 2 µg Poly(Glu-Tyr) in 100 µL PBS overnight
at 4 degrees (PolyEY stock 10.0 mg/ml in PBS at -80).

10 2. Wash using the PBSRINSE program of the Titertek Plate Washer.
Block plate with 150 µL of TBB for 60 min. The PBS/BLOCK program will add
blocking buffer after PBS rinse.

3. Wash plate twice with PBS, once with 50 mM Hepes buffer pH 7.4
(use washer program 1STWASH). Plates can sit in HEPES buffer if other buffers
15 need to be prepared. The Kinase Dilution Buffer, ATP mixture, and TBS-W should
all be prepared before proceeding.

4. Add 25 µL drugs (in 4% DMSO) or DMSO controls (4% in water) to
plate.

If starting with a 10 mM stock of compound in 100% DMSO, make a 25 fold
20 dilution into water. The concentration of the drug dilution plate will be 400 µM in
4% DMSO. If creating serial dilutions, dilute in 4% DMSO/ water down the drug
plate. Final compound concentration (highest) in the assay plate will be 100 µM in
1% DMSO

25 5. Add 50 µl of diluted kinase into all wells. Purified kinase is diluted
into "Kinase Dilution Buffer" to achieve a concentration of 5 ng/well.
(Concentration of present purified kinase batch is 0.18 mg/mL). For the current
batch, add 3.33 µL to 6 mLs of KDB.

30 • **Kinase is unstable when diluted. Start kinase reaction as soon
after dilution as possible.**

6. Add 25 μ L of 0.5 M EDTA to the negative controls wells.
7. Add 25 μ L ATP/ $MnCl_2$ mixture to entire plate.

5 **The reaction time is 10 minutes and is the most critical part of the assay. The assay must be stopped with 25 μ L 0.5 M EDTA in a manner similar to the addition of ATP.**

- 10 • Additions of the enzyme, ATP, primary and secondary antibodies can be added with the Titertek Multidrop. The tubing must be flushed with MilliQue H_2O and primed with reagent. Dispensing volumes are accurate to 20 μ L.

- 15 9. Wash plate 4x with TBS-W (Washer program WASH1).

10. Detect substrate phosphorylation HRP conjugated anti-Ptyr diluted 1:6,000 in Antibody Dilution Buffer. Add 100 μ L per well and incubate at room temperature, with shaking, for one hour.

- 20 11. Wash plate 3X with TBS-W and 1X with PBS (Washer Program LASTWASH). Residual Tween20 from the washing buffer can inhibit HRP activity and decrease the delta.

- 25 12. Add 100 μ l of ABTS solution to each well using the Proline Biohit Repeating Pipetor or Titertek Multidrop .

15. If necessary, stop the development reaction with the addition of 20 μ l 10% SDS per well.

- 30 16. Read plate on Dynatech MR7000 elisa reader.
Test Filter: 410 nM

Reference Filter: 630 nM

Plate template for Placement of Controls

										0	1	2
OS												EG
OS												EG
OS												EG
OS												EG
EG												OS
EG												OS
EG												OS
EG												OS

5

Cellular/Biologic Assays

EXAMPLE 120: GENERAL PROCEDURE FOR BRDU INCORPORATION ASSAYS

10 The following assays use cells engineered to express a receptor of interest and the evaluate the effect of a compound of interest on the activity of ligand-induced DNA synthesis by determining BrdU incorporation into the DNA.

The following materials, reagents, and procedure are general to each of the following BrdU incorporation assays. Variances in specific assays are noted.

15 Materials and Reagents:

1. The appropriate ligand.
2. The appropriate engineered cells.

3. BrdU Labeling Reagent: 10 mM, in PBS (pH7.4) (Boehringer Mannheim, Germany).
4. FixDenat: fixation solution (ready to use) (Boehringer Mannheim, Germany).
5. Anti-BrdU-POD: mouse monoclonal antibody conjugated with peroxidase (Boehringer Mannheim, Germany).
6. TMB Substrate Solution: tetramethylbenzidine (TMB, Boehringer Mannheim, Germany).
7. PBS Washing Solution : 1X PBS, pH 7.4.
8. Albumin, Bovine (BSA), fraction V powder (Sigma Chemical Co., USA).

General Procedure:

1. Cells are seeded at 8000 cells/well in 10% CS, 2mM Gln in DMEM, in a 96 well plate. Cells are incubated overnight at 37 °C in 5% CO₂.
2. After 24 hours, the cells are washed with PBS, and then are serum starved in serum free medium (0%CS DMEM with 0.1% BSA) for 24 hours.
3. On day 3, the appropriate ligand and the test compound are added to the cells simultaneously. The negative control wells receive serum free DMEM with 0.1% BSA only; the positive control cells receive the ligand but no test compound. Test compounds are prepared in serum free DMEM with ligand in a 96 well plate, and serially diluted for 7 test concentrations.
4. After 18 hours of ligand activation, diluted BrdU labeling reagent (1:100 in DMEM, 0.1% BSA) is added and the cells are incubated with BrdU (final concentration=10 µM) for 1.5 hours.
5. After incubation with labeling reagent, the medium is removed by decanting and tapping the inverted plate on a paper towel. FixDenat solution is added (50 µL/well) and the plates are incubated at room temperature for 45 minutes on a plate shaker.
6. The FixDenat solution is thoroughly removed by decanting and tapping the inverted plate on a paper towel. Milk is added (5% dehydrated milk in

PBS, 200 μ L/well) as a blocking solution and the plate is incubated for 30 minutes at room temperature on a plate shaker.

7. The blocking solution is removed by decanting and the wells are washed once with PBS. Anti-BrdU-POD solution (1:200 dilution in PBS, 1% BSA) is added (50 μ L/well) and the plate is incubated for 90 minutes at room temperature on a plate shaker.

8. The antibody conjugate is thoroughly removed by decanting and rinsing the wells 5 times with PBS, and the plate is dried by inverting and tapping on a paper towel.

9. TMB substrate solution is added (100 μ L/well) and incubated for 20 minutes at room temperature on a plate shaker until color development is sufficient for photometric detection.

10. The absorbance of the samples are measured at 410 nm (in "dual wavelength" mode with a filter reading at 490 nm, as a reference wavelength) on a Dynatech ELISA plate reader.

EXAMPLE 121: EGF-Induced BrdU Incorporation Assay

The procedure is the same as the general procedure, outlined above, except for the following changes:

Materials and Reagents:

1. Mouse EGF, 201 (Toyobo Co., Ltd., Japan).
2. 3T3/EGFRc7.

EXAMPLE 122: EGF-Induced Her-2-driven BrdU Incorporation Assay

The procedure is the same as the general procedure, outlined above, except for the following changes:

Materials and Reagents:

1. Mouse EGF, 201 (Toyobo Co., Ltd., Japan).
2. 3T3/EGFr/Her2/EGFr (EGFr with a Her-2 kinase domain).

EXAMPLE 123: EGF-Induced Her-4-driven BrdU Incorporation Assay

The procedure is the same as the general procedure, outlined above, except for the following changes:

5

Materials and Reagents:

1. Mouse EGF, 201 (Toyobo Co., Ltd., Japan).
2. 3T3/EGFr/Her4/EGFr (EGFr with a Her-4 kinase domain).

10 EXAMPLE 124: PDGF-Induced BrdU Incorporation Assay

The procedure is the same as the general procedure, outlined above, except for the following changes:

Materials and Reagents:

15

1. Human PDGF B/B (Boehringer Mannheim, Germany).
2. 3T3/EGFRc7.

EXAMPLE 125: FGF-Induced BrdU Incorporation Assay

The procedure is the same as the general procedure, outlined above, except for the following changes:

20

Materials and Reagents:

1. Human FGF2/bFGF (Gibco BRL, USA).
2. 3T3c7/EGFr

25

EXAMPLE 126: IGF1-Induced BrdU Incorporation Assay

The procedure is the same as the general procedure, outlined above, except for the following changes:

30

Materials and Reagents:

1. Human, recombinant (G511, Promega Corp., USA)

2. 3T3/IGF1r.

EXAMPLE 127: Insulin-Induced BrdU Incorporation Assay

The procedure is the same as the general procedure, outlined above, except for the following changes:

5

Materials and Reagents:

1. Insulin, crystalline, bovine, Zinc (13007, Gibco BRL, USA).

EXAMPLE 128: HGF-Induced BrdU Incorporation Assay

10 The procedure is the same as the general procedure, outlined above, except for the following changes:

Materials and Reagents:

1. Recombinant human HGF (Cat. No. 249-HG, R&D Systems, Inc.
15 USA).
2. BxPC-3 cells (ATCC CRL-1687).

Procedure:

1. Cells are seeded at 9000 cells/well in RPMI 10% FBS in a 96 well
20 plate. Cells are incubated overnight at 37°C in 5% CO₂.
2. After 24 hours, the cells are washed with PBS, and then are serum
starved in 100 µl serum-free medium (RPMI with 0.1% BSA) for 24 hours.
3. On day 3, 25 µl containing ligand (prepared at 1 µg/ml in RPMI with
0.1% BSA; final HGF conc. = 200 ng/ml) and test compounds are added to the cells.
25 The negative control wells receive 25 µl serum-free RPMI with 0.1% BSA only; the
positive control cells receive the ligand (HGF) but no test compound. Test
compounds are prepared at 5 times their final concentration in serum-free RPMI
with ligand in a 96 well plate, and serially diluted for 7 test concentrations.
Typically, the highest final concentration of test compound is 100 µM, and 1:3
30 dilutions are used (i.e. final test compound concentration range = 0.137-100 µM).

4. After 18 hours of ligand activation, 12.5 µl of diluted BrdU labeling reagent (1:100 in RPMI, 0.1% BSA) is added to each well and the cells are incubated with BrdU (final concentration = 10 µM) for 1 hour.

5. Same as General Procedure.

5 6. Same as General Procedure.

7. The blocking solution is removed by decanting and the wells are washed once with PBS. Anti-BrdU-POD solution (1:100 dilution in PBS, 1% BSA) is added (100 µl/well) and the plate is incubated for 90 minutes at room temperature on a plate shaker.

10 8. Same as General Procedure.

9. Same as General Procedure.

10. Same as General Procedure.

EXAMPLE 129: HUV-EC-C Assay

15 The following protocol may also be used to measure a compound's activity against PDGF-R, FGF-R, VEGF, aFGF or Flk-1/KDR, all of which are naturally expressed by HUV-EC cells.

DAY 0

20 1. Wash and trypsinize HUV-EC-C cells (human umbilical vein endothelial cells, (American Type Culture Collection; catalogue no. 1730 CRL). Wash with Dulbecco's phosphate-buffered saline (D-PBS; obtained from Gibco BRL; catalogue no. 14190-029) 2 times at about 1 mL/10 cm² of tissue culture flask. Trypsinize with 0.05% trypsin-EDTA in non-enzymatic cell dissociation solution
25 (Sigma Chemical Company; catalogue no. C-1544). The 0.05% trypsin was made by diluting 0.25% trypsin/1 mM EDTA (Gibco; catalogue no. 25200-049) in the cell dissociation solution. Trypsinize with about 1 mL/25-30 cm² of tissue culture flask for about 5 minutes at 37 °C. After cells have detached from the flask, add an equal volume of assay medium and transfer to a 50 mL sterile centrifuge tube (Fisher
30 Scientific; catalogue no. 05-539-6).

2. Wash the cells with about 35 mL assay medium in the 50 mL sterile centrifuge tube by adding the assay medium, centrifuge for 10 minutes at approximately 200 g, aspirate the supernatant, and resuspend with 35 mL D-PBS. Repeat the wash two more times with D-PBS, resuspend the cells in about 1 mL assay medium/15 cm² of tissue culture flask. Assay medium consists of F12K medium (Gibco BRL; catalogue no. 21127-014) + 0.5% heat-inactivated fetal bovine serum. Count the cells with a Coulter Counter (Coulter Electronics, Inc.) and add assay medium to the cells to obtain a concentration of 0.8-1.0x10⁵ cells/mL.
3. Add cells to 96-well flat-bottom plates at 100 µL/well or 0.8-1.0x10⁴ cells/well; incubate ~24h at 37 °C, 5% CO₂.

DAY 1

1. Make up two-fold drug titrations in separate 96-well plates, generally 50 µM on down to 0 µM. Use the same assay medium as mentioned in day 0, step 2 above. Titrations are made by adding 90 µL/well of drug at 200 µM (4X the final well concentration) to the top well of a particular plate column. Since the stock drug concentration is usually 20 mM in DMSO, the 200 µM drug concentration contains 2% DMSO.

Therefore, diluent made up to 2% DMSO in assay medium (F12K + 0.5% fetal bovine serum) is used as diluent for the drug titrations in order to dilute the drug but keep the DMSO concentration constant. Add this diluent to the remaining wells in the column at 60 µL/well. Take 60 µL from the 120 µL of 200 µM drug dilution in the top well of the column and mix with the 60 µL in the second well of the column. Take 60 µL from this well and mix with the 60 µL in the third well of the column, and so on until two-fold titrations are completed. When the next-to-the-last well is mixed, take 60 µL of the 120 µL in this well and discard it. Leave the last well with 60 µL of DMSO/media diluent as a non-drug-containing control. Make 9 columns of titrated drug, enough for triplicate wells each for 1) VEGF (obtained from Pepro Tech Inc., catalogue no. 100-200, 2) endothelial cell growth factor (ECGF) (also known as acidic fibroblast growth factor, or aFGF) (obtained from Boehringer Mannheim Biochemica, catalogue no. 1439 600); or, 3) human

PDGF B/B (1276-956, Boehringer Mannheim, Germany) and assay media control. ECGF comes as a preparation with sodium heparin.

2. Transfer 50 μL /well of the drug dilutions to the 96-well assay plates containing the $0.8\text{--}1.0 \times 10^4$ cells/100 μL /well of the HUV-EC-C cells from day 0 and incubate ~ 2 h at 37°C , 5% CO_2 .

3. In triplicate, add 50 μL /well of 80 $\mu\text{g/mL}$ VEGF, 20 ng/mL ECGF, or media control to each drug condition. As with the drugs, the growth factor concentrations are 4X the desired final concentration. Use the assay media from day 0 step 2 to make the concentrations of growth factors. Incubate approximately 24 hours at 37°C , 5% CO_2 . Each well will have 50 μL drug dilution, 50 μL growth factor or media, and 100 μL cells, = 200 μL /well total. Thus the 4X concentrations of drugs and growth factors become 1X once everything has been added to the wells.

DAY 2

1. Add ^3H -thymidine (Amersham; catalogue no. TRK-686) at 1 μCi /well (10 μL /well of 100 $\mu\text{Ci/mL}$ solution made up in RPMI media + 10% heat-inactivated fetal bovine serum) and incubate ~ 24 h at 37°C , 5% CO_2 . Note: ^3H -thymidine is made up in RPMI media because all of the other applications for which we use the ^3H -thymidine involve experiments done in RPMI. The media difference at this step is probably not significant. RPMI was obtained from Gibco BRL, catalogue no. 11875-051.

DAY 3

1. Freeze plates overnight at -20°C .

DAY 4

1. Thaw plates and harvest with a 96-well plate harvester (Tomtec Harvester 96^(R)) onto filter mats (Wallac; catalogue no. 1205-401); read counts on a Wallac Betaplate^(TM) liquid scintillation counter.

In Vivo Animal Models

EXAMPLE 130: Xenograft Animal Models

5 The ability of human tumors to grow as xenografts in athymic mice
(*e.g.*, Balb/c, nu/nu) provides a useful in vivo model for studying the biological
response to therapies for human tumors. Since the first successful
xenotransplantation of human tumors into athymic mice, (Rygaard and Povlsen,
1969, Acta Pathol. Microbial. Scand. 77:758-760), many different human tumor cell
lines (*e.g.*, mammary, lung, genitourinary, gastro-intestinal, head and neck,
10 glioblastoma, bone, and malignant melanomas) have been transplanted and
successfully grown in nude mice. The following assays may be used to determine
the level of activity, specificity and effect of the different compounds of the present
invention.

Suitable cell lines for subcutaneous xenograft experiments include C6 cells
15 (glioma, ATCC # CCL 107), A375 cells (melanoma, ATCC # CRL 1619), A431
cells (epidermoid carcinoma, ATCC # CRL 1555), Calu 6 cells (lung, ATCC # HTB
56), PC3 cells (prostate, ATCC # CRL 1435) and NIH 3T3 fibroblasts genetically
engineered to overexpress EGFR, PDGFR, IGF-1R or any other test kinase. The
following protocol can be used to perform xenograft experiments:

20 Female athymic mice (BALB/c, nu/nu) are obtained from Simonsen
Laboratories (Gilroy, CA). All animals are maintained under clean-room conditions
in Micro-isolator cages with Alpha-dri bedding. They receive sterile rodent chow
and water ad libitum.

Cell lines are grown in appropriate medium (for example, MEM,
25 DMEM, Ham's F10, or Ham's F12 plus 5% - 10% fetal bovine serum (FBS) and 2
mM glutamine (GLN)). All cell culture media, glutamine, and fetal bovine serum
are purchased from Gibco Life Technologies (Grand Island, NY) unless otherwise
specified. All cells are grown in a humid atmosphere of 90-95% air and 5-10%
CO₂ at 37 °C. All cell lines are routinely subcultured twice a week and are
30 negative for mycoplasma as determined by the Mycotect method (Gibco).

Trypsin-EDTA and pelleted at 450 x g for 10 min. Pellets are resuspended in sterile PBS or media (without FBS) to a particular concentration and the cells are implanted into the hindflank of the mice (8 - 10 mice per group, $2 - 10 \times 10^6$ cells/animal). Tumor growth is measured over 3 to 6 weeks using venier calipers. Tumor volumes are calculated as a product of length x width x height unless otherwise indicated. P values are calculated using the Students t-test. Test compounds in 50 - 100 μ L excipient (*e.g.*, DMSO, or VPD:D5W) are delivered by IP injection at different concentrations generally starting at day one after implantation.

The following tumor invasion model has been developed and may be used for the evaluation of therapeutic value and efficacy of the compounds identified to selectively inhibit KDR/FLK-1 receptor or CDK2.

8 week old nude mice (female) (Simonsen Inc.) were used as experimental animals. Implantation of tumor cells was performed in a laminar flow hood. For anesthesia, Xylazine/Ketamine Cocktail (100 mg/kg ketamine and 5 mg/kg Xylazine) are administered intraperitoneally. A midline incision is done to expose the abdominal cavity (approximately 1.5 cm in length) to inject 10^7 tumor cells in a volume of 100 μ L medium. The cells are injected either into the duodenal lobe of the pancreas or under the serosa of the colon. The peritoneum and muscles are closed with a 6-0 silk continuous suture and the skin was closed by using wound clips. Animals were observed daily.

After 2-6 weeks, depending on gross observations of the animals, the mice are sacrificed, and the local tumor metastases, to various organs (lung, liver,

brain, stomach, spleen, heart, muscle) are excised and analyzed (measurements of tumor size, grade of invasion, immunochemistry, and in situ hybridization).

EXAMPLE 134: Measurement of Cell Toxicity

- 5 Therapeutic compounds should be more potent in inhibiting protein kinase activity than in exerting a cytotoxic effect. A measure of the effectiveness and cell toxicity of a compound can be obtained by determining the therapeutic index: IC_{50}/LD_{50} . IC_{50} , the dose required to achieve 50% inhibition, can be measured using standard techniques such as those described herein. LD_{50} , the dosage which
- 10 results in 50% toxicity, can also be measured by standard techniques (Mossman, 1983, J. Immunol. Methods, 65:55-63), by measuring the amount of LDH released (Korzeniewski and Callewaert, 1983, J. Immunol. Methods, 64:313; Decker and Lohmann-Matthes, 1988, J. Immunol. Methods, 115:61), or by measuring the lethal dose in animal models. Compounds with a large therapeutic index are preferred.
- 15 The therapeutic index should be greater than 2, preferably at least 10, more preferably at least 50.

EXAMPLE 135: The Activity of the Compounds of the Invention

- 20 The biological or biochemical activity of some of the compounds of the invention were tested using the assays described above. The IC_{50} values were measured for several of the compounds of the invention. The results are shown in Table 10 below.

Table 10

Compound No.	bio flkGST IC_{50} (μ M)	bio EGF IC_{50} (μ M)	bio PDGF IC_{50} (μ M)
IN-001	0.86	>100	0.11
IN-002	0.03	2.87	0.38
IN-003	1.56	61.29	
IN-004	0.47	2.57	21
IN-005	0.31	16.27	
IN-006	0.16	4.69	1.45
IN-007	0.4	37.37	<0.78
IN-008	0.22	1.97	0.29
IN-009	9.42	>100	<0.78

Compound No.	bio flkGST IC ₅₀ (μM)	bio EGF IC ₅₀ (μM)	bio PDGF IC ₅₀ (μM)
IN-010	0.35	88.8	<0.78
IN-011	0.52	1.36	<0.78
IN-012	1.95	>100	<0.78
IN-013	0.1	41.26	0.008
IN-014	0.003	43.47	0.15
IN-015	0.99	70.62	0
IN-016	0.05	>10	0.028
IN-017	0.01	4.67	0.07
IN-018	5.41	>100	<0.78
IN-019	0.16	9.55	5.65
IN-022	0.14	>20	
IN-023	0.24	>20	
IN-024	<0.01	>20	
IN-025	0.14	>20	
IN-026	0.76	>20	
IN-027	2.01	>20	
IN-028	0.66	6.06	
IN-029	0.76	>20	
IN-030	<0.009	>20	
IN-031	0.53	>20	
IN-032	1.47	>20	
IN-033	<0.009	8.2	
IN-034	13.98	>20	
IN-035	>20	>20	
IN-036	>20	>20	
IN-037	4.38	>20	
IN-038	>10	>20	
IN-039	0.17	>100	0.91
IN-040	0.21	>100	9.3
IN-041	0.21	8.43	
IN-042	1.34	7.84	
IN-043	0.27	4.35	
IN-044	0.18	1.69	
IN-045	0.12	8.02	
IN-046	1.05	1.23	
IN-047	0.39	0.87	
IN-048	1.16	2.13	
IN-049	0.13	4.38	
IN-050	0.71	2.4	
IN-051	0.37	3.52	
IN-052	1.02	1.16	
IN-053	1.21	15.2	
IN-054	4.28	>20	
IN-055	1.5	3.5	

Compound No.	bio flkGST IC ₅₀ (μM)	bio EGF IC ₅₀ (μM)	bio PDGF IC ₅₀ (μM)
IN-056	9.96	15.53	
IN-057	>20	>20	
IN-058	8.49	>20	
IN-059	0.63	>20	
IN-060	8.55	>20	
IN-061	0.39	4.46	
IN-062	15.38	>20	
IN-063	14.33	>20	
IN-064	17.82	>20	
IN-065	11.67		89.41
IN-066	>20	>100	>100
IN-067	3.99	>100	>100
IN-068	>20	>100	>100
IN-069	12.77	>100	>100
IN-070	2.33	>100	79.22
IN-071	11.47	22.04	69.14
IN-072	>20	>100	>100
IN-073	>20	>100	>100
IN-074	1.58	>100	>100
IN-075	6.51	>100	>100

CONCLUSION

One skilled in the art would readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. The molecular complexes and the methods, procedures, treatments, molecules, specific compounds described herein are presently representative of preferred embodiments are exemplary and are not intended as limitations on the scope of the invention. Changes therein and other uses will occur to those skilled in the art which are encompassed within the spirit of the invention are defined by the scope of the claims.

It will be readily apparent to one skilled in the art that varying substitutions and modifications may be made to the invention disclosed herein without departing from the scope and spirit of the invention.

All patents and publications mentioned in the specification are indicative of the levels of those skilled in the art to which the invention pertains. All patents and publications are herein incorporated by reference to the same extent as if each

individual publication was specifically and individually indicated to be incorporated by reference.

5 The invention illustratively described herein suitably may be practiced in the absence of any element or elements, limitation or limitations which is not specifically disclosed herein. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention that in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments and optional features, modification and variation of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention as defined by the appended claims.

15 In addition, where features or aspects of the invention are described in terms of Markush groups, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group. For example, if X is described as selected from the group consisting of bromine, chlorine, and iodine, claims for X being bromine and claims for X being bromine and chlorine are fully described.

20 The invention has been described broadly and generically herein. Each of the narrower species and subgeneric groupings falling within the generic disclosure also form part of the invention. This includes the generic description of the invention with a proviso or negative limitation removing any subject matter from the genus, regardless of whether or not the excised material is specifically recited herein.